

North Dakota 2008 Integrated Section 305(b) Water Quality Assessment Report and Section 303(d) List of Waters Needing Total Maximum Daily Loads



The S.S. Ruby charter boat tours the Red River of the North in the Fargo, ND and Moorhead, MN area. The S.S. Ruby is owned and operated by River Keepers, a non-profit organization established in 1990 to protect and preserve the integrity and natural environment of the Red River of the North in the F-M area.

**Submitted to the US EPA
August 4, 2008**

**Approved
September 29, 2008**





UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
REGION 8

1595 Wynkoop Street
Denver, CO 80202-1129
Phone 800-227-8917
<http://www.epa.gov/region08>

Ref: 8EPR-EP

SEP 29 2008

Mr. Dennis Fewless
Director
Division of Water Quality
North Dakota Department of Health
918 East Divide Ave., 4th Floor
Bismarck, ND 58501-1947

Re: Clean Water Act Section 303(d) Total
Maximum Daily Load (TMDL) Waterbody
List

Dennis

Dear Mr. Fewless:

Thank you for your submittal of North Dakota's 2008 Integrated Report for Surface Water Quality Assessment dated August 4, 2008. EPA has conducted a complete review of the Clean Water Act Section 303(d) waterbody list and supporting documentation and information. Based on this review EPA has determined that North Dakota's 2008 list of water quality limited segments (WQLSs) still requiring TMDLs meets the requirements of Section 303(d) of the Clean Water Act (CWA) and EPA's implementing regulations. Therefore, by this order, EPA hereby APPROVES North Dakota's 2008 Section 303(d) list. Please see the enclosure for a description of the statutory and regulatory requirements and a summary of EPA's review of North Dakota's compliance with each requirement.

EPA's approval of North Dakota's 2008 Section 303(d) list extends to all waterbodies in category 5 of the list with the exception of those waters that are within Indian Country, as defined in 18 U.S.C. Section 1151. EPA is taking no action to approve or disapprove the State's list with respect to those waters at this time. EPA, or eligible Indian Tribes, as appropriate, will retain responsibilities under Section 303(d) for those waters.

The public participation process sponsored by the North Dakota Department of Health included publishing ads in newspapers across the state requesting public input in developing the draft list and requesting water quality data, an official public notice on the list availability, use of the North Dakota Department of Health website, and mail or email requests to many entities asking for both comments and additional data or information on waters. We commend the State for its thorough public participation process.



Under current regulations, the next Section 303(d) list is required to be submitted on April 1, 2010. Although current regulations require lists to be submitted every 2 years, in April of even years, states may submit Section 303(d) lists more frequently as they deem necessary. All additions, deletions and modifications to the list will require EPA approval.

Again, thank you for the efforts related to the development of North Dakota's Section 303(d) waterbody list for the 2008-2010 biennium. If you have questions, the most knowledgeable person on my staff is Kris Jensen and she may be reached at (303) 312-6237.

Sincerely,



Carol L. Campbell
Assistant Regional Administrator
Office of Ecosystems Protection and
Remediation

Enclosure

cc: Mike Ell, NDDH
Karen Hamilton, Vern Berry, Karl Hermann, Eric Steinhaus, EPA

**North Dakota 2008 Integrated
Section 305(b) Water Quality Assessment Report and
Section 303(d) List of Waters Needing
Total Maximum Daily Loads**

John Hoeven, Governor
Terry Dwelle, M.D., State Health Officer



North Dakota Department of Health
Division of Water Quality
Gold Seal Center, 4th Floor
918 East Divide Ave.
Bismarck, North Dakota 58501-1947

701.328.5210

CONTENTS

PART I. EXECUTIVE SUMMARY	I-1
PART II. INTRODUCTION	II-1
A. Section 305(b) Water Quality Assessment Report	II-1
B. Section 303(d) TMDL List of Water Quality-limited Waters.....	II-1
PART III. BACKGROUND.....	III-1
A. Atlas	III-1
B. Total Waters	III-2
C. Water Pollution Control Program	III-4
Chapter 1. Water Quality Standards Program.....	III-4
Chapter 2. Point Source Control Program	III-6
Chapter 3. Nonpoint Source (NPS) Pollution Management Program	III-12
Resource Assessment.....	III-17
Prioritization	III-18
Assistance	III-18
Coordination	III-19
Information and Education	III-20
Program Evaluation	III-21
Chapter 4. Total Maximum Daily Load (TMDL) Program.....	III-28
Chapter 5. Coordination with Other Agencies.....	III-30
D. Cost/Benefit Assessment.....	III-31
E. Special State Concerns and Recommendations.....	III-32
PART IV. SURFACE WATER MONITORING AND ASSESSMENT	
METHODOLOGY	IV-1
A. Surface Water Quality Monitoring Program.....	IV-1
Chapter 1. Monitoring Goals and Objectives	IV-1
Chapter 2. Monitoring Program, Projects and Studies	IV-2
Ambient Water Quality Monitoring Network for Rivers and Streams.....	IV-2
Biological Monitoring Program.....	IV-6
Lake Water Quality Assessment Program	IV-9
Fish Tissue Contaminant Surveillance Program.....	IV-12
NPS Pollution Management Program Monitoring.....	IV-12
Support Projects and Special Studies.....	IV-14
Complaint and Fish Kill Investigations	IV-14
Stream Flow	IV-15

B. Assessment Methodology	IV-16
Chapter 1. Introduction	IV-16
Chapter 2. Assessment Database (ADB)	IV-16
Chapter 3. Beneficial Use Designation	IV-17
Chapter 4. Sufficient and Credible Data Requirements	IV-17
Chapter 5. Existing and Available Water Quality Data	IV-18
Rivers and Streams	IV-18
Lake and Reservoirs	IV-20
Fish Tissue Data	IV-21
Other Agency/Organization Data	IV-22
Chapter 6. Beneficial Use Assessment Methodology	IV-22
Chapter 7. Assessment Categories	IV-22
PART V. SECTION 305(b) WATER QUALITY ASSESSMENT	V-1
A. Rivers and Streams Water Quality Assessment	V-1
Chapter 1. Assessment Category Summary	V-1
Chapter 2. Water Quality Summary	V-3
Chapter 3. Environmental Monitoring and Assessment Program	
Western Pilot Project Results for North Dakota	V-7
Introduction	V-7
Study Area Description	V-7
Sample Site Determination	V-8
Extent of Resource Assessed	V-9
Reference Site Determination	V-10
Condition Indicators	V-11
Stressor Indicators	V-18
B. Lakes and Reservoirs Water Quality Assessment	V-26
Chapter 1. Assessment Category Summary	V-26
Chapter 2. Water Quality Summary	V-27
Chapter 3. Trophic Status	V-31
Chapter 4. Control Methods	V-32
Chapter 5. Restoration/Rehabilitation Efforts	V-32
Chapter 6. Acid Effects on Lakes and Reservoirs	V-32
Chapter 7. Toxic Effects on Lakes and Reservoirs	V-33
C. Wetlands Assessment Program	V-34
Chapter 1. Background	V-34
Chapter 2. Extent of Wetland Resources	V-35
Chapter 3. Integrity of Wetland Resources	V-37
Chapter 4. Wetland Water Quality Standards	V-39
Chapter 5. Wetland Monitoring and Assessment Program	V-40
Current and Historic Program	V-40
Regional Scale Wetland Assessment Pilot Project	V-40
Other Program Plans	V-40

D. Public Health/Aquatic Life Concerns	V-42
--	------

**PART VI. NORTH DAKOTA SECTION 303(d) LIST OF WATER
QUALITY-LIMITED WATERS NEEDING TMDLS..... V1-1**

A. Background	VI-1
B. Prioritization of TMDL-Listed Waters	VI-2
C. Public Participation Process	VI-3
D. Listing of Impaired Waters Needing TMDLs	VI-4
E. De-listing of 2006-Listed TMDL Waters.....	VI-4
F. TMDL Development and Monitoring Schedule.....	VI-5

PART VII. GROUND WATER ASSESSMENT VII-1

A. Ground Water Extent and Uses.....	VII-1
Chapter 1. Aquifer Description.....	VII-1
Chapter 2. Ground Water Use.....	VII-3
B. Ground Water Contamination Sources	VII-4
Chapter 1. Contaminant Source Description	VII-4
Chapter 2. Ground Water Contaminant Source Databases	VII-4
Concentrated Animal Feeding Operations (CAFO) Database.....	VII-4
Underground Injection Control (UIC) Program Class V Database	VII-5
Spill Response/Contaminant Release Database.....	VII-5
Ambient Ground Water Quality Monitoring Database.....	VII-5
C. Ground Water Protection Programs	VII-6
Chapter 1. Wellhead Protection and Source Water Protection Programs.....	VII-6
D. Ground Water Quality.....	VII-8
Chapter 1. Ambient Ground Water Monitoring Program.....	VII-8
Chapter 2. Underground Injection Control (UIC) Program.....	VII-8
Chapter 3. Additional Ground Water-Related Projects	VII-9
Facility Location Reviews	VII-9
Water Appropriation and Monitoring	VII-9
Contaminant Release Sites.....	VII-9
Pesticide Use Exemption Evaluations	VII-9
Emergency Response and Spills	VII-10

REFERENCES..... VIII-1

APPENDICES

Appendix A. Water Quality Assessment Methodology for North Dakota's Surface Waters	A-1
Appendix B. Agency and Organization Data Request.....	B-1
Appendix C. Estimated Weighted Average Methyl-Mercury Concentrations in Fish for Lake Sakakawea.....	C-1
Appendix D. Estimated Weighted Average Methyl-Mercury Concentrations in Fish for Lake Oahe and the Missouri River	D-1
Appendix E. Estimated Weighted Average Methyl-mercury Concentrations in Fish for Devils Lake	E-1
Appendix F. Estimated Weighted Average Methyl-mercury Concentrations in Fish for the Red River of the North.....	F-1
Appendix G. Public Notice Statement Requesting Public Comment on the State of North Dakota's Draft 2008 Section 303(d) List	G-1
Appendix H. Public Comments Received on the State of North Dakota's Draft 2008 Section 303(d) List and the State Responses.	H-1

TABLES

Table III-1. Atlas.....	III-1
Table III-2. Status and Budgets for Projects Supported Under the 2003, 2006 and 2007 Section 319 Grants (1/1/03-11/30/07).....	III-14
Table III-3. Section 319 Allocation and Expenditures per Project Category (1/1/03-11/30/07)	III-19
Table III-4. Agencies/Organizations Represented on the North Dakota NPS Pollution Task Force	III-20
Table III-5. BMPs Supported Under the 2003, 2006 and 2007 Grants (1/1/03-11/30/07)	III-22
Table III-6. STEPL Estimates – Nitrogen, Phosphorus and Sediment Load Reductions in 2006/2007	III-24
Table IV-1. Ambient Water Quality Monitoring Network Sites	IV-3
Table IV-2. Ambient Water Quality Monitoring Parameters	IV-6
Table IV-3. Assessment Categories for the Integrated Report	IV-23
Table IV-4. Biological Integrity Scoring and Aquatic Life Use Support Criteria Based on the Lake Agassiz Plain Ecoregion Fish IBI	IV-29
Table IV-5. Macroinvertebrate IBI Scores for Reference Sites in the Lake Agassiz Plain Ecoregion of North Dakota	IV-29
Table IV-6. Biological Integrity Scoring and Aquatic Life Use Support Criteria Based on the Lake Agassiz Plain Ecoregion Macroinvertebrate IBI.....	IV-29
Table IV-7. State Water Quality Standards for Chloride, Sulfate and Nitrate (NDDoH, 2001)	IV-31
Table V-1. Assessment Category Summary for Rivers and Streams in North Dakota (Miles).....	V-2
Table V-2. Individual Use Support Summary for Rivers and Streams in North Dakota (Miles).....	V-3
Table V-3. Impairment Summary for Rivers and Streams in North Dakota	V-4

Table V-4. Impairment Source Summary for Rivers and Streams in North Dakota	V-6
Table V-5. Landscape, Physical Habitat and Chemical Metrics Used in the North Dakota Reference Site Index	V-10
Table V-6. Multi-metric Index Threshold Values Used to Determine Condition Classes for Biological Indicators in the Cultivated Plains Region of North Dakota	V-12
Table V-7. Multi-metric Index Threshold Values Used to Determine Condition Classes for Biological Indicators in the Rangeland Plains Region of North Dakota.....	V-12
Table V-8. Threshold Values Used to Determine Condition Classes for Three Chemical Stressors in the Cultivated Plains Region of North Dakota.....	V-19
Table V-9. Threshold Values Used to Determine Condition Classes for Three Chemical Stressors in the Rangeland Plains Region of North Dakota	V-19
Table V-10. Threshold Values Used to Determine Condition Classes for the Mercury in Fish Tissue Stressor in North Dakota	V-19
Table V-11. Threshold Values Used to Determine Condition Classes for Four Physical Habitat Stressors in the Cultivated Plains Region of North Dakota	V-22
Table V-12. Threshold Values Used to Determine Condition Classes for Four Physical Habitat Stressors in the Rangeland Plains Region of North Dakota.....	V-22
Table V-13. Assessment Category Summary for Lakes and Reservoirs in North Dakota (Acres).....	V-26
Table V-14. Individual Use Support Summary for Lakes and Reservoirs in North Dakota (Acres).....	V-27
Table V-15. Impairment Summary for Lakes and Reservoirs in North Dakota.....	V-28
Table V-16. Impairment Source Summary for Lakes and Reservoirs in North Dakota.....	V-30
Table V-17. Trophic Status Summary for Lakes and Reservoirs in North Dakota	V-31
Table V-18. Definitions of Functions for Temporary and Seasonal Prairie Pothole Wetlands	V-38
Table VI-1. 2008 List of Section 303(d) TMDL Waters for the Souris River Basin in North Dakota	VI-6

Table VI-2. 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota	VI-9
Table VI-3. 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota	VI-29
Table VI-4. 2008 List of Section 303(d) TMDL Waters for the James River Basin in North Dakota	VI-41
Table VI-5. 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008	VI-46
Table VII-1. 2003 Reported Ground Water Use in North Dakota.....	VII-3

FIGURES

Figure III-1. Major Hydrologic Basins in North Dakota	III-3
Figure III-2. Average BOD-5 day and TSS Concentrations for Wastewater Discharges in North Dakota.....	III-7
Figure III-3. Cumulative Cost Category Expenditures Under the 2003, 2006 and 2007 Grants (1/1/03-11/30/07)	III-26
Figure III-4. BMP Category Expenditures Under the 2003, 2006 and 2007 Grants (1/1/03-11/30/07)	III-26
Figure III-5. Map Depicting Areas of Responsibility for Regional TMDL/Watershed Liaison Staff	III-29
Figure IV-1. Ambient Water Quality Monitoring Network Sites for Rivers and Streams	IV-5
Figure IV-2. Map of Reach-Indexed Assessment Units Delineated in the Souris River Basin	IV-18
Figure IV-3. Macroinvertebrate Sampling Sites in North Dakota (1995-2000).....	IV-22
Figure IV-4. A Graphic Representation of Carlson's TSI	IV-24
Figure V-1. Landcover Categories and Ecoregions in North Dakota.....	V-8
Figure V-2. North Dakota Stream Length Assessed.....	V-9
Figure V-3. EMAP Western Pilot Project Sampling Sites in North Dakota	V-10
Figure V-4. Condition Class Summary for Perennial Streams in North Dakota Based on Macroinvertebrate MMI Scores	V-13
Figure V-5. Condition Class Summary for Perennial Streams in the Cultivated Plains Region of North Dakota Based on Fish MMI Scores	V-14
Figure V-6. Perennial Stream Length (Kilometers) Estimates for Each Family of Fish in the Rangeland Plains Region of North Dakota	V-15
Figure V-7. Abundance of Primary Sportfish in the Rangeland Plains Region of North Dakota.....	V-16
Figure V-8. Percent Abundance of Each Family of Fish in the Rangeland Plains Region of North Dakota.....	V-16

Figure V-9. Condition Class Summary for Perennial Streams in North Dakota Based on Periphyton MMI Scores	V-18
Figure V-10. Perennial Stream Length (Kilometers) in the Cropland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Three Chemical Stressors	V-20
Figure V-11. Perennial Stream Length (Kilometers) in the Rangeland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Three Chemical Stressors	V-20
Figure V-12. Perennial Stream Length (Kilometers) in North Dakota Estimated to be in Good, Fair and Poor Condition Based on Three Chemical Stressors	V-21
Figure V-13. Perennial Stream Length (Kilometers) in North Dakota Estimated to be in Good and Poor Condition Based on the Mercury in Fish Tissue Stressor ..	V-22
Figure V-14. Perennial Stream Length (Kilometers) in the Cropland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Four Physical Habitat Stressors	V-23
Figure V-15. Perennial Stream Length (Kilometers) in the Rangeland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Four Physical Habitat Stressors	V-23
Figure V-16. Perennial Stream Length (Kilometers) in North Dakota Estimated to be in Good, Fair and Poor Condition Based on Four Physical Habitat Stressors	V-23
Figure V-17. Prairie Pothole Region	V-34
Figure VI-1. Graphical Depiction of 2008 List of Impaired Waters Needing TMDLs in the Souris River Basin	VI-8
Figure VI-2. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Upper Red River Basin	VI-27
Figure VI-3. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Lower Red River Basin.....	VI-28
Figure VI-4. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Lake Sakakawea/Missouri River Basin	VI-39
Figure VI-5. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Lake Oahe/Missouri River Basin.....	VI-40

Figure VI-6. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the James River Basin.....	VI-45
Figure VII-1. Major Glacial Drift Aquifers in North Dakota	VII-1
Figure VII-2. Location and Extent of North Dakota’s Primary Bedrock Aquifers	VII-2

PART I. EXECUTIVE SUMMARY

The Clean Water Act (CWA) contains several sections which require states to report on the quality of their waters. Section 305(b) (*State Water Quality Assessment Report*) requires a comprehensive biennial report; and Section 303(d) requires, from time to time, a list of a state's water quality-limited waters needing total maximum daily loads (TMDLs). The primary purpose of the Section 305(b) *State Water Quality Assessment Report* is to assess and report on the extent to which beneficial uses of the state's rivers, streams, lakes, reservoirs and wetlands are met. Section 305(b) of the Clean Water Act requires states to submit this assessment report every two years; the information presented in this report is for the reporting period of 2006-2007. The Section 305(b) report is a summary report that presents information on use impairment and the causes and sources of impaired or threatened uses for the state as a whole. While the Section 305(b) report is considered a summary report, Section 303 and its accompanying regulations (CFR Part 130 Section 7) require each state to list individual waterbodies (i.e., lakes, reservoirs, rivers, streams and wetlands) which are considered water quality limited and which require load allocations, waste load allocations and TMDLs. This list has become known as the "TMDL list" or "Section 303(d) list."

The North Dakota Department of Health (hereafter referred to as the department) currently recognizes 247 public lakes and reservoirs. Of the 247 public lakes and reservoirs recognized as public waters and included in the Assessment Database (ADB), only 198 are included in the state's water quality standards as classified lakes and therefore are assigned designated beneficial uses. The remaining 50 lakes and reservoirs, while included in the state's estimate of total lake acres, are not classified and therefore were not assessed for this report. Based on the state's Assessment Database (ADB), the 138 reservoirs have an areal surface of 543,156 acres. Reservoirs comprise about 71 percent of North Dakota's total lake/reservoir surface acres. Of these, 480,731 acres or 62 percent of the state's entire lake and reservoir acres are contained within the two mainstem Missouri River reservoirs (Lake Sakakawea and Lake Oahe). The remaining 136 reservoirs share 62,425 acres, with an average surface area of 459 acres. The 109 natural lakes in North Dakota cover 218,616 acres, with approximately 117,697 acres or 54 percent attributed to Devils Lake. The remaining 108 lakes average 934 acres, with half being smaller than 250 acres. There is an estimated 54,607 miles of rivers and streams in the state. Estimates of river stream miles in the state are based on the National Hydrography Dataset (NHD).

For purposes of 2008 Section 305(b) reporting and Section 303(d) listing, the U.S. Environmental Protection Agency (EPA) is encouraging states to submit an integrated report and to follow its integrated reporting guidance (EPA, 2005). Key to integrated reporting is an assessment of all of the state's waters and placement of those waters into one of five categories. The categories represent varying levels of water quality standards attainment, ranging from Category 1, where all of a waterbody's designated uses are met, to Category 5, where a pollutant impairs a waterbody and a TMDL is required.

Eighty-four percent (4,004 miles) of the rivers and streams assessed for this report fully support the beneficial use designated as aquatic life. Of the streams assessed as fully supporting aquatic life use, a little more than 60 percent (2,394 miles) are considered threatened. In other words, if water quality trends continue, the stream may not fully support its use for aquatic life in the

future. The remaining 16 percent of rivers and streams assessed for this report were assessed as not supporting aquatic life use.

Nonpoint source (NPS) pollution (e.g., siltation/sedimentation and stream habitat loss or degradation) was the primary cause of aquatic life use impairment. Other forms of pollution causing impairment are trace element contamination, flow alteration and oxygen depletion. The primary sources of pollutants affecting aquatic life use in the state are cropland erosion and runoff, animal feeding operations and poor grazing management. Other sources linked to aquatic life use impairment are point source discharges, urban runoff and hydrologic modifications (e.g., upstream impoundments, low-head dams, channelization, flow regulation and diversion, riparian vegetation removal, wetland drainage).

Recreation use was assessed on 6,617 miles of rivers and streams in the state. Recreation use was fully supporting, fully supporting but threatened and not supporting on 1,536 miles, 3,421 miles and 1,660 miles, respectively. Fecal coliform bacteria data collected from monitoring stations across the state were the primary indicators of recreation use attainment. For this reason, pathogens (as reflected by fecal coliform bacteria) are the primary cause of recreation use impairment in North Dakota. The primary sources of fecal coliform bacteria contamination are animal feeding operations and riparian area grazing.

Drinking water supply use is classified for 5,560 miles of rivers and streams in the state. Of the 1,738 miles assessed for this report, only 86 miles (4.9 percent) were assessed as threatened for drinking water supply use. The primary threats are taste and odor problems.

A total of 4,095 miles of rivers and streams were identified as capable of supporting a sport fishery from which fish could be used for consumption. Based on the EPA fish tissue of 0.3 micrograms (μg) methyl-mercury/gram of fish tissue, only the Red River of the North was assessed as not supporting fish consumption. While there are many potential sources of methyl-mercury (both anthropogenic and natural), to date there have been no specific causes or sources identified for the mercury present in North Dakota fish.

A total of 197 lakes and reservoirs, representing 700,315.89 surface acres, were assessed for this report. The remaining 50 lakes and reservoirs, included in the ADB but not assessed, represent 61,455.61 acres or only 5.5 percent of the total lake and reservoir acres in the state. One-hundred-twenty-four (124) lakes and reservoirs, representing 686,115.1 acres, were assessed as fully supporting aquatic life use; in other words, they are considered capable of supporting and maintaining a balanced community of aquatic organisms. Of this total, 30 lakes and reservoirs, representing 376,606.3 acres, are considered threatened. A threatened assessment means that if water quality and/or watershed trends continue, it is unlikely these lakes will continue to support aquatic life use. The lakes and reservoirs will begin to experience more frequent algal blooms and fish kills. They will display a shift in trophic status from a mesotrophic or eutrophic condition to a hypereutrophic condition. Only three lakes, totaling 171.8 acres, were assessed as not supporting aquatic life use. One of the primary causes of aquatic life impairment to lakes and reservoirs is low dissolved oxygen (DO) in the water column. Low DO in lakes can occur in summer (summer kills) but usually occurs in the winter under ice-cover conditions. When fish kills occur, low DO-tolerant fish species (e.g., carp, bullhead, white suckers) will be favored, resulting in a lake dominated by these rough fish species. Pollutants which stimulate the

production of organic matter, such as plants and algae, can also cause aquatic life impairment. Two secondary pollutant causes are excessive nutrient loading and siltation.

Major sources of nutrient loading to the state's lakes and reservoirs are erosion and runoff from cropland; runoff from animal feeding operations (e.g., concentrated livestock feeding and wintering operations); and hydrologic modifications. Hydrologic modifications, such as wetland drainage, channelization and ditching, increase the runoff and delivery rates to lakes and reservoirs, in effect increasing the size of a lake's watershed.

Recreation use (e.g., swimming, waterskiing, boating, sailing, sunbathing) was assessed for 686,250.1 lake and reservoir acres in the state. Of this total, two (2) lakes, representing 5,546.8 acres, were assessed as not supporting use for recreation. The primary cause of use impairment is excessive nutrient loading, which results in nuisance algal blooms and noxious aquatic plant growth. Sources of nutrients causing algal blooms and weed growth were described earlier. Thirty-seven (37) lakes and reservoirs, totaling 135,366.4 acres, were assessed as threatened.

One-hundred-ninety-six (196) lakes and reservoirs, representing 699,430.6 acres, were assigned the use for fish consumption. Of these, only Devils Lake and Lake Sakakawea had sufficient methyl-mercury fish tissue data and fish population survey data necessary to calculate weighted average concentrations and to assess fish consumption use. Based on these data, both were assessed as not supporting fish consumption use. The remaining 194 lakes and reservoirs which support a sport fishery were not assessed for this report. Sources of methyl-mercury in fish remain largely unknown. Potential sources of mercury include natural sources and atmospheric deposition.

Five reservoirs (Lake Sakakawea, Lake Ashtabula, Homme Dam, Bisbee Dam and Mt. Carmel Reservoir) are currently used either directly or indirectly as municipal drinking water supplies, while two others (Patterson Lake and Renwick Dam) serve as back-up water supplies in the event the primary water supplies should fail. Homme Dam, Mt. Carmel Reservoir and Lake Sakakawea were assessed as fully supporting drinking water supply use. Drinking water supply use was not assessed for the remaining lakes and reservoirs.

Section 303(d) of the CWA and its accompanying regulations require each state to list waterbodies (i.e., lakes, reservoirs, rivers, streams and wetlands) which are considered water quality limited and require load allocations, waste load allocations and TMDLs. This list has become known as the "TMDL list" or "Section 303(d) list." A waterbody is considered water quality limited when it is known that its water quality does not meet applicable standards or is not expected to meet applicable standards. Waterbodies can be water quality limited due to point source pollution, NPS pollution or both.

In considering whether or not applicable water quality standards are being met, the state should not only consider the narrative and numeric criteria set forth in the standards but also the classified uses defined for the waterbody and whether the use or uses are fully supported or not supported due to any pollutant source or cause. Where a waterbody is water quality limited, the state is required to determine in a reasonable time frame the reduction in pollutant loading necessary for that waterbody to meet water quality standards, including its beneficial uses. The process by which the pollutant-loading capacity of a waterbody is determined and the load is

allocated to point and nonpoint sources is called a total maximum daily load (TMDL). While the term “total maximum daily load” implies that loading capacity is determined on a daily time scale, TMDLs can range from meeting an instantaneous concentration (i.e., an acute standard) to computing an acceptable annual phosphorus load for a lake or reservoir.

When a state prepares its list of water quality-limited waterbodies, it is required to prioritize waterbodies for TMDL development and to identify those waterbodies which will be targeted for TMDL development within the next two years. Factors to be considered when prioritizing waterbodies for TMDL development include: (1) the severity of pollution and the uses which are impaired; (2) the degree of public interest or support for the TMDL, including the likelihood of implementation of the TMDL; (3) recreational, aesthetic and economic importance of the waterbody; (4) the vulnerability or fragility of a particular waterbody as an aquatic habitat, including the presence of threatened or endangered species; (5) immediate programmatic needs, such as wasteload allocations needed for permit decisions or load allocations for Section 319 NPS project implementation plans; and (6) national policies and priorities identified by EPA.

After considering each of the six factors, the state has developed a two-tiered priority ranking. Assessment units (AUs) listed as “High” priority are: (1) lakes and reservoirs and river and stream segments for which TMDLs are scheduled to be completed and submitted to EPA in the next two years; or (2) lakes and reservoirs and river and stream segments for which TMDL development projects are scheduled to be started in the next two years. The majority of these “High” priority AUs were identified as such, based largely on their degree of public support and interest and the likelihood of implementation of the TMDL once completed. “Low” priority AUs are those river and stream segments and lakes and reservoirs that are scheduled for completion in the next eight years.

The 2008 TMDL list is represented by 225 AUs (32 lakes and reservoirs and 193 river and stream segments) and 389 individual waterbody/beneficial use/pollutant combinations. For purposes of TMDL development, each waterbody/beneficial use/pollutant combination requires a TMDL. Of this total, the department has targeted 80 waterbodies or 112 waterbody/beneficial use/pollutant combinations for completion in the next two to four years. These “high” priority waterbodies are AUs for which the monitoring is either completed, near completion or has recently been initiated. Based on an anticipated TMDL completion schedule of approximately 40 additional waterbody/beneficial use/pollutant combinations per year following 2010, the department expects to complete TMDLs for all 2008-listed waters in the next eight years. With the continued commitment to adequate TMDL development staffing and with a continuation in the growth of funding for TMDL development projects in the state, the department is confident it will meet its TMDL development schedule.

PART II. INTRODUCTION

The Clean Water Act (CWA) contains several sections which require states to report on the quality of their waters. Section 305(b) (*State Water Quality Assessment Report*) requires a comprehensive biennial report, and Section 303(d) requires, from time to time, a list of a state's water quality-limited waters needing total maximum daily loads (TMDLs). In its regulations implementing Section 303(d), the U.S. Environmental Protection Agency (EPA) has defined "time to time" to mean April 1 of every even-numbered year. While due at the same time, states have historically submitted separate reports to EPA under these two sections. However, in guidance provided to the states by EPA dated July 29, 2005 (EPA, 2005), EPA suggested that states combine these two reports into one integrated report. The following is a brief summary of the requirements of each reporting section.

A. Section 305(b) Water Quality Assessment Report

The primary purpose of this *State Water Quality Assessment Report* is to assess and report on the extent to which beneficial uses of the state's rivers, streams, lakes, reservoirs and wetlands are met. Section 305(b) of the Clean Water Act requires states to submit this assessment report every two years; the information presented in this report is for the reporting period of 2006-2007. The Section 305(b) report is a summary report that presents information on use impairment and the causes and sources of impaired or threatened uses for the state as a whole.

This report is not a trends report, nor should the data or information in this report be used to assess water quality trends. Factors which complicate and prohibit comparisons between reporting years include changes in the number of sites, the quality of data upon which assessment information is based and changes to the estimated river and stream miles.

B. Section 303(d) TMDL List of Water Quality-limited Waters

While the Section 305(b) report is considered a summary report, Section 303 and its accompanying regulations (CFR Part 130 Section 7) require each state to list individual waterbodies (i.e., lakes, reservoirs, rivers, streams and wetlands) which are considered water quality limited and which require load allocations, waste load allocations and TMDLs. This list has become known as the "TMDL list" or "Section 303(d) list."

A waterbody is considered water quality limited when it is known that its water quality does not or is not expected to meet applicable water quality standards. Waterbodies can be water quality limited due to point sources of pollution, nonpoint sources (NPS) of pollution or both.

In considering whether or not applicable water quality standards are being met, the state should not only consider the narrative and numeric criteria set forth in the standards to protect specific uses, but also the classified uses defined for the waterbody and whether the use or uses are fully supported or not supported due to any pollutant source or cause. Therefore, a waterbody could be considered water quality limited when it can be demonstrated that a beneficial use (e.g., aquatic life or recreation) is impaired, even when there are no demonstrated exceedances of either the narrative or numeric criteria. In cases where there is use impairment and no exceedance of the numeric standard, the state should provide information as to the cause of the

impairment. Where the specific pollutant (e.g., copper or phosphorus) is unknown, a general cause category (e.g., metals or nutrients) should be included with the waterbody listing.

Section 303(d) of the CWA and accompanying EPA regulations and policy only require impaired and threatened waterbodies to be listed and TMDLs developed when the source of impairment is a pollutant. Pollution, by federal and state definition, is “any man-made or man-induced alteration of the chemical, physical, biological and radiological integrity of water.” Based on the definition of a pollutant provided in Section 502(6) of the CWA and in 40 CFR 130.2(d), pollutants would include temperature, ammonia, chlorine, organic compounds, pesticides, trace elements, nutrients, biochemical oxygen demand (BOD), sediment and pathogens. Waterbodies impaired by habitat and flow alteration and the introduction of exotic species would not be included in the Section 303(d) TMDL list, as these impairment categories would be considered pollution and not pollutants. In other words, all pollutants are pollution, but not all pollution is a pollutant.

Where a waterbody is water quality limited, the state is required to determine, in a reasonable timeframe, the reduction in pollutant loading necessary for that waterbody to meet water quality standards, including its beneficial uses. The process by which the pollutant loading capacity of a waterbody is determined and the load is allocated to point and nonpoint sources is called a total maximum daily load (TMDL). While the term “total maximum daily load” implies that loading capacity is determined on a daily time scale, TMDLs can range from meeting an instantaneous concentration (i.e., an acute standard) to computing an acceptable annual phosphorus load for a lake or reservoir.

Section 303(d) requires states to submit their lists of water quality-limited waterbodies “from time to time.” Federal regulations have clarified this language; therefore, beginning in 1992 and by April 1 of every even-numbered year thereafter, states are required to submit a revised list of waters needing TMDLs. North Dakota’s last TMDL list was submitted to EPA on April 13, 2006 and was approved by EPA on June 27, 2006.

This Section 303(d) list includes waterbodies not meeting water quality standards, waterbodies needing TMDLs and waterbodies which have been removed from the 2006 list. Reasons for removing a waterbody from the 2006 list include: (1) a TMDL was completed for the waterbody/pollutant combination; (2) the applicable water quality standard is now attained and/or the original basis for the listing was incorrect; (3) the applicable water quality standard is now attained due to a change in the water quality standard and/or assessment methodology; (4) the applicable water quality standard is now attained due to restoration activities; or (5) sufficient data and/or information lacking to determine water quality status and/or the original basis for listing was incorrect.

PART III. BACKGROUND

A. Atlas

Table III-1. Atlas

Topic	Value
State Population ¹	639,715
State Surface Area (Sq. Miles)	70,700
Total Miles of Rivers and Streams ²	54,607.04
Total Miles of Rivers and Streams by Stream Class ³	
Class I, IA and II Streams	5,973.18
Class III Streams	48,633.86
Total Miles of Rivers and Streams by Basin	
Red River (including Devils Lake)	11,990.94
Souris River	3,670.18
Upper Missouri (Lake Sakakawea)	13,877.43
Lower Missouri (Lake Oahe)	22,276.60
James River	2,791.89
Border Miles of Shared Rivers and Streams ⁴	429.84
Total Number of Lakes and Reservoirs ⁵	247
Number of Natural Lakes	109
Number of Manmade Reservoirs	138
Total Acres of Lakes and Reservoirs	761,771.51
Acres of Natural Lakes	218,615.85
Acres of Manmade Reservoirs ⁶	543,155.66
Total Acres of Lakes and Reservoirs by Lake Class ⁷	
Class 1	481,841.29
Class 2	62,890.46
Class 3	145,602.15
Class 4	9,096.70
Class 5	885.30
Unclassified	61,455.61
Acres of Freshwater Wetlands ⁸	2,500,000

¹ Based on U.S. Census Bureau estimates provided on July 1, 2007

² Total miles are based on rivers and streams entered into the Assessment Database (ADB) and reach indexed to the 1:100,000 scale National Hydrography Dataset (NHD).

³ Stream classes are defined in the *Standards of Quality for Waters of the State* (North Dakota Department of Health, 2006). In general, Classes I, IA and II streams are perennial, while Class III streams are intermittent or ephemeral.

⁴ Includes the Bois de Sioux River and the Red River of the North

⁵ Number includes only the lakes and reservoirs which are publicly owned and are in the ADB.

⁶ Estimates based on surface acreage at full pool elevation.

⁷ Lake and reservoir classes are defined in the *Standards of Quality for Waters of the State* (North Dakota Department of Health, 2006).

⁸ Estimate provided by Dahl, T.E., *Wetlands - Losses in the United States: 1780's to 1980's*, Washington, D.C., U.S. Fish and Wildlife Service Report to Congress, 1990.

B. Total Waters

The North Dakota Department of Health (hereafter referred to as the department) currently recognizes 247 public lakes and reservoirs. Of the 247 public lakes and reservoirs recognized as public waters and included in the Assessment Database (ADB), only 198 are included in the state's water quality standards as classified lakes and therefore are assigned designated beneficial uses (Table III-1). The remaining 50 lakes and reservoirs, while included in the state's estimate of total lake acres, are not classified and therefore were not assessed for this report.

Of the 247 public lakes and reservoirs included in the ADB, there are 138 manmade reservoirs and 109 natural lakes. All lakes and reservoirs included in this assessment are considered significantly publicly owned. Reservoirs are defined as waterbodies formed as a result of dams or dugouts constructed on natural or manmade drainages. Natural lakes are waterbodies having natural lake basins. A natural lake can be enhanced with outlet control structures, diversions, or dredging. Based on the state's Assessment Database (ADB), the 138 reservoirs have an areal surface of 543,156 acres. Reservoirs comprise about 71 percent of North Dakota's total lake/reservoir surface acres. Of these, 480,731 acres or 62 percent of the state's entire lake and reservoir acres are contained within the two mainstem Missouri River reservoirs (Lake Sakakawea and Lake Oahe). The remaining 136 reservoirs share 62,425 acres, with an average surface area of 459 acres.

The 109 natural lakes in North Dakota cover 218,616 acres, with approximately 117,697 acres¹ or 54 percent attributed to Devils Lake. The remaining 108 lakes average 934 acres, with half being smaller than 250 acres.

There are an estimated 54,607 miles of rivers and streams in the state. Estimates of river stream miles in the state are based on rivers and streams entered into the ADB and reach indexed to the 1:100,000 scale National Hydrography Dataset (NHD).

In this report, the state has been divided into five basins: Red River (including Devils Lake), Souris River, Upper Missouri River (Lake Sakakawea), Lower Missouri River (Lake Oahe) and James River (Figure III-1). The atlas provided in Table III-1 provides a basin-by-basin estimate of total river and stream miles.

¹ The estimated surface area for Devils Lake is based on a lake elevation of 1446 mean sea level (msl), which is the elevation at which water overflows to Stump Lake.




Figure III-1. Major Hydrologic Basins in North Dakota

C. Water Pollution Control Program

Chapter 1. Water Quality Standards Program

State water quality standards describe the policy of the state which is to protect, maintain and improve the quality of water for use as public and private water supplies; for propagation of wildlife, fish and aquatic life; and for domestic, agricultural, industrial, recreational and other legitimate beneficial uses.

The state classifies its surface water resources into five categories. The assignment of a waterbody into a particular classification is based on the water quality of record (1967), existing uses at that time, hydrology and natural background factors.

Water quality standards also identify specific numeric criteria for chemical, biological and physical parameters. The specific numeric standard assigned to each parameter ensures protection of the beneficial uses for that classification. The water quality standards also contain general conditions, termed “narrative standards,” applicable to all waters of the state. These general conditions contain provisions not specifically addressed in numeric criteria. These conditions add an extra level of protection for water quality.

The department has also developed a narrative biological goal for all waters of the state. The goal is to restore all surface waters to a condition similar to that of sites or waterbodies determined to be regional reference sites. The goal is non-regulatory; however, it may be used in combination with other information in determining whether aquatic life uses are attained. The state is also in the process of developing “biological criteria.” These criteria will define ecological conditions in state waters and set goals for their attainment.

In addition to numeric and narrative standards and the beneficial uses they protect, a third element of water quality standards is antidegradation. The fundamental concept of antidegradation is the protection of waterbodies which currently have better water quality than applicable standards. Antidegradation policies and procedures are in place to maintain high quality water resources and prevent them from being degraded to the level of water quality standards.

State water quality standards have established three categories or tiers of antidegradation protection. Category 1 is a very high level of protection and automatically applies to all Class I and IA rivers and streams, all Class 1, 2 and 3 lakes and reservoirs, and wetlands that are functioning at their optimal level. Category 1 may also apply to some Class II and III rivers and streams, but only if it can be demonstrated that there is remaining pollutant assimilative capacity, and both aquatic life and recreation uses are currently being supported. Category 2 antidegradation protection applies to Class 4 and 5 lakes and reservoirs and to Class II and III rivers and streams not meeting the criteria for Category 1. Category 3 is the highest level of protection and is reserved for Outstanding State Resource Waters. Waterbodies may only be designated Category 3 after they have been determined to have exceptional value for present and future potential for public water supplies, propagation of fish or aquatic biota, wildlife, recreation, agriculture, industry, or other legitimate beneficial uses.

The U.S. EPA requires the department to review and update, as necessary, the state water quality standards based on new information and EPA guidance a minimum of every three years. This process is termed the “triennial review.” Issues currently being considered for this review are beneficial use designations for wetlands and associated numeric criteria. Currently, wetlands are considered waters of the state and are protected by general conditions.

The department is also in the process of developing nutrient criteria which are needed to address the eutrophication of the state’s surface waters. Excessive nutrients typically manifest themselves as elevated amounts of algae in lakes and reservoirs and as epiphytic algae in streams and rivers. In preparation for the development of nutrient criteria, the department has developed a plan for developing technically defensible nutrient criteria specific to the unique resources of North Dakota. The Nutrient Criteria Development Plan describes the anticipated conceptual approach for developing nutrient water quality criteria. The plan specifically focuses on lotic systems (i.e., small to large Wadeable and non-Wadeable streams and rivers) and lentic systems (i.e., lakes and reservoirs). The plan is intended to provide clear and meaningful guidance for the development of nutrient criteria within North Dakota. The report does not represent a binding commitment, and modification of the plan will likely be needed as new information becomes available or unanticipated issues arise.

The approach described by the Nutrient Criteria Development Plan has enabled North Dakota to explore in detail the feasibility of implementing various development concepts. The department, through funding provided by EPA Headquarters, is currently performing a pilot project on establishing numeric standards for lentic systems. This project will result in a proposed state-wide classification system for all lake and reservoir systems based on an intensive examination and analysis of database information. The project will identify a major geographic region of the state and assess nutrient criteria for the lakes within that region. Outcomes of the regional assessment will determine what numeric endpoints should be set for different types of lakes and reservoirs (i.e., small versus large water bodies).

Chapter 2. Point Source Control Program

The department regulates all releases of wastewater from point sources into waters of the state. The regulation of all point source discharges is the responsibility of the department's Division of Water Quality. The North Dakota Pollutant Discharge Elimination System (NDPDES) Program requires all point source dischargers (municipal and industrial) to obtain a permit. NDPDES permits outline technology-based and/or water quality-based limits for wastewater discharges. There are approximately 400 facilities (25 percent industrial and 75 percent municipal) that are permitted for discharges of treated wastewater.

The NDPDES Program also includes coverage for stormwater discharges associated with industrial and construction activities. There are approximately 382 facilities covered under general permits for stormwater discharges from industrial activities. Included in these general permits are requirements for monitoring and sampling of stormwater discharges. All discharge data is evaluated and used to update the standard pollution prevention practices that are currently used in the state. These facilities must implement pollution prevention plans which are intended to improve the quality of stormwater discharges.

There are approximately 1070 facilities covered for construction stormwater and 18 municipal separate storm sewer system (MS4) stormwater permits. The department continues to implement the Stormwater Phase II regulations (effective December 8, 1999) to the maximum extent possible. The department also works with the regulated small MS4s and the Red River Work Group on issues relating to stormwater discharges. The focus of activities with MS4s continues to be on the development of ordinances or other regulatory mechanisms for local construction site erosion and sediment control and post construction controls (MCMs 4 and 5).

The permitting procedure for small construction is being revised to better address building/construction in subdivisions. Several of the forms and guidance materials for the industrial permit and the construction permit were revised or created to assist permit holders. A stormwater sampling guide was developed and posted on the department's website, and a new construction stormwater pollution prevention plan guide should be completed in 2008. The department continues to provide stormwater education, including an annual conference on stormwater issues.

Many of the wastewater treatment systems in North Dakota consist of impoundments or lagoons. The availability of land and the low operation and maintenance costs are the main reasons for their use and acceptance in North Dakota. These wastewater stabilization pond systems discharge intermittently, and the discharges are short in duration. The average discharge duration is less than six days in length with the majority of the discharges occurring in the spring and fall. A facility that receives permission to discharge treated wastewater is required to monitor the discharge for quality and quantity data. This information is submitted to the department in monthly, quarterly, or semi-annual reports which are tracked and monitored for compliance with the conditions outlined in the permit.

The overall quality of wastewater is commonly indicated by 5-day biochemical oxygen demand (BOD-5) and total suspended solids (TSS). Typically, high concentrations of BOD-5 and TSS indicate poor treatment system performance which can present an environmental concern.

Treated wastewater from many of the state's permitted facilities is discharged over land or through ditches before it reaches waters of the state. In such cases, it is likely the reported concentrations for BOD-5 and TSS are further reduced prior to entering a waterbody.

Figure III-2 shows the mean annual concentrations of BOD-5 and TSS reported for wastewater discharges in North Dakota. Data used to generate this graph are for the years 1985 through December 2007. The overall trend in the mean annual concentrations of these two pollutants appears to be decreasing, which generally means wastewater treatment systems in the state are doing a good job of operating. It should be noted that the slight increase in BOD-5 noted for 2007 may be due to excessive rains that occurred in certain areas of the state during the year.




Figure III-2. Average Annual BOD-5 day and TSS Concentrations for Wastewater Discharges in North Dakota (1985-2007)

Western North Dakota continued to receive below-normal precipitation, while eastern North Dakota is received normal to slightly above normal precipitation. Localized flooding in certain areas compounded wastewater treatment and storage problems, resulting in bypasses and lagoon overflows. Several communities in the state initiated major improvements to their wastewater collection and treatment systems. The number of discharges and total volume of water discharged annually has leveled off to near-normal levels compared to the upward trend that started in 1993. The NDPDES Program also requires all permitted industrial and municipal facilities to report spills and releases of wastewater. Most releases were related to mechanical failure and/or excessive precipitation events.

Generally, development of Total Maximum Daily Loads (TMDLs) has not been required for point source discharges in North Dakota. TMDL development activity occurs mainly in rural watersheds dealing with nonpoint source pollution issues. There is effective internal coordination during the development of TMDLs and waste load allocation (WLA) requirements in NDPDES permits, and no formal tracking mechanism is required or necessary in the NDPDES Program at this time. At the present time, no permits have been modified or reissued to implement WLAs in approved TMDLs. With the cooperation of the cities of Fargo and Moorhead, the department and the Minnesota Pollution Control Agency are in the process of finalizing a bacteria TMDL for the Red River in the Fargo area. The department is also finalizing a low-flow TMDL for the James River near Jamestown. Results of these TMDLs will be used to determine if modifications to NDPDES permits are needed for the cities of Fargo and Jamestown, respectively.

Toxic pollutants in wastewater discharges are a concern, particularly for the larger cities and industries in North Dakota. They are regulated through the Industrial Pretreatment Program. The department received primacy (effective September 9, 2005) from EPA Region VIII to implement the program in North Dakota. The cities of Grand Forks, Fargo, Bismarck, Mandan and West Fargo have approved pretreatment programs. The department is working closely with pretreatment personnel from select industries and municipalities and is in the process of organizing a pretreatment workgroup in the state.

All waters of the state “shall be free from substances attributable to municipal, industrial or other discharges in concentrations or combinations which are toxic or harmful to humans, animals, plants or resident biota.” This narrative water quality standard is enforced, in part, through Whole Effluent Toxicity (WET) requirements the NDPDES permits. All major municipal and industrial permittees must monitor their discharges for WET on a regular basis. At a minimum, these municipalities and industries sample quarterly for WET testing with results submitted for the department’s review. Failure of WET tests can result in toxicity identification evaluations (TIEs) to determine the cause of the toxicity in the effluent. TIEs that have been completed in the state have resulted in major and minor improvements to wastewater treatment systems.

Several cities and industries have selected biological treatment methods to improve their wastewater treatment systems and the quality of their discharge water. The biological treatment system at the Amoco Refinery in Mandan is providing consistent, advanced treatment of its wastewater. The Devils Lake “Lemna” system was specifically designed to remove phosphorus from the wastewater. This treatment system generally provides an advanced level of nutrient removal; however, flooding in the Devils Lake basin since 1993 has taxed the system beyond its design capabilities. An interim phosphorus limit/goal has been established to compensate for the adverse operating conditions which currently prevail.

The wetland treatment system for the city of Minot continues to provide low ammonia concentrations in the final effluent. The city is capable of continuously discharging a quality effluent during non-ice conditions which adds to the river flow and enhances aesthetic river quality. This is extremely beneficial since the Souris River has a history of poor river quality and low/no-flow conditions during several months of the year.

American Crystal Sugar uses a combination of lagoons and constructed wetlands for wastewater treatment and polishing/finishing at both its Hillsboro and Drayton plants. The final effluent from these facilities surpasses the federal effluent criteria for sugar beet processing plants. The 1.5 million-gallon-per-day (MGD) anaerobic digester and clarifier at the Hillsboro plant maximizes the performance of the existing aerobic digester, resulting in a reduction of the feed water strength while maintaining a constant temperature throughout the season. American Crystal Sugar can then route this high quality water to its wetlands earlier in the season, maximizing the wetland's ability to treat the facility's wastewater prior to discharge.

The Minn-Dak Farmers Cooperative sugar beet processing plant uses both mechanical and facultative lagoons for wastewater treatment at its Wahpeton facility. The wastewater receives additional treatment/polishing in the large discharge reservoir from which the final effluent is discharged through an in-stream diffuser to the Red River. Minn-Dak coordinates its discharges with the Cargill Corn Milling plant since both facility permits contain receiving stream quality requirements for sulfate, chloride and total dissolved solids (TDS).

Cargill Corn Milling (ProGold) produces high fructose corn syrup at its facility near Wahpeton. The plant discharges select waste streams to the Red River on a continuous basis with ponds available to store wastewater when treatment is inadequate or when the river would be adversely affected. Wastewater high in TDS is stored in two ponds on site. To meet the requirements of ProGold's permit, these ponds must be discharged according to conditions in the Red River and coordinated with discharges from downstream users and the Minn-Dak Farmers Cooperative Plant. The background water quality in the Red River has been the most limiting factor for coordinating discharges from the ponds, particularly when flows are predominantly from Lake Traverse.

The city of Fargo continues to provide a quality effluent to the Red River. Wastewater treatment consists of pretreatment/odor control, primary clarification, trickling filters, nitrification filters, final clarification and disinfection. Residuals management (biosolids) consists of digesters, sludge drying beds and belt presses. The processed solids are used as cover at the municipal landfill. Fargo still maintains its six 90-acre wastewater stabilization ponds which can be used for storage during times of flooding or when an upset occurs in the treatment plant. The city is moving forward with the construction of a new force main interceptor to transport wastewater from south Fargo to the treatment plant located in the north part of town. With the addition of the new force main, the "infiltration and inflow" (I & I) issues in downtown Fargo that occur during heavy rain events will be greatly reduced.

The city of Bismarck continues with upgrades to its wastewater treatment plant. The facility master plan consists of short-term and long-term improvements to the facility. In phase one, three large storage tanks for biosolids retention were constructed at the plant. In addition to providing the necessary storage during the winter months, the tanks also assist with BOD-5 permit compliance. The second phase of improvements consists of construction of the new pre-treatment facility which started in 2007. Other planned improvements to the facility include an additional primary clarifier, as well as updates to the existing primary clarifiers, trickling filter, final clarifier and control systems.

The city of Grand Forks started operation of its new wastewater treatment facility in late fall

2002. The activated sludge plant uses a European technology of “Micro-Bubble” flotation and is designed for 15 MGD. Plant operations staff experienced minor problems associated with startup and limited knowledge of this type of system. Plant staff and the contractors continue to fine tune the process controls to provide optimal wastewater treatment. The effluent from the treatment plant is routed to the stabilization ponds which the city continues to operate. In the future, the city should be discharging on a continual basis to the river.

The Mandan wastewater treatment plant consists of a “biolac wave oxidation” process which includes extended aeration for BOD removal, nitrification and sludge stabilization. The whole process was constructed in the city’s old primary aerated lagoon cell. The plant has averaged more than 90 percent removal of BOD and TSS. This plant is the first in the state to use ultraviolet disinfection of the treated wastewater.

The Jamestown mechanical wastewater treatment plant was designed to treat agricultural process wastes which are blended with domestic waste from the city. The excess oil and grease from the potato processing facility was addressed with the addition of a large grease and sand interceptor at the head works to the treatment plant. The city also has the capability of treating and storing wastewater in its lagoon system. Results of the low-flow TMDL for the James River will be used to determine if modifications to Jamestown’s NDPDES permits are needed.

The department continues to work on addressing stormwater noncompliance. The focus was on non-filers and on permitted facilities where there is water quality degradation and/or a threat to public health. Routine inspections resulted in informal enforcement letters requesting additional information and/or requiring repairs to best management practices (BMPs). In addition, the department issues formal warning letters citing apparent noncompliance with permit rules and water quality statutes. Notices of Violation (NOVs) and Consent Agreements are issued through the Attorney General’s office. The department initiated four consent agreements with penalties assessed. Penalties ranged from \$10,000 to \$24,000, which included both upfront and suspended penalties. For each case, the collected penalty exceeded any economic benefit of non-compliance.

The department is addressing all animal feeding operations impacting water quality through mechanisms or existing programs in the state. In addition, owners/operators are required to obtain permit coverage or eliminate the unacceptable conditions causing the operation to pollute and provide the information to EPA as needed.

The department incorporated the February 12, 2003 federal CAFO rules into the state program. This consisted of updates to the North Dakota Pollutant Discharge Elimination System (NDPDES) rules (NDAC 33-16-01) and Control of Pollution from Animal Feeding Operations rules (NDAC 33-16-03.1). These rules became final on January 7, 2005.

Because of the 9th District Court decision and pending revisions to the 2003 federal CAFO rules, the state has not issued an NDPDES permit at this time but continues to permit animal feeding operations facilities under the current state program (NDAC 33-16-03.1). Once the federal rules are final, the department will initiate the process to amend its state rules to be consistent with any new changes to the federal CAFO rules. For all state-permitted CAFOs, the permit facility data, permit event data and inspection data are entered into the state data base system. CAFO

inspections are performed yearly, and information is provided to EPA on a regular basis.

The department provides educational materials to livestock producers and the public on the impacts that livestock manure has on waters of the state. Several times each year, the department participates in presentations to producer groups. The department works closely with the Natural Resources Conservation Service (NRCS) and local health units on livestock manure systems. The department coordinates with the North Dakota Department of Agriculture and the North Dakota Stockmen's Association on assessing potential water quality impacts at livestock facilities. The department also meets with individual producers on site to determine what impacts the facility may have on water quality and discuss ways to prevent water quality impacts, if needed

The department works closely with local zoning boards and county commissions to help them recognize sensitive areas where livestock operations may cause problems and to encourage them to limit the expansion of operations in these areas. The department spearheaded a task force consisting of planning and zoning boards, producer groups and environmental groups to develop a model zoning ordinance for concentrated animal feeding operations.

The Operator Training Program is an important aspect of water quality protection. North Dakota regulations require a certified operator for municipalities with populations of greater than 500. The goal of the program is to conduct an inspection of each municipal treatment system at least once a year. These inspections verify proper system operation and reaffirm to the operator the importance of proper operation in protecting the state's water resources. The department also conducts annual wastewater operator training and certification seminars. In addition to the seminars, the program provides individual training and assistance to facilities encountering treatment problems.

Contracts were awarded to several health districts in the state to provide assistance in water pollution investigations. The contracts run through the state fiscal year (July 1 - June 30) and are for a two-year period. Activities associated with these contracts are water and wastewater inspections, odor readings at animal feeding operations and initial response to spills and releases to waters of the state.

Chapter 3. Nonpoint Source (NPS) Pollution Control Program

Surface water and ground water are two of North Dakota's most valuable natural resources. Water quality is affected by both natural and cultural, point source and nonpoint source (NPS) pollution, with NPS pollution being the major factor affecting surface water quality in the state. Ground water quality has remained relatively unaffected by major sources of pollution. However, some aquifers have experienced minor water quality impairments (see Part VII. Ground Water Assessment).

All rivers, streams, reservoirs and lakes assessed within the state are impacted to some degree by NPS pollution. Generally, most surface water quality impacts are associated with agricultural activities in these watersheds. Ground water impacts result from the improper use of agricultural chemicals, leaking underground petroleum storage tanks and pipelines, wastewater impoundments, oil and gas exploration activities, septic systems and improperly located and maintained solid waste disposal sites.

NPS pollution control efforts to maintain or improve the beneficial uses of North Dakota's water resources are primarily accomplished through the North Dakota NPS Pollution Management Program. The voluntary NPS Program is dependent on the formation of partnerships and coordination with local resource managers to effectively reduce and/or prevent NPS pollution from impairing beneficial uses of the state's water resources. Over the long term, through these coordinated efforts, the cumulative benefits of the local projects will help the department achieve its mission and long-term goal as identified in the North Dakota NPS Pollution Management Program Plan. The NPS Program's mission statement and long-term goal are as follows:

North Dakota NPS Program Mission: "To protect or restore the chemical, physical and biological integrity of the waters of the state by promoting locally sponsored, incentive-based, voluntary programs where those waters are threatened or impaired due to nonpoint sources of pollution."

North Dakota NPS Management Program Long Term Goal: "To initiate a balanced program focused on the restoration and maintenance of the beneficial uses of the state's water resources (i.e., streams, rivers, lakes, reservoirs, wetlands, aquifers) impaired by NPS pollution."

To achieve the long-term goal, an average of five watershed restoration projects will be targeted for implementation each year. The objective is to initiate 75 watershed restoration projects by 2013. To maintain program balance and strengthen support for the watershed initiatives, financial and technical resources will be used to complete NPS assessments or TMDLs on additional waterbodies and implement various public education projects. In most cases, these projects will be initiated and managed by local entities such as soil conservation districts (SCDs) or water resource boards.

The local or state projects supported with Section 319 funding can be placed under one of four different categories. These project categories are: (1) development phase projects; (2) educational projects; (3) technical support projects; and (4) watershed projects. Under each of these categories, there may also be one or more different project types or subcategories.

The primary purposes of the development phase projects are to identify beneficial use impairments or threats within specific waterbodies and determine the extent to which those threats or impairments are due to NPS pollution. Typically, development phase projects involve an inventory of existing data and supplemental monitoring to allow a thorough assessment of the targeted waterbody and its watershed. Through these efforts, the local project sponsors are able to: (1) determine the extent to which beneficial uses are being impaired by NPS pollution; (2) identify specific sources and causes of the pollutants; (3) establish preliminary pollutant reduction goals or TMDLs; and (4) identify management measures needed to restore or maintain the beneficial uses of the waterbody. Projects under this category include NPS Assessment Projects and TMDL Development Projects.

Educational projects are designed to increase public awareness and understanding of various NPS pollution issues and/or the solutions to specific NPS pollution concerns. The focus of these educational efforts may range from a local source or cause of NPS pollution to statewide measures that can be initiated to reduce NPS pollution. Educational tools typically include brochures, all media (TV, radio, newspaper), workshops, “how to” manuals, tours, exhibits and demonstrations. Two types of educational projects are currently being delivered in the state. The first are demonstration projects that focus on the development of on-the-ground demonstrations for educational purposes. The other type of educational project is public outreach, which focuses on the distribution of information on various local and/or state NPS pollution issues.

Projects designed to deliver technical or financial assistance to other ongoing NPS pollution management projects are identified as “Technical Support Projects.” These projects or programs are either offered statewide or targeted toward a “project area” that includes multiple NPS projects. The primary purpose of these projects is to deliver a specific service or “tool” to locally sponsored NPS projects. Specific types of assistance or management tools being delivered by the technical support projects include engineering designs, manure management planning, digitized soils, land use satellite imagery and wetland restoration/creation support.

The watershed project category includes the most comprehensive projects currently implemented through the NPS Program. These projects are typically long-term efforts designed to address documented NPS pollution impacts and beneficial use impairments within priority watersheds. Common objectives for watershed projects include: (1) protection and/or restoration of impaired beneficial uses through voluntary implementation of BMPs; (2) dissemination of information on local NPS pollution concerns and effective solutions to those concerns; and (3) evaluation of progress toward identified use attainment or NPS pollutant reduction goals. In nearly all cases, the goals and objectives of the watershed projects are based on data collected through some type of development project (e.g., NPS Assessment Project, TMDL development).

Section 319 funding is the primary source of financial support for projects addressing NPS pollution. Through the 2003, 2006 and 2007 Section 319 Grants (Active Grants), the NPS Program has provided funding to 72 local and state projects. The budgets and status of the locally sponsored projects and NPS Program staffing are provided in Table III-2.

Table III-2. Status and Budgets for Projects Supported Under the Fiscal Year 2003, 2006 and 2007 Section 319 Grants (1/1/03 -11/30/07)

Development Phase – NPS Assessment

Project Name	Status	319 Allocation	Local Match	Total Budget
Bear/Bonehill Creek Assessment	Completed	\$ 15,253	\$ 10,169	\$ 25,422
Cass Co. – Three Rivers Assessment Project	Active	\$ 136,372	\$ 90,915	\$ 227,287
Lake Hoskins Water Quality Assessment	Completed	\$ 18,066	\$ 12,044	\$ 30,110
McDowell Dam Alum Treatment Demo	Completed	\$ 47,664	\$ 31,776	\$ 79,440
Ransom Co. Sheyenne River Assessment	Completed	\$ 79,480	\$ 52,987	\$ 132,467
Red River Basin Volunteer Monitoring Network	Completed	\$ 47,829	\$ 31,886	\$ 79,715
Rice Lake Water Quality Improvement Project	Completed	\$ 448,200	\$298,800	\$ 747,000
Stutsman Co. Subwatershed Assessment Project	Active	\$ 11,845	\$ 7,897	\$ 19,742
Turtle River Assessment	Active	\$ 129,079	\$ 86,053	\$ 215,132
Unobligated Development Phase Fund	Active	\$ 354,699	\$236,466	\$ 591,165
Upper Goose River Watershed Assessment Project	Completed	\$ 82,159	\$ 54,773	\$ 136,932
Subtotal		\$1,370,646	\$913,764	\$2,284,410

Development Phase – TMDL Development

Project Name	Status	319 Allocation	Local Match	Total Budget
Armourdale Dam TMDL	Completed	\$ 4,055	\$ 2,703	\$ 6,758
Blacktail & McGregor TMDL Development Projects	Completed	\$ 14,998	\$ 9,999	\$ 24,997
Carbury Dam TMDL	Completed	\$ 6,184	\$ 4,123	\$ 10,307
Dickinson Dike TMDL Development – Phase II	Completed	\$ 2,800	\$ 1,867	\$ 4,667
Dickinson Dike TMDL Development – Phase III	Completed	\$ 6,455	\$ 4,303	\$ 10,758
Dickinson Dike TMDL Development - Phase I	Completed	\$ 6,853	\$ 4,569	\$ 11,422
McDowell Watershed TMDL	Completed	\$ 22,688	\$ 15,125	\$ 37,813
Northgate Dam TMDL	Completed	\$ 14,245	\$ 9,497	\$ 23,742
Subtotal		\$ 78,278	\$ 52,185	\$ 130,463

Education – Demonstration

Project Name	Status	319 Allocation	Local Match	Total Budget
Kelly Creek Water Quality Improvement Demonstration	Completed	\$ 7,860	\$ 5,240	\$ 13,100
SW North Dakota NPS/Water Quality I&E Project	Active	\$1,337,086	\$ 891,391	\$2,228,477
Subtotal		\$1,344,946	\$ 896,631	\$2,241,577

Table III-2 (cont.). Status and Budgets for Projects Supported Under the Fiscal Year 2003, 2006 and 2007 Section 319 Grants (1/1/03 -11/30/07)

Education – Public Outreach

Project Name	Status	319 Allocation	Local Match	Total Budget
Digital Taxonomic Keys for Aquatic Insects in ND	Completed	\$ 72,324	\$ 48,216	\$ 120,540
Envirothon Program	Active	\$ 142,948	\$ 95,299	\$ 238,247
Foster County – TREES Program	Active	\$ 630,523	\$ 420,349	\$1,050,872
Groundwater Pesticide Assessment Educational Program	Active	\$ 24,000	\$ 16,000	\$ 40,000
NDSU Livestock Waste Information & Assistance Program	Active	\$1,246,738	\$ 831,159	\$2,077,897
Project WET	Active	\$ 514,067	\$ 342,711	\$ 856,778
Statewide ECO ED Camp	Active	\$ 786,138		\$1,310,230
Subtotal		\$3,416,738	\$2,277,825	\$5,694,563

Local Project Support (TA or FA)

Project Name	Status	319 Allocation	Local Match	Total Budget
Adams Co. Livestock Manure Management Program	Active	\$1,009,584	\$ 673,056	\$ 1,682,640
Dairy Pollution Prevention Program	Active	\$2,477,358	\$1,651,572	\$ 4,128,930
Groundwater Sensitivity Mapping	Completed	\$ 329,704	\$ 219,803	\$ 549,507
Livestock Facility Assistance Program	Active	\$1,029,240	\$ 686,160	\$ 1,715,400
ND Waterbank Program	Completed	\$ 239,035	\$ 159,357	\$ 398,392
NDSU Satellite Imagery for WQ Protection	Completed	\$ 150,167	\$ 100,111	\$ 250,278
NPS BMP Team	Active	\$1,265,481	\$ 843,654	\$ 2,109,135
Project Safe Send – Dept. of Agriculture	Completed	\$ 140,895	\$ 93,930	\$ 234,825
Stockmen's Association Manure Management Specialist	Active	\$1,386,326	\$ 924,217	\$ 2,310,543
Subtotal		\$8,027,790	\$5,351,860	\$13,379,650

NPS Assessment – Multi Year Grant Award

Project Name	Status	319 Allocation	Local Match	Total Budget
Cannonball River Watershed Assessment – Phase II	Completed	\$ 3,020	\$ 2,013	\$ 5,033
Devils Lake Basin Assessment (00 WRAS)	Completed	\$ 3,864	\$ 2,576	\$ 6,440
NDSU Deep Soil Nitrogen Assessment	Completed	\$ 15,960	\$ 10,640	\$ 26,600
Nine Township Assessment (Knife River)	Completed	\$ 31,286	\$ 20,857	\$ 52,143
Pembina River Basin Assessment (99 WRAS)	Completed	\$ 71,632	\$ 47,755	\$ 119,387
Rocky Run Watershed Assessment – Phase I	Completed	\$ 0	\$ 0	\$ 0
UND Aquifer Denitrification Assessment	Completed	\$ 9,388	\$ 26,259	\$ 65,647
Subtotal		\$ 165,150	\$ 110,100	\$ 275,250

Table III-2 (cont.). Status and Budgets for Projects Supported Under the Fiscal Year 2003, 2006 and 2007 Section 319 Grants (1/1/03 -11/30/07)

NPS Program Staffing And Support

Project Name	Status	319 Allocation	Local Match	Total Budget
NPS Program Staffing & Support	Active	\$ 2,250,600	\$1,500,400	\$3,751,000
Subtotal		\$ 2,250,600	\$1,500,400	\$3,751,000

Watershed Project

Project Name	Status	319 Allocation	Local Match	Total Budget
Antelope Creek Watershed/Wild Rice Riparian Project	Active	\$ 880,949	\$ 587,299	\$ 1,468,248
Barnes Co. Sheyenne River Watershed (01 WRAS)	Active	\$ 1,228,114	\$ 818,743	\$ 2,046,857
Bear Creek Watershed	Active	\$ 877,402	\$ 584,935	\$ 1,462,337
Beaver Creek Watershed (99 WRAS)	Active	\$ 2,384,678	\$ 1,589,785	\$ 3,974,463
Bone Hill Creek Watershed	Active	\$ 633,660	\$ 422,440	\$ 1,056,100
Buffalo Springs & Lightening Creek Watersheds	Completed	\$ 250,587	\$ 167,058	\$ 417,645
Cannonball River TMDL Implementation Project	Active	\$ 165,065	\$ 110,043	\$ 275,108
Cedar Lake Watershed	Completed	\$ 205,105	\$ 136,737	\$ 341,842
Chanta Peta Watershed (00 WRAS)	Completed	\$ 109,153	\$ 72,769	\$ 181,922
Cottonwood Creek Watershed (99 & 02 WRAS)	Active	\$ 615,708	\$ 410,472	\$ 1,026,180
Crooked Creek Watershed (00 WRAS)	Completed	\$ 144,149	\$ 96,099	\$ 240,248
Deep Creek Watershed	Active	\$ 596,958	\$ 397,972	\$ 994,930
Dickey/LaMoure Livestock Manure Management Program	Active	\$ 933,900	\$ 622,600	\$ 1,556,500
Griggs Co. 319 Water Quality Project (99 WRAS)	Completed	\$ 702,570	\$ 468,380	\$ 1,170,950
Hay Creek Watershed - Phase IV	Completed	\$ 17,317	\$ 11,545	\$ 28,862
Hay Creek Watershed - Phase V	Completed	\$ 212,922	\$ 141,948	\$ 354,870
James River Headwaters Watershed	Active	\$ 685,000	\$ 456,667	\$ 1,141,667
Lake Hoskins Watershed	Active	\$ 230,142	\$ 153,428	\$ 383,570
Lower Pipestem Creek Watershed (02 WRAS)	Active	\$ 2,047,192	\$ 1,364,795	\$ 3,411,987
Maple Creek Watershed (00 WRAS)	Active	\$ 781,709	\$ 521,139	\$ 1,302,848
Middle Cedar Creek Watershed (00 WRAS)	Completed	\$ 345,714	\$ 230,476	\$ 576,190
Mirror Lake Watershed	Completed	\$ 71,856	\$ 47,904	\$ 119,760
Morton Co. Livestock Manure Management Program	Active	\$ 861,200	\$ 574,133	\$ 1,435,333
Nine Townships Watershed - Implementation Phase	Active	\$ 760,888	\$ 507,259	\$ 1,268,147
Pheasant Lake/Elm River Watershed (03 WRAS)	Active	\$ 934,834	\$ 623,223	\$ 1,558,057
Powers Lake Watershed (03 WRAS)	Active	\$ 538,205	\$ 358,803	\$ 897,008
Red River Riparian Project - Phases II & III (03 WRAS)	Active	\$ 1,603,428	\$ 1,068,952	\$ 2,672,380
Rocky Run Watershed - Phase II (02 WRAS)	Completed	\$ 443,710	\$ 295,807	\$ 739,517
Sheyenne River & Dead Colt Watersheds (Ransom Co.)	Active	\$ 635,919	\$ 423,946	\$ 1,059,865
Upper Cannonball Manure Management Program	Active	\$ 990,830	\$ 660,553	\$ 1,651,383
Upper Sheyenne Watershed (02 WRAS)	Completed	\$ 39,647	\$ 26,431	\$ 66,078
Wild Rice Watershed (99 & 00 WRAS)	Active	\$ 1,420,061	\$ 946,707	\$ 2,366,768
Subtotal		\$22,348,572	\$14,899,048	\$37,247,620
Grand Totals		\$39,002,720	\$26,001,813	\$65,004,533

Statewide delivery of the NPS Program is accomplished through six main goals identified in the NPS Program Management Plan. These goals, organized as individual sections of the management plan, are as follows:

- Resource Assessment - This section addresses the NPS Program's existing inventory/assessment system and future needs to improve or expand assessment efforts.
- Prioritization - This section discusses existing and future prioritization methods or strategies within the NPS Program.
- Assistance - This section focuses on "how" the financial and technical assistance available through the program is delivered to state/local project sponsors.
- Coordination - Development and maintenance of partnerships with private and local/state/federal agencies and organizations are described in this section.
- Information/Education - The program's multi-year strategy for public outreach and information dissemination is described under this section.
- Evaluation/Monitoring - Program and local project evaluation/monitoring efforts are addressed in this section.

Resource Assessment

Resource Assessment Goal: To accurately and thoroughly assess beneficial use support and the sources and causes of use impairments within the state's watersheds.

Resource assessment is being implemented at both the statewide and local levels. On a statewide basis, data (e.g., water quality, biological) collected by state and local staff are utilized to evaluate and document water quality and beneficial use trends of numerous waterbodies. At the local level, resource managers collect watershed-specific data to identify beneficial use and water quality impairments, establish waterbody priorities, develop watershed strategies and/or measure benefits of applied BMPs.

The locally sponsored NPS assessment or TMDL development projects are the primary means used to identify watershed priorities and management measures needed to address NPS pollution impairments. The local NPS assessments, commonly referred to as "development projects," provide the foundation for all watershed projects by identifying specific sources and causes of NPS pollutants impairing or threatening beneficial uses. This information is used to establish watershed priorities as well as to develop multi-year project implementation plans (PIPs) that address the identified beneficial use impairments. When applicable, department staff members also coordinate with the local sponsors to utilize the assessment data to develop TMDLs.

There are two sources of Section 319 financial support for assessment level projects. Short-term (i.e., 1-2 years) NPS assessment projects are supported with Section 319 funds available through the NPS Program's "Development Fund." Section 319 funds available under the development fund are unexpended funds reallocated from other NPS projects that were completed under budget. If the waterbody is also listed on the TMDL List, alternative funding sources (e.g.,

604[b], 104[b][3]) may also be used to support the assessment activities. For the multi-year or basin-wide NPS assessments, the local sponsors participate in the annual Section 319 grant application process to secure Section 319 support (base or incremental funding).

Since January 1, 2003, financial and/or technical assistance has been provided to 23 different assessment phase projects. Specific assessment phase projects are listed in Table III-2.

Prioritization

Prioritization Goal: Based on the most current inventory and assessment data, prioritize the state's waterbodies/watersheds for future NPS pollution assessment or abatement efforts.

The NPS Program utilizes a "process" rather than a "physical list" (with the exception of the TMDL List) to identify local waterbody priorities. On a statewide basis, waterbodies included on the TMDL List are considered high priority waterbodies for the development and implementation of watershed assessments. At the local level, the TMDL-listed waterbodies are also considered a high priority, although local resource managers may also establish priority rankings for other waterbodies not included on the TMDL List. For waterbodies lacking data and/or omitted from the TMDL List, a two-step process is used to establish the priorities. The first step involves a review of current information (e.g., local feedback, 305[b] reports, land use imagery) to establish a preliminary ranking for each subwatershed in the project area. These rankings are used to indicate the type of management or assessment activities needed in each subwatershed. The second step focuses on the development of a priority schedule for the implementation of the appropriate subwatershed assessment or management activities.

Typically, most waterbodies require the collection of additional data to identify beneficial use impairments and/or determine the sources and causes of pollutants impairing beneficial uses. For these waterbodies, the local sponsors coordinate with NPS Program staff to determine data collection needs and to establish a priority schedule for assessing the waterbodies. Following this prioritization process, financial and/or technical assistance can be provided to the sponsors to develop and implement quality assurance project plans (according to the priority schedule) to collect the necessary data. If sufficient data is already available on a waterbody to identify beneficial use impairments and the sources and causes of pollution, the local resource managers can seek Section 319 financial support to actively address the NPS pollutants impairing beneficial uses.

Assistance

Assistance Goal: Provide sufficient financial and technical assistance to local resource managers (e.g., SCDs, water resource boards) to ensure accurate identification of beneficial use and water quality impairments resulting from NPS pollution and effective development and completion of projects that will restore and/or maintain the beneficial uses of waterbodies impacted by NPS pollution.

NPS Program financial and/or technical assistance generally starts during the early stages of project development and continues throughout the implementation of the projects. Types of technical assistance being provided to local projects on an annual basis include project oversight, sample analysis, PIP review and comment, sample collection and project management training, quality assurance project plan development, distribution of educational materials and biological

monitoring support. Section 319 funding is the primary type of financial support for the NPS Program and locally sponsored NPS projects initiated in the state.

Since January 1, 2003, approximately 7 percent of the NPS Program budget has been used to support NPS Program staff. The balance of expenditures (i.e., 93 percent) has been used to support locally sponsored NPS pollution management projects. These local projects can be grouped under one of seven NPS project categories. Specific projects supported under each category are listed in Table III-2. Table III-3 lists the cumulative expenditures and distribution of costs for NPS program staffing and the different NPS project categories during the period of January 1, 2003 through November 30, 2007.

**Table III-3. Section 319 Allocations and Expenditures per Project Category
(1/1/03 -11/30/07)**

Project Category	319 Allocation	319 Expenditures	Percent of Total 319 Expenditures
Development Phase - NPS Assessment	\$ 1,370,646	\$ 943,921	4.43%
Development Phase - TMDL Development	\$ 78,278	\$ 78,276	0.37%
Education - Demonstration	\$ 1,344,946	\$ 1,074,238	5.04%
Education - Public Outreach	\$ 3,416,738	\$ 2,216,505	10.39%
Local Project Support (TA or FA)	\$ 8,027,790	\$ 4,289,685	20.11%
NPS Assessment - Multi Year Grant Award	\$ 165,150	\$ 165,148	0.77%
NPS Program Staffing and Support	\$ 2,250,600	\$ 1,512,769	7.09%
Watershed Projects	\$ 22,348,572	\$ 11,046,337	51.8%
Totals	\$ 39,002,720	\$ 21,326,879	

Coordination

Coordination Goal: Increase the effectiveness of NPS pollution management in the state by coordinating project development and implementation efforts with local, state and federal agencies and private organizations involved with natural resource management in the state.

Initiation and maintenance of a coordinated effort with appropriate entities is one of the most important activities within the project areas. At the onset of planning, the lead sponsors are encouraged to solicit the involvement of all groups or agencies that may have an interest in the planned project. For most projects, the involvement of multiple entities has helped ensure expertise is available and, in some cases, helped projects gain additional financial support.

Given the agricultural focus of most projects, local SCDs are the lead sponsors for most (70 percent) of the current projects. The SCDs provide the local leadership necessary to implement and manage projects as well as a “familiar face” to ensure effective communication with producers. However, as the NPS Program has expanded and diversified, more projects are being sponsored by other local and regional organizations (e.g., universities, state agencies, lake associations, resource conservation and development councils, water resource boards).

The NPS Task Force has also helped strengthen coordination among NPS projects and similar programs sponsored by other state or federal agencies and organizations. During the annual review process, the Task Force members become aware of the goals and objectives of the local NPS projects. This, in turn, gives them the opportunity to recognize and develop new partnerships that may strengthen projects/programs managed by their agency or organization. Conversely, during the review process, the local sponsors also gain a better understanding of what the Task Force member agencies can offer to their NPS pollution management projects. Organizations represented on the North Dakota NPS Source Pollution Task Force are listed in Table III-4.

Table III-4. Agencies/Organizations Represented on the North Dakota NPS Pollution Task Force.

Agency/Organization	Agency/Organization
Energy & Environmental Research Center	NDSU Extension Service
ND Farmers Union	USDA Farm Services Agency
USFS Dakota Prairies Grassland	ND Farm Bureau
ND Game & Fish Dept.	Bureau of Land Management
US Geological Survey	US Fish & Wildlife Service
ND Geological Survey	USDA Rural Development
US Bureau of Reclamation	ND Forest Service
ND Association of Soil Conservation Districts	State Soil Conservation Committee
ND Department of Agriculture	ND Water Resource Districts Association
US EPA Region VIII	Medora Grazing Association
ND Pork Producers	Grain Growers Association
ND Wildlife Federation	Rural Water Systems
USDA - Ag Research Station	USDA - NRCS
ND Parks & Recreation Dept.	ND Natural Resources Trust
ND State Water Commission	ND Stockmen's Association
ND Department of Health	

Information and Education

Information and Education Goal: Increase North Dakotans' understanding of the water quality and beneficial use impairments associated with NPS pollution, and strengthen public support for the voluntary implementation of NPS pollution control activities.

A variety of educational efforts are supported by the NPS Program to increase public awareness of NPS pollution issues as well as to strengthen support for current and future NPS pollution management projects. These educational efforts can include activities such as workshops, demonstrations, tours, fact sheets, radio ads and videos. Generally, the information/education (I/E) efforts are sponsored and implemented by SCDs, resource conservation and development councils or the NDSU Extension Service. Although the goals and target audiences of the educational projects may vary, these state/locally sponsored I/E projects cumulatively form a balanced statewide NPS pollution education program. Specific I/E projects supported under the 2003, 2006 and 2007 Grants are listed in Table III-2.

On an annual basis, NPS Program staff members are also involved in numerous educational events. These efforts can include presentations at local tours and workshops, display booths at county fairs and agricultural shows, instruction at ECO ED camps, assistance with Envirothon competitions, newsletter articles and dissemination of various materials.

Program Evaluation

Evaluation Goal: Evaluate the successes and failures of the NPS Pollution Management Program and identify the necessary updates to the NPS Pollution Management Program to maintain successful delivery of financial and technical assistance to local and state agencies and private organizations addressing NPS pollution.

The overall success of the NPS Program is evaluated at both the state and local levels. At the state level, success is being measured by the degree of progress toward goals set forth in the management plan. Locally, progress toward project-specific goals and objectives will be used to evaluate the accomplishments of the individual projects.

The long-term goal of the NPS Program is to deliver a balanced program focused on the restoration and maintenance of beneficial uses impaired by NPS pollution. The 1998 305(b) Report and Section 303(d) list are the baseline documents that will be used to measure progress toward this goal. Initiation of watershed restoration projects in 75 of the “impaired” watersheds included on the 1998 303(d) list is the main objective associated with the long-term goal. This objective is scheduled to be met by 2013. With 23 NPS assessment and/or TMDL development projects and 31 watershed restoration projects supported under the Active Grants, the NPS Program is on track to initiate 75 watershed restoration projects by the target date. It should be noted, however, that the objective is to initiate the restoration projects by 2013. Past experience has indicated that many of the watershed restoration projects initiated by 2013 may not actually be completed until 2020-2023. Consequently, the full benefits of the watershed restoration efforts may not be realized until 2023 and beyond.

A variety of water quality and land use data are collected annually to document improvements within the NPS watershed project areas. During an average year, approximately 500 water quality samples are collected from STORET sites within the active watershed project areas. The main parameters typically monitored include nitrogen, phosphorus, TSS and fecal coliform bacteria. Stream discharge is also measured at many of the STORET sites to determine pollutant loadings. Upon completion of a project, a summary of the water quality data is developed and incorporated into the final project report. To gauge land use improvements, the number and type of BMPs applied are also tracked by the local NPS projects. Table III-5 lists the amounts and costs of the BMPs applied within the NPS project areas since January 1, 2003. Sixty percent of the total BMP costs listed in Table III-5 were supported with Section 319 funds.

**Table III-5. BMPs Supported Under the Fiscal Year 2003, 2006 and 2007
Grants (1/1/03 - 11/30/07)**

Category/Practice	Amount	Units	Total Cost
<i>Cropland Management</i>			
GPS Equipment (Nutrient Management)	3.00	Number	\$ 5,726.05
Nutrient Management	127,225.90	Acres	\$ 568,041.66
Pasture/Hayland Planting	371.80	Acres	\$ 11,471.53
Pest Management	36,503.20	Acres	\$ 151,851.45
Residue Management (Mulch Till)	52,790.90	Acres	\$ 392,181.68
Residue Management (No-Till and Strip Till)	96,047.70	Acres	\$ 1,098,211.48
Soil Test (Nutrient Management)	36.00	Number	\$ 2,022.20
Subtotal			\$ 2,229,506.05
<i>Erosion Control</i>			
Critical Area Planting	685.40	Acres	\$ 191,108.24
Grade Stabilization	1.00	Number	\$ 2,694.81
Grassed Waterway	550.00	Linear Feet	\$ 13,711.50
Miscellaneous (Erosion Control)	1.00	Misc	\$ 4,228.70
Sediment Basin	2.00	Number	\$ 122,483.34
Water and Sediment Control	2.00	Number	\$ 12,755.00
Subtotal			\$ 346,981.59
<i>Grazing Management</i>			
Alternative Power Source (Livestock Watering Only)	5.00	Number	\$ 27,187.37
Fencing	1,119,754	Linear Feet	\$ 853,743.78
Heavy Use Protection	1.00	Number	\$ 1,400.00
Mechanical Treatment	45.00	Acres	\$ 373.50
Miscellaneous (Grazing Management)	4.00	Misc	\$ 13,621.36
Pasture/Hayland Planting	8,914.80	Acres	\$ 313,584.64
Pipelines	409,896	Linear Feet	\$ 938,718.24
Pond	53.00	Number	\$ 89,847.67
Prescribed Grazing	8,406.70	Acres	\$ 1,600.00
Range Planting	41.90	Acres	\$ 2,259.72
Solar Pumps	4.00	Number	\$ 22,476.46
Spring Development	4.00	Number	\$ 35,525.86
Trough and Tank	229.00	Number	\$ 283,276.93
Use Exclusion	10.00	Acres	\$ 3,321.66
Well (Livestock Only)	59.00	Number	\$ 331,859.55
Subtotal			\$ 2,918,796.74
<i>Livestock Manure Management System (Full System)</i>			
Miscellaneous (Full Manure Management System)	5.00	Misc	\$ 115,087.62
Phase I Waste Management System	37.00	System(s)	\$ 2,209,337.43
Phase II Waste Management System	28.00	System(s)	\$ 1,354,900.38
Phase III Waste Management System	6.00	System(s)	\$ 242,291.58
Waste Management System (Coordinated With EQIP)	24.00	System(s)	\$ 1,294,161.44
Waste Management System (Full System Completed)	21.00	System(s)	\$ 1,527,507.70
Subtotal			\$ 6,743,286.15

**Table III-5 (cont.). BMPs Supported Under the Fiscal Year 2003, 2006 and 2007
Grants (1/1/03 - 11/30/07).**

Category/Practice	Amount	Units	Total Cost
<i>Livestock Manure Management System (Partial Sys)</i>			
Building Relocation, Moving Costs (Ag Waste)	1.00	Number	\$ 40,267.27
Bunk Line Fencing (Ag Waste)	1,920.00	Linear Feet	\$ 4,800.00
Diversion	1,360.00	Linear Feet	\$ 27,267.76
Engineering Services - Construction Phase	1.00	System(s)	\$ 11,192.00
Engineering Services - Preconstruction	3.00	System(s)	\$ 9,067.63
Manure Removal (Ag Waste)	1.00	System(s)	\$ 1,360.00
Miscellaneous (Partial Manure Management System)	1.00	Misc	\$ 3,722.26
Perimeter Fencing (Ag Waste)	10,705.00	Linear Feet	\$ 19,438.80
Runoff Management System	1.00	System(s)	\$ 95,589.38
Site Prep (Ag Waste)	1.00	System(s)	\$ 3,625.00
Soil Test (Ag Waste)	5.00	Number	\$ 4,344.60
Waste Storage Facility	1.00	System(s)	\$ 2,750.00
Waste Utilization	9,232.92	Acres	\$ 198,881.61
Water Supply (Ag Waste)	6.00	Number	\$ 3,000.00
Watering Facility (Ag Waste: Tank, Pipeline, Well)	2.00	Number	\$ 12,667.60
Windbreak Fencing (Ag Waste)	7,386.00	Linear Feet	\$ 14,521.76
Subtotal			\$ 452,495.67
<i>Miscellaneous Practices</i>			
Cultural Resource Review	11.00	Number	\$ 10,284.26
Engineering Services - Construction Phase	6.00	System(s)	\$ 22,388.36
Engineering Services - Post Construction	2.00	System(s)	\$ 14,883.96
Engineering Services - Preconstruction	13.00	System(s)	\$ 58,838.12
Miscellaneous (Full Manure Management System)	1.00	Misc	\$ 7,925.50
Miscellaneous (Miscellaneous Practices)	12.00	Misc	\$ 32,938.11
Septic System Renovation	4.00	System(s)	\$ 24,136.45
Site Preparation - Heavy w/Chemical (Trees, G13)	2.00	Acres	\$ 340.00
Soil Investigations	1.00	Number	\$ 738.70
Solar Pumps	5.00	Number	\$ 15,840.78
Urban Stormwater Management	1.00	System(s)	\$ 268,134.95
Well Decommissioning	27.00	Number	\$ 24,877.96
Subtotal			\$ 481,327.15
<i>Riparian Area Management</i>			
Engineering Services - Construction Phase	1.00	System(s)	\$ 7,906.88
Engineering Services - Preconstruction	3.00	System(s)	\$ 12,320.26
Riparian Forest Buffer	495.62	Acres	\$ 175,966.45
Riparian Herbaceous Cover	18.00	Acres	\$ 18,059.01
Stream Channel Stabilization	44,845.00	Linear Feet	\$ 236,867.81
Streambank and Shoreline Stabilization	15,144.00	Linear Feet	\$ 491,096.43
Timber Stand Improvement (Scarification)	14.80	Acres	\$ 4,864.55
Tree Handplants	1,833.00	Number	\$ 2,233.00
Sub-total			\$ 949,314.39

**Table III-5 (cont.). BMPs Supported Under the Fiscal Year 2003, 2006 and 2007
Grants (1/1/03 - 11/30/07)**

Category/Practice	Amount	Units	Total Cost
<i>Upland Tree Planting</i>			
Cultural Resource Review	1.00	Number	\$ 1,529.27
Mechanical Treatment	3.20	Acres	\$ 64.00
Site Preparation - Heavy w/Chemical (Trees, G13)	32.20	Acres	\$ 901.60
Tree Handplants	2,446.00	Number	\$ 4,196.44
Tree/Shrub Establishment	150,625.34	Linear Feet	\$ 43,881.57
Weed Control For Tree Establishment (Chem or Mech)	32.20	Acres	\$ 615.00
Windbreak/Shelterbelt	131,242.00	Linear Feet	\$ 60,666.31
Subtotal			\$ 111,854.19
<i>Vegetative Buffers</i>			
Filter Strip	48.50	Acres	\$ 6,079.25
Subtotal			\$ 6,079.25
<i>Wetland Restoration/Creation</i>			
Wetland Creation	23.00	Acres	\$ 52,448.94
Wetland Restoration	855.60	Acres	\$ 223,554.27
Subtotal			\$ 276,003.21
Grand Total			\$14,515,644.39

Despite the implementation of multiple BMPs and the collection of extensive water quality and quantity data, documentation of annual pollutant load reductions continues to be very difficult. This is particularly true within the large watershed projects (i.e., greater than 50,000 acres). Given variables such as rainfall timing and amounts and cropping changes, it is anticipated that more than 10 years of data may be needed to accurately document pollutant load reductions within most watershed projects. For the short term, annual pollutant load reductions within some watershed projects are being estimated with the Spreadsheet Tool for the Estimation of Pollutant Load (STEPL) model. In recent years, the Animal Feedlot Runoff Risk Index (AFRRI) worksheet has also been used by projects focused on manure management improvement. The STEPL model provides annual estimates for nitrogen, phosphorus and sediment load reductions associated with crop residue management practices and/or manure management systems. The AFRRI worksheet only provides the estimated nitrogen and phosphorus load reductions associated with manure management systems. Due to the data limitations of both models, the estimated load reductions for other BMPs such as nutrient management and riparian buffers are not calculated or included in any reported load reduction estimates. However, even with these limitations, the results generated by the models still indicate that the local Section 319 projects are having a very positive affect on annual nitrogen, phosphorus and/or sediment loadings. The project-specific load reductions estimated with the models for 2006 and 2007 are listed in Table III-6.

Table III-6. STEPL Estimates - Nitrogen, Phosphorus and Sediment Load Reductions in 2006/2007

Project Name	Nitrogen Load Reduction (lbs/yr)	Phosphorus Load Reduction (lbs/yr)	Sediment Load Reduction (tons/yr)
Lower Pipestem Watershed	3,702	795	75
Cottonwood Creek Watershed	78,199	37,734	2,142
Rocky Run Watershed	12,003	2,526	175
Maple Creek Watershed	68,608	12,672	1,824
Nine Townships Watersheds	1,296	251	117
Beaver Creek Watershed	633	111	41
Powers Lake Watershed	1,068	212	105
Wild Rice River Watershed	5,853	1,291	10
Adams County Livestock Manure Management Program	15,729	3,539	0*
Bear Creek Watershed	6,668	3,301	0*
Dairy Pollution Prevention Program	101,070	38,529	0*
Stockmen's Association - Environmental Services Program	117,815	56,617	0*
Sheyenne River Watershed (Barnes Co.)	1,741	849	0*
Sheyenne River/Dead Colt Watershed (Ransom Co.)	7,518	1,692	0*
Lake Hoskins Watershed	1,105	196	87
Bone Hill Creek Watershed	4,921	2,399	0*
Pheasant Lake Watershed	233	40	15
Buffalo Springs/Lightning Creek Watershed	3,941	1,894	0
Middle Cedar Creek Watershed	9,870	2,248	158
Total	441,973	166,896	4,749

* Livestock manure management systems were the only BMPs installed by these projects. The AFFRI worksheet was used to estimate load reductions. The AFFRI worksheet does not estimate sediment load reductions associated with manure management systems.

Documenting the type and amount of BMPs applied is another valuable measure of project and program success. As indicated in Figure III-3, 39 percent of total Section 319 expenditures under the Active Grants have been associated with the implementation of BMPs. The most common BMPs implemented with this financial support have included cropland nutrient management practices, manure management systems and grazing management practices. The main NPS pollutants addressed by the BMPs include nitrogen, phosphorus, sediment and fecal coliform bacteria. Figure III-4 shows the total BMP expenditures associated with each BMP Category. Specific BMPs implemented since January 1, 2003 are listed in Table III-5.




Figure III-3. Cumulative Cost Category Expenditures Under the Fiscal Year 2003, 2006, and 2007 Grants (1/1/03 –11/30/07)




Figure III-4. BMP Category Expenditures Under the Fiscal Year 2003, 2006 and 2007 Grants (1/1/03 – 11/30/07)

Although it continues to be difficult to measure actual statewide benefits of the NPS Program, the estimates generated by the AFRRI worksheet and STEPL model, as well as the number of applied BMPs, do indicate Section 319 funding is having a positive impact on water quality in the state. Over the long term, as the applied BMPs mature and additional projects are initiated, the actual water quality data collected locally and statewide should begin to reflect reductions in NPS pollution. Continued coordination with USDA and other state, federal and local natural resource agencies will also be a key factor for ensuring measurable reductions in NPS pollution are realized statewide by 2013.

Chapter 4. Total Maximum Daily Load (TMDL) Program

Section 303(d) of the CWA and its accompanying regulations (CFR Part 130, Section 7) require each state to list waterbodies (i.e., lakes, reservoirs, rivers, streams and wetlands) that are considered water quality limited and require load allocations, waste load allocations and total maximum daily loads (TMDLs). This list has become known as the “TMDL list” or “Section 303(d) list.”

A waterbody is considered water quality limited when it is known that its water quality does not or is not expected to meet applicable standards. Waterbodies can be water quality limited due to point source pollution, NPS pollution or both. When a state prepares its list of water quality-limited waterbodies, it is required to prioritize waterbodies for TMDL development and to identify those “High” priority waterbodies that will be targeted for TMDL development within the next two to four years. Factors to be considered when prioritizing waterbodies for TMDL development include: (1) the severity of pollution and the uses which are impaired; (2) the degree of public interest or support for the TMDL, including the likelihood of implementation of the TMDL; (3) recreational, aesthetic and economic importance of the waterbody; (4) the vulnerability or fragility of a particular waterbody as an aquatic habitat, including the presence of threatened or endangered species; (5) immediate programmatic needs, such as waste load allocations needed for permit decisions or load allocations for Section 319 NPS project implementation plans; and (6) national policies and priorities identified by EPA.

After considering each of the six factors, the state has developed a two-tiered priority ranking. Assessment units (AUs) listed as “High” priority are: (1) lakes and reservoirs and river and stream segments for which TMDLs are scheduled to be completed and submitted to EPA in the next two years; or (2) lakes and reservoirs and river and stream segments for which TMDL development projects are scheduled to be started in the next two years. The majority of these “High” priority AUs were identified as such based largely on their degree of public support and interest and the likelihood of implementation of the TMDL once completed. “Low” priority AUs are those river and stream segments and lakes and reservoirs that are scheduled for completion in the next eight years.

The responsibility for TMDL development for Priority 1 and 2 waterbodies in North Dakota lies primarily with the department’s Division of Water Quality - Surface Water Quality Management Program. To facilitate the development of TMDLs, the department created three regional offices located in Fargo, Dickinson and Towner, N.D. (Figure III-5). The focus of the regional TMDL/Watershed Liaison staff is to work with local stakeholders in the development of TMDL water quality assessments and TMDLs based on the 303(d) list. Technical support for TMDL development projects and overall program coordination are provided by Surface Water Quality Management Program staff located in Bismarck, North Dakota.

Typically, TMDL development projects involve monitoring and assessment activities which will:

- Quantify the amount of a pollutant that the impaired water can assimilate and still meet water quality standards.
- Identify all sources of the pollutant contributing to the water quality impairment or threat.
- Calculate the pollutant loading entering the waterbody from each source.

- Calculate the reduction needed in the pollutant load from each source necessary for attainment of water quality standards.

The goals, objectives, tasks and procedures associated with each TMDL development project are described in project-specific Quality Assurance Project Plans.




Figure III-5. Map Depicting Areas of Responsibility for Regional TMDL/Watershed Liaison Staff

Equally as important as the development of TMDLs is their implementation. The regional TMDL liaisons provide technical assistance to local SCDs and water resource boards in the development of NPS pollution management projects that address TMDL-listed waterbodies. The liaisons also provide technical expertise to local stakeholder groups and assist with youth and adult information/education events in their regions.

Chapter 5. Coordination with Other Agencies

North Dakota has two rivers of international significance. The Souris River originates in the Canadian province of Saskatchewan, loops through North Dakota and returns to the province of Manitoba (Figure III-1). The Red River of the North originates at the confluence of the Bois de Sioux and Ottertail Rivers at Wahpeton, North Dakota. The Red River flows north, forming the boundary between North Dakota and Minnesota before entering Manitoba. The department participates in two cross-border cooperative efforts to jointly manage and protect these rivers.

To ensure an ecosystems approach to transboundary water issues and to achieve greater operational efficiencies in the conduct of the International Joint Commission (IJC) and its responsibilities, the IJC has combined the ongoing responsibilities of the International Souris River Board of Control and the Souris River aspects of the International Souris-Red River Engineering Board into the International Souris River Board (ISRB). The ISRB operates under a directive from the IJC dated April 11, 2002. Part of the ISRB's mission is to assist the IJC in preventing and resolving disputes related to the transboundary waters of the Souris River basin.

The other international water quality effort in which the department is involved is the International Red River Board. Created by the International Joint Commission (IJC), the board monitors Red River water quality. The board also informs the IJC of trends and exceedances of water quality objectives, documents discharges and control measures, establishes a spill contingency plan and identifies future water quality issues. Board activities are detailed in annual reports. Other members of the board include Environment Canada, Manitoba Water Stewardship, EPA, USGS, U.S. Bureau of Reclamation and the Minnesota Pollution Control Agency.

The department monitors water quality in Devils Lake and distributes historical and current data to various federal and state agencies. Information and technical expertise is provided to sponsoring agencies that are planning mitigation measures for rising lake levels.

The Red River Basin Commission (RRBC) was formed in 2002 to initiate a grass roots effort to address land and water issues in a basin-wide context. The RRBC was formed as a result of a merger between The Red River Basin Board, The International Coalition and the Red River Water Resources Council.

The RRBC is not intended to replace governmental agencies or local boards that have water management responsibilities in the basin. Rather, it was created to develop a comprehensive plan on a scale never before attempted. Another purpose of the RRBC is to foster the inter-jurisdictional coordination and communication needed to implement such a plan and to resolve disputes that inevitably will arise among varied interests during the planning process.

The RRBC is made up of a 41-member board of directors, comprised of mainly representatives of local government, including the cities, counties, rural municipalities, watershed boards, water resource districts and joint powers boards, as well as representation from First Nations, a water supply cooperative, a lake improvement association and environmental groups. There also are four at-large members. The governors of North Dakota and Minnesota and the premier of the province of Manitoba have also appointed members to the board.

D. Cost/Benefit Assessment

Costs associated with municipal point source pollution control have been extensive. Capital investments in the form of additions to and construction of new wastewater treatment facilities account for the largest expenditure of funds. While the Clean Water State Revolving Fund (CWSRF) and other state and federal programs have been the major sources of funding, many communities have upgraded wastewater treatment facilities at their own expense.

In the last two years, approximately \$53 million has been obligated from the CWSRF for the construction of wastewater system improvements. The cumulative amount invested in wastewater system improvements since passage of the Clean Water Act in 1972 is approximately \$455 million. In addition to the capital costs, an estimated \$20 million per year is spent operating and maintaining wastewater treatment systems in the state.

While the costs of construction and maintenance of municipal wastewater treatment systems are relatively easy to compile, monetary benefits cannot be so easily quantified. Qualitative benefits include the reduction or elimination of waste loads to receiving waters (Figure III-2, page III-6) and the elimination of public health threats such as malfunctioning drain-field systems and sewer backups.

Federal, state and local governments have also made significant investments in NPS pollution controls. Since 1999, the state's Section 319 NPS Pollution Control Program has provided more than \$39 million in financial support to more than 72 state and local projects, including more than \$22.3 million to 32 watershed restoration projects. In addition to the Section 319 investment in these watershed projects, project sponsors have provided more than \$14.8 million in local match to these watershed projects (Table III-2, page III-14). A variety of agricultural and other BMPs have been implemented through these watershed projects (Table III-5, page III-22). Total costs of these BMPs were more than \$14.5 million.

The water quality benefits of these Section 319 NPS Pollution Control Program expenditures can be described through documented watershed reductions in nitrogen, phosphorus and sediment. Using EPA's STEPL model, Section 319 cost-shared BMPs are resulting in significant nitrogen, phosphorus and sediment reductions. Based only on crop residue management practices and manure management systems cost-shared in watershed projects through 2005, it is estimated that nitrogen, phosphorus and sediment loading will be reduced by more than 441,000 pounds, 166,000 pounds and 4,700 tons per year, respectively (Table III-6, page III-25).

E. Special State Concerns and Recommendations

The “watershed approach” is not a new or unique concept in water quality protection programs. The concept of conducting watershed planning and management first arose with Section 208 of the original 1972 Clean Water Act. The watershed approach is also a key element in EPA’s Clean Water Action Plan. This cooperative approach involves state, tribal, federal and local governments and the public identifying the watersheds with the most critical water quality problems and then working together to focus resources and implement effective strategies to solve those problems.

It is the department’s recommendation that a watershed approach be implemented for all of its water quality monitoring, assessment and nonpoint source pollution control programs. The department will continue to work with local governmental entities (e.g., SCDs, water resource boards, county commissions, cities) in the implementation of watershed restoration projects throughout the state.

As the dominant land use in North Dakota, agriculture has been the primary focus of the state’s NPS Pollution Management Program. Over the past seven years, the department has directed a majority of Section 319 funds to projects addressing agricultural NPS pollution (see Part III. C. Chapter 3). Given the magnitude and complexity of the agricultural industry, the department has developed a close working relationship with the NRCS to ensure sufficient resources are available to adequately address NPS pollution within the state. The combined resources from both the Section 319 Program and the NRCS have proven essential for a balanced NPS Pollution Management Program.

To maintain this coordinated effort, continued funding through Section 319 and the NRCS programs will be necessary. While NRCS programs (e.g., Environmental Quality Incentives Program) can provide funding for BMP cost-share, this funding is only available on an annual basis and producers and project sponsors are required to compete for this funding on a statewide basis. Section 319 provides long-term (five- to 10-year) funding to address water quality problems at the watershed scale. Section 319 funding is also used to hire watershed coordinators who are dedicated to the goals of each watershed project. These coordinators are responsible for providing much needed technical assistance to producers in their watersheds, assistance that would not be available through any other funding source.

The state has recently made a significant investment in NPS pollution control. Since 2001, the state has contributed \$800,000 to the Section 319 funded watershed projects. Using state “Water Development Trust” funds either appropriated by the state legislature or obligated by the State Water Commission, these funds have provided a much needed source of the state/local match required by the Section 319 Program. The state should continue to maintain funding to support NPS pollution management projects throughout the state and to explore ways to expand state funding to support these efforts.

Public awareness of environmental issues, along with the trend toward larger, more concentrated livestock operations, has brought increased concern over these operations and their potential impacts to water quality. The department continues to work closely with the NRCS and others to provide assistance to implement approved livestock waste systems. Without consistent funding

from federal programs like the Section 319 NPS Pollution Management Program and the NRCS Environmental Quality Incentives Program that are addressing animal feeding operations, efforts to bring impaired waters into compliance could be greatly hampered.

The department has taken an active approach in implementing its Stormwater Program. The department continues to work with regulated small MS4s and the Red River Work Group on issues relating to stormwater discharges. Runoff from construction stormwater has been a major concern of EPA.

States need flexibility when managing their stormwater management programs so they can find the best fit for their respective conditions. As long as the stormwater requirements are being met and no water quality violations occur in the state, EPA should refrain from program micromanagement. The department also believes that EPA's "one-size-fits all" approach is not the best way to address construction storm water issues. Each state has its own unique set of conditions when it comes to topography, soils and associated BMPs. For example, BMPs that are used on locations with tighter (clay) soils and flat topography may not work in till or sandy soils with steeper slopes. A one-size-fits-all approach that does not recognize these differences can lead to over-regulation and inefficiencies in program implementation.

The department continues to develop and expand its biological assessment program. It is generally believed that the instream biological community (e.g., fish, aquatic insects and algae) exposed to pollutant stresses on a continual basis is the best measure of aquatic life use. In 2005, the department initiated a two-year biological assessment project in the Red River basin using a probabilistic study design. Once completed, this project will provide an unbiased estimate of biological condition in the Red River basin of North Dakota. Data collected as part of this study will also be used to refine existing fish and macroinvertebrate Indices of Biological Integrity. In subsequent years, the department plans to continue its biological assessment program in the Souris, James and Missouri Rivers basins. This plan will only become a reality, however, if supplemental funding for monitoring programs is maintained by Congress and the EPA.

The department has primacy for most Clean Water Act programs. These include the NDPDES Permit Program, Industrial Pretreatment Program, Storm Water Management Program, Animal Waste Management Program, Clean Water State Revolving Loan Fund Program, Source Water Protection Program, Nonpoint Source Pollution Management Program, Total Maximum Daily Load Program, Clean Lakes Program, Surface Water Monitoring Program, Water Quality Standards, Section 401 Certification and Groundwater Monitoring and Assessment. In order to effectively implement these programs, the department relies on federal funding authorized and appropriated by Congress and provided by EPA. Competing federal priorities (e.g., disaster relief and the "War on Terror") have called into question the federal government's commitment to Clean Water Act programs. Recent cuts in EPA grants to states and rescission orders have put a strain on programs that are already suffering from funding shortfalls. If this trend continues and federal funding continues to decline, the state may have to consider returning some low priority CWA programs to the EPA.

Delays in EPA grant awards to the state are also becoming more problematic. It is not unusual for EPA grant awards to take six to eight months from the time of application to when the grant is awarded. These delays ultimately result in delays in implementing on-the-ground projects or

programs. These delays also strain the department's relationships with local project sponsors. EPA needs to find ways to streamline the granting process by providing a consistent and timely funding source for all Clean Water Act programs. These improvements will ultimately lead to better long-term water quality planning and more effective implementation.

The state's water quality standards define the water quality policy of the state which is to protect, maintain and improve the quality of water for use as public and private water supplies; for propagation of wildlife, fish and aquatic life; and for domestic, agricultural, industrial, recreational and other legitimate beneficial uses. These standards identify specific numeric criteria for chemical, biological and physical parameters. The specific numeric standard assigned to each parameter ensures protection of the beneficial uses for that classification. Numeric standards have been established for bacteria, sulfate, chloride, ammonia, numerous trace elements and organic chemicals.

While nutrients and sediment are the two most prevalent pollutants affecting water quality in the state, no specific criteria exist for them in state water quality standards. EPA has developed guidance and is requiring states to develop a strategy or plan for the development of nutrient criteria. In the absence of a state plan, EPA has said it will promulgate nutrient criteria for the states. Through support provided by an EPA Nutrient Criteria grant, the department recently completed its "Nutrient Criteria Development Plan." This plan provides the blueprint for the development of nutrient criteria for the state's rivers, streams, lakes and rivers.

There are currently no consistent methods for the development of "clean" sediment criteria for the nation's rivers and streams. Without specific criteria or standards for sediment, it is difficult, if not impossible, to set TMDL goals for waterbodies impacted by sedimentation. EPA needs to expand efforts to develop technical guidance for the development of sediment criteria. EPA should also continue funding state efforts to implement its "Nutrient Criteria Development Plans" as well as state efforts to develop scientifically defensible "clean" sediment criteria.

Appropriation of water for consumptive use reduces river flows and subsequently contributes to impaired water quality. Water quality and water quantity are inextricably linked. Reduction in flow reduces the dilution potential and limits the assimilative capacity of the river or stream. Current state appropriation policy contributes to an increasing challenge to meet ambient water quality criteria. The increase in the number of impaired and threatened waterbodies suggests a link to reduced flows. Changes in the natural flow regime of rivers and streams through water withdrawals can also negatively affect instream habitat for fish and other aquatic biota and the aquatic food web.

In North Dakota, a large portion of the potable groundwater resource underlies agricultural areas. The department, in conjunction with the State Water Commission, is involved in several projects designed to evaluate and monitor the effects of agricultural practices on groundwater quality and quantity. The department also reviews water appropriation permits to assess potential impacts to groundwater quality. The department will need to allocate sufficient resources to continue providing project oversight and monitoring, reviewing appropriation permits and working with producers regarding irrigation and chemigation practices to protect groundwater resources.

Careful attention must be paid to the water quality and supply issues associated with the continued energy development, for example, in-situ fossil fuel recovery (oil and coal bed methane development) and the production of ethanol and biodiesel. Sufficient resources must be allocated to avoid impacts to water quality.

Certain areas of the state have experienced increased population growth, and additional funds and resources will be required to ensure waters of the state are protected in populous areas.

The North Dakota Department of Health continues its work to maintain and improve surface and ground water quality in the state. It has taken considerable funding, time and dedication to protect water quality from point and nonpoint sources. For example, more than \$100 million will be spent by North Dakota's three largest cities in the next four years to maintain secondary treatment of wastewater. An additional \$5 million is spent annually on NPS projects, and intensive, annual monitoring continues on the state's most vulnerable aquifers. To maintain this level of effort, both state and federal funding must be continued at current or increased levels.

While efforts to protect water quality have been successful, more remains to be done to achieve the goal of restoring and maintaining the chemical, physical and biological integrity of the state's and nation's waters.

PART IV. SURFACE WATER MONITORING AND ASSESSMENT METHODOLOGY

A. Surface Water Quality Monitoring Program

Chapter 1. Monitoring Goals and Objectives

North Dakota's surface water quality monitoring program is detailed in a report entitled *North Dakota's Water Quality Monitoring Strategy for Surface Waters: 2005-2014* (NDDoH, 2005). This document describes the department's strategy to monitor and assess its surface water resources, including rivers and streams, lakes and reservoirs and wetlands. This strategy also fulfills requirements of Clean Water Act Section 106(e)(1) that requires the EPA, prior to awarding a Section 106 grant to a state, to determine that the state is monitoring the quality of its waters, compiling and analyzing data on the quality of its waters and including those data in its Section 305(b) report. An EPA guidance document entitled *Elements of a State Water Monitoring and Assessment Program* (EPA, March 2003) outlines 10 key elements of a state monitoring program necessary to meet the prerequisites of the CWA. The 10 key elements are:

- Monitoring Program Strategy
- Monitoring Objectives
- Monitoring Design
- Core and Supplemental Water Quality Indicators
- Quality Assurance
- Data Management
- Data Analysis/Assessment
- Reporting
- Programmatic Evaluation
- General Support and Infrastructure Planning

The department's water quality monitoring goal for surface waters is ***“to develop and implement monitoring and assessment programs that will provide representative data of sufficient spatial coverage and of known precision and accuracy that will permit the assessment, restoration and protection of the quality of all the state's waters.”*** In support of this goal and the water quality goals of the state and of the Clean Water Act, the department has established 10 monitoring and assessment objectives. The following objectives have been established to meet the goals of this strategy. They are:

- Provide data to establish, review and revise water quality standards.
- Assess water quality status and trends.
- Determine beneficial use support status.
- Identify impaired waters.
- Identify causes and sources of water quality impairments.
- Provide support for the implementation of new water management programs and for the modification of existing programs.
- Identify and characterize existing and emerging problems.
- Evaluate program effectiveness.
- Respond to complaints and emergencies.
- Identify and characterize reference conditions.

Chapter 2. Monitoring Programs, Projects and Studies

In order to meet the goals and objectives outlined above, the department has taken an approach which integrates several monitoring designs, both spatially and temporally. Monitoring programs include fixed station sites, stratified random sites, rotating basin designs, statewide networks, chemical parameters and biological attributes. In some cases, department staff members conduct the monitoring, while in other instances monitoring activities are contracted to other agencies such as soil conservation districts, the USGS or private consultants. In the following sections, current monitoring activities are documented in the form of narrative descriptions. These include the project or program purpose (objectives), monitoring design (selection of monitoring sites), selected parameters and the frequency of sample collection.

Ambient Water Quality Monitoring Network for Rivers and Streams

The department's Ambient Water Quality Monitoring Network for Rivers and Streams was established in the 1960s. The primary purpose of this network is to provide data for trend analysis, general water quality characterization and pollutant loading calculations. Although the network has undergone several modifications since that time, the network currently consists of 34 fixed-station ambient monitoring sites located on 19 rivers (Table IV-1 and Figure IV-1). Sites are both wadeable and non-wadeable. Where practical, these sites are co-located with USGS flow-gauging stations. Samples are collected and analyzed for water chemistry and bacteria at each of these sites every six weeks during the open-water period (generally from early April through November) and once during the winter under ice cover (generally in late January or early February). Parameters include major ions, trace elements, total suspended solids, total and dissolved nutrients (phosphorus, nitrogen, ammonia, nitrate-nitrite, Total Kjeldahl Nitrogen), total and dissolved organic carbon, and fecal coliform and E. coli bacteria, (Table IV-2). Field measurements are taken for dissolved oxygen (DO), temperature, conductivity and pH.

Through a cooperative agreement with the USGS, a new component was added to the network in September 2003 and May 2007. Equipment was installed at the USGS gauging stations at Fargo (USGS site 05054000; NDDoH site 385414) and Grand Forks (USGS site 05082500; NDDoH site 384156) that monitors field parameters continuously. Data are collected through the deployment of a continuous recording YSI Model 600 multi-probe sonde and datalogger. Output from the sonde is transmitted via telemetry and the data posted "real-time" on the USGS North Dakota district web site. The USGS is also collecting water quality samples 10 times per year from these sites that are analyzed for major cations and anions, total suspended solids (TSS), total phosphorus, total nitrogen, ammonia, nitrate-nitrite and fecal coliform bacteria. As this data set has increased, regression relationships have been developed for select water quality variables (e.g., TSS, TDS, total phosphorus and total nitrogen) using the continuously recorded field parameters. These regression relationships have now been used to provide "real-time" concentration estimates of TSS, total phosphorus, total nitrogen and TDS that are posted on the USGS North Dakota District web site (<http://nd.water.usgs.gov>). As the data set increases for the Grand Forks site, regression relationships will be developed and "real-time" concentration estimates provided for this site as well.

Table IV-1. Ambient Water Quality Monitoring Network Sites.

Station ID	River	Location
380161	Souris River	above Minot
380021	Des Lacs River	at Foxholm
380095	Souris River	at Verendrye
385055	Bois de Sioux	near Doran, MN
380083	Red River	at Brushville, MN
380031	Wild Rice River	near Abercrombie
385414 ^{1,2}	Red River	at Fargo
385040	Red River	near Harwood
380010	Sheyenne River	at Warwick
380009	Sheyenne River	3 mi E of Cooperstown
380153	Sheyenne River	below Baldhill Dam
380007	Sheyenne River	at Lisbon
385001	Sheyenne River	near Kindred
384155	Maple River	at Mapleton
380156	Goose River	at Hillsboro
384156 ^{1,2}	Red River	at Grand Forks
380037 ¹	Turtle River	at Manvel
380039 ¹	Forest River	at Minto

¹ Sampling conducted by the USGS through a cooperative agreement

² USGS “real-time” station

Table IV-1 (cont.). Ambient Water Quality Monitoring Network Sites.

Station ID	River	Location
380157 ¹	Park River	at Grafton
380158 ¹	Pembina River	at Neche
384157 ¹	Red River	at Pembina
384130	James River	at Grace City
380013	James River	at Jamestown
380012	James River	at LaMoure
380022	Little Missouri River	at Medora
380059	Little Missouri River	S of Watford City on Hwy 85 bridge
384131	Knife River	near Golden Valley
380060	Spring Creek	at Zap
380087	Knife River	at Hazen
380160	Heart River	above Lake Tschida
380151	Heart River	near Mandan
380077	Cedar Creek	at Raleigh
380105	Cannonball River	near Raleigh
380067	Cannonball River	S of Breien

¹ Sampling conducted by the USGS through a cooperative agreement




Figure IV-1. Ambient Water Quality Monitoring Network for Sites Rivers and Streams.

Table IV-2. Ambient Water Quality Monitoring Parameters.

Field Measurements	Laboratory Analysis			
	General Chemistry	Trace Elements	Nutrients	Biological
Temperature	Sodium	Aluminum	Ammonia, total	Fecal coliform
pH	Magnesium	Antimony	Nitrate-nitrite, total	E. coli
DO	Potassium	Arsenic	Kjeldahl Nitrogen, total	Enterococcus sp.
Specific Conductance	Calcium	Barium	Nitrogen, total	
	Manganese	Beryllium	Phosphorus, total	
	Iron	Boron	Organic Carbon, total	
	Chloride	Cadmium	Ammonia, dissolved	
	Sulfate	Chromium	Nitrate-nitrite, dissolved	
	Carbonate	Copper	Kjeldahl Nitrogen, dissolved	
	Bicarbonate	Lead	Nitrogen, dissolved	
	Hydroxide	Nickel	Phosphorus, dissolved	
	Alkalinity	Silver	Organic carbon, dissolved	
	Hardness	Selenium		
	TDS	Thallium		
	TSS	Zinc		

Biological Monitoring Program

Historic Program

In response to a recognized need for more and better water quality assessment information, the department initiated a biological monitoring program in 1993. This initial program, a cooperative effort with the Minnesota Pollution Control Agency and the USGS's Red River National Water Quality Assessment Program, was conducted in 1993 and 1994 and involved approximately 100 sites in the Red River Basin. The result of this initial program was the development of the index of biological integrity (IBI) for fish in the Red River Basin. This program continued in the Red River Basin in 1995 and 1996 with the sampling of an additional 100-plus biological monitoring sites. The Upper Red River Basin, including the Sheyenne River and its tributaries, was sampled in 1995, while the Lower Red River Basin was sampled in 1996. From this initial work the program expanded to the Souris River Basin in 1997, the James River Basin in 1998 and the Missouri River Basin in 1999 and 2000. Beginning in 1995, biological monitoring was expanded to include macroinvertebrate sampling in addition to fish. The purpose of this biological monitoring program was to (1) develop an IBI for fish and

macroinvertebrates and (2) provide an assessment of aquatic life use attainment for those stream reaches that were assessed.

Environmental Monitoring and Assessment Program Western Pilot Project

The rotating basin monitoring program was discontinued in 2001 while the department focused its resources in support of sampling for EPA's Environmental Monitoring and Assessment Program (EMAP) Western Pilot Project. The EMAP Western Pilot Project was the second regional pilot project within EMAP focusing on multiple resources. The first of these regional pilot projects focused on the mid-Atlantic region (Maryland, Delaware, Pennsylvania, Virginia and West Virginia). The EMAP Western Pilot Project was a five-year effort (2000-2004) targeted for the western conterminous United States. The pilot involved three EPA Regions (VIII, IX and X) and 12 states (North Dakota, South Dakota, Montana, Wyoming, Colorado, Utah, Arizona, Nevada, Idaho, California, Washington and Oregon). The purpose of the EMAP Western Pilot Project was to: (1) develop the monitoring tools (e.g., biological indicators, stream survey design methods and description[s] of reference condition) necessary to produce unbiased estimates of the ecological condition of rivers and streams that are applicable for the west; and (2) demonstrate those tools in assessments of ecological condition of rivers and streams across multiple geographic regions in the west. In addition to state- and regional-specific assessment questions, the goal of the EMAP Western Pilot's Surface Water Project is to provide answers to three general assessment questions: (1) What proportion of the perennial river and stream miles in the western United States are in acceptable (or poor) biological condition? (2) What is the relative importance of potential stressors (e.g., habitat modification, sedimentation, nutrients, temperature, toxic contaminants, grazing, urbanization) in rivers and streams across the west? (3) What are the stressors associated with the perennial rivers and streams in poor condition? In addition to answering these questions for the western 12-state region of the United States, the EMAP sampling design will allow these questions to be answered in each of the three EPA regions in the west, in each participating state and in several more spatially-intensive "focus areas" in each region. Within North Dakota, these areas are the Upper Missouri River Basin and the Northern Glaciated Plains Ecoregion.

Field sampling for the project began in 2000 and continued through 2003. Based on the EMAP study design, 64 probability-based sites (representing 4,278 perennial stream miles) were sampled within the state. Sites were chosen by EMAP staff based on a random site-selection process. By randomly selecting sites, results can be extrapolated to the entire resource population of concern (in this case, all perennial rivers and streams in the west, EPA Region VIII, North Dakota, the Missouri River Basin and the Northern Glaciated Plains Ecoregion). In addition to the 64 random sites, an additional 47 sites were chosen as targeted "reference" and "trashed" sites. Reference sites exemplify river and stream reaches that are considered "least impaired" with respect to anthropogenic (human) disturbance or stress, while "trashed" sites are believed to be impaired due to one or more anthropogenic stressors (e.g., nutrients, habitat, toxics).

Results of the EMAP Western Pilot Project for North Dakota, along with all of the other states in the region, have been summarized in a report that will be published by EPA Region 8.. These results have also been summarized in this report (see Part V. Rivers and Streams Water Quality Assessment, Chapter 3. EMAP Western Pilot Project Results for North Dakota).

Current Program

Beginning in the spring of 2005 through 2007, the department conducted a biological monitoring and assessment project in the Red River Basin. This project was a joint effort with the Minnesota Pollution Control Agency which sampled the Minnesota side of the Red River Basin. The purposes of this project are to: (1) assess (using biological, physical and chemical data) the current biological condition of perennial, Wadeable rivers and streams in the North Dakota and Minnesota portions of the Red River basin; (2) assess the current status of aquatic life use attainment of the perennial, Wadeable streams of the Red River basin; (3) develop and refine indices of biological integrity for the fish and macroinvertebrate communities; and (4) investigate potential stressors to impaired aquatic life uses.

Sampling consisted of macroinvertebrates, fish, physical habitat and water chemistry. Sampling in 2005 was limited to the Lake Agassiz Plain ecoregion; however, due to above normal precipitation in June and July 2005, only nine sites (three reference and six probabilistic) were sampled for fish and physical habitat. A total of 41 sites (eight reference, nine trashed, eight duplicate Minnesota and 16 probabilistic) were sampled for macroinvertebrates in September 2005. Due, in part, to delays in securing the state FY05 supplemental grant carry-over funds and to staffing shortages caused by untimely employee resignations, sampling was again limited in 2006. Fish were not collected in 2006, and only 17 sites were sampled in the Northern Glaciated Plains ecoregion for macroinvertebrates. All sampling activities were completed in 2007. In the Lake Agassiz Plain ecoregion, a total of 24 random, 10 targeted reference and 10 targeted impaired sites were sampled for the fish indicator. A total of 25 random, 10 targeted reference and 10 targeted impaired sites were visited for the macroinvertebrate indicator in the Lake Agassiz Plain ecoregion. Within-year and among-year replicate samples were also collected as a measure of variability. In the Northern Glaciated Plains ecoregion, field sampling was conducted only for macroinvertebrates. A total of 25 random, 10 targeted reference and 10 targeted impaired sites were sampled for macroinvertebrates. Within-year and among-year samples were once again collected as a measure of variability. Fish were not sampled in this ecoregion.

National Rivers and Streams Assessment

In 2008 and 2009, the department will be participating in the EPA-sponsored National Rivers and Streams Assessment (NRSA). The NRSA is a probabilistic assessment of the condition of the nation's rivers and streams and is designed to:

- Assess the condition of the nation's rivers and streams.
- Establish a baseline to compare future rivers and streams surveys for trends assessments.
- Evaluate changes in condition from the 2004 Wadeable Streams Assessment.
- Help build state and tribal capacity for monitoring and assessment and promote collaboration across jurisdictional boundaries.

The NRSA is one in a series of water assessments being conducted by states, tribes, the EPA and other partners. In addition to rivers and streams, the water assessments will also focus on coastal waters, lakes and wetlands in a revolving sequence. The purpose of these assessments is to

generate statistically valid reports on the condition of our nation's water resources and identify key stressors to these systems.

The goal of the NRSA is to address two key questions about the quality of the nation's rivers and streams:

- What percent of the nation's rivers and streams are in good, fair and poor condition for key indicators of water quality, ecological health and recreation?
- What is the relative importance of key stressors such as nutrients and pathogens?

The NRSA is designed to be completed during the index period of late May through September. Field crews will collect a variety of measurements and samples from predetermined sampling reaches (located with an assigned set of coordinates) and from randomly selected stations along the sampling reach. The field crews will also document the physical habitat conditions along the sampling reach.

Reference Site Network Sampling

The department is currently in the process of implementing its "Nutrient Criteria Development Strategy." This strategy, completed in June 2007, forms the blueprint for the development of nutrient criteria for the state's rivers, streams, lakes, reservoirs and wetlands. One outcome of this strategy is a "reference" condition approach to the development of nutrient criteria for the state's rivers and streams. The "reference" condition approach is also a key component of the state's biological monitoring and assessment program. The purpose of this project is to establish a core set of reference sites throughout the state that can be used to support nutrient criteria development and biological indicators used in the state's bioassessment program. Within each level III ecoregion in the state, a minimum of 10 "reference" or "least impaired" sites will be selected and sampled. In addition, a set of "impaired" or "trashed" sites will also be selected and sampled in each ecoregion. Biological indicators and nutrient criteria will be tested by comparing the results from the "trashed" sites to the "reference" sites. Each site will be sampled once for fish, periphyton and/or macroinvertebrates. Water chemistry variables, including nutrients, will also be sampled at each site. As part of the Red River Basin Biological Assessment Project, "reference" and "trashed" sites were sampled in the Lake Agassiz Plain and Northern Glaciated Plains ecoregions in 2005, 2006 and 2007. In 2008, an additional 30 "reference" and "trashed" sites will be sampled in the Northern Glaciated Plains, Northwestern Glaciated Plains and Northwestern Great Plains ecoregions.

Lake Water Quality Assessment Program

Historic Program

The department currently recognizes 247 lakes and reservoirs for water quality assessment purposes. Of this total, 138 are manmade reservoirs and 109 are natural lakes. All lakes and reservoirs included in this assessment are considered significantly publicly owned.

Reservoirs are defined as waterbodies formed as a result of dams or dugouts constructed on natural or manmade drainages. Natural lakes are waterbodies having natural lake basins. A

natural lake can be enhanced with outlet control structures, diversions or dredging. Based on the state's ADB, the 137 reservoirs have an areal surface of 543,156 acres. Reservoirs comprise about 71 percent of North Dakota's total lake/reservoir surface acres. Of these, 480,731 acres or 62 percent of the state's entire lake and reservoir acres are contained within the two mainstem Missouri River reservoirs (Lake Sakakawea and Lake Oahe). The remaining 136 reservoirs share 62,425 acres, with an average surface area of 459 acres.

The 109 natural lakes in North Dakota cover 218,616 acres, with approximately 117,697 acres¹ or 54 percent attributed to Devils Lake. The remaining 108 lakes average 934 acres, with half being smaller than 250 acres.

Through a grant from the U.S. EPA Clean Lakes Program, the department initiated the Lake Water Quality Assessment (LWQA) Project from 1991-1996. During that time, the department completed sampling and analysis for 111 lakes and reservoirs in the state. The objective of the assessment project was to describe the general physical and chemical condition of the state's lakes and reservoirs, including trophic status.

The lakes and reservoirs targeted for assessment were chosen in conjunction with the North Dakota Game and Fish Department (NDGF). Criteria used during the selection process were geographic distribution, local and regional significance, fishing and recreational potential and relative trophic condition. Lakes without much historical monitoring information were given the highest priority.

The results from the LWQA Project were prepared in a functional atlas-type format. Each lake report discusses the general description of the waterbody, general water quality characteristics, plant and phytoplankton diversity, trophic status estimates and watershed condition.

From 1997-2000, LWQA Project activities were integrated into the department's rotating basin monitoring strategy. Lake Darling and the Upper Des Lacs Reservoir were sampled in 1997 as the department focused its monitoring activities in the Souris River Basin. Pipestem Dam and Jamestown Reservoir were sampled in 1998; Lake Sakakawea was sampled in 1999; and Bowman-Haley Reservoir, Patterson Lake and Lake Tschida were sampled in 2000.

Devils Lake and Lake Sakakawea Monitoring

In addition to inclusion in the annual LWQA Project, Devils Lake and Lake Sakakawea have received special attention. Devils Lake has increased in elevation 26 feet since 1993. In response to questions about water quality changes resulting from these water level increases, the department initiated a comprehensive water quality monitoring program in 1993 for Devils Lake. Devils Lake is currently sampled four times per year, including once during the winter.

While Devils Lake has increased in elevation over the last 10 years, Lake Sakakawea's lake level has dropped significantly since 2002. This drop has been due to drought conditions in the upper Missouri River Basin of Montana resulting in reduced runoff and by the U.S. Army Corps of

¹ The estimated surface area for Devils Lake is based on a lake elevation of 1446 mean sea level (msl), which is the elevation water overflows to Stump Lake.

Engineers' operating policies, which favor downstream navigation interests over the health and condition of the upper Missouri River reservoirs. Of particular concern in North Dakota is the quality of Lake Sakakawea's cold water fishery. Since 2002, the department and the NDGF have cooperated in a project to monitor the condition of the lake. Sampling consists of weekly DO/temperature profiles and water quality samples collected once each month at seven locations.

While not a significant component of the state's lake assessment program, the department also cooperates and assists lake associations and citizen groups with volunteer lake monitoring and assessment projects. When a group or association requests assistance, department staff will meet with the group to define the overall goals and objectives of the project. Based on these goals and objectives, the department will prepare a sampling plan and provide training in sampling methods. The group is responsible for day-to-day monitoring activities, and the department provides laboratory analysis of all samples collected.

NDGF Cooperative District LWQA Project

Many of the lake/reservoir assessments conducted as part of the LWQA Project are now nine to 15 years old. Since that time, there has been a severe drought and significant statewide flooding, both of which may have affected water quality. These climatic factors, along with normal eutrophication, make the assessments conducted as part of the LWQA Project highly questionable.

Working cooperatively with the NDGF Fisheries Division, the department reinitiated a targeted, statewide LWQA Program in 2005. Through this program, 60 lakes and reservoirs were sampled in 2005, ten in 2006 and six in 2007. Samples were collected at least twice during the summer (May/June, July/August or September/October) and once during the winter. The purposes of this project are to: (1) characterize general water quality conditions; (2) assess trophic conditions; (3) determine trends; and (4) assess whether beneficial uses are being met. The results from this project are being summarized in short reports for each lake or reservoir.

Survey of the Nation's Lakes

In 2007, the U.S. EPA, in partnership with the department and other state agencies, initiated the Survey of the Nation's Lakes to answer key environmental questions about the quality of the nation's lakes. The survey will provide a snapshot of the condition of our nation's lake resource on a broad geographic scale. Results from this assessment will allow water quality managers, the public, state agencies and others to say, with known statistical confidence, what proportion of the nation's lakes are in poor biological condition and identify key stressors affecting this resource. Data collected from the lakes will be analyzed on both a regional and national scale. The information generated from this survey fills an important gap in meeting the requirements of the Clean Water Act. The goals of the lakes survey are to:

- Provide regional and national estimates of the condition of lakes in good, fair and poor condition.
- Explore the relative importance of key stressors such as nutrients and pathogens and their extent across the population.
- Establish a baseline to compare future surveys for trends assessment and to evaluate trends since the 1970's National Eutrophication Study.

- Help build state and tribal capacity for monitoring and assessment.

To answer these questions and to achieve the goals of the program, the lakes survey focused on identifying and measuring relevant lake quality indicators in three basic categories: ecological integrity, trophic status and recreational condition. Data collected on stressors will be analyzed to explore associations between stressors and ecological condition.

For the purposes of this survey, lakes are defined as natural or manmade freshwater lakes, ponds and reservoirs in the conterminous U.S. Additional criteria included lake size greater than 10 acres (4 hectares), lake depth greater than 1 meter, and lake area greater than 1000 square meters of open water. Water bodies that were excluded include the Great Lakes (surveyed as part of the National Coastal Condition Assessment), the Great Salt Lake and other naturally saline systems, and water treatment or disposal ponds.

The lake sampling locations were selected using a modern probabilistic survey design approach. In North Dakota, the department, working in cooperation with the USGS, conducted lake sampling at 38 lakes. Four of the state's 38 lakes were replicate sampled for a total of 42 lakes sampled in North Dakota in 2007.

Fish Tissue Contaminant Surveillance Program

Program Background

The purpose of the Fish Tissue Surveillance Program is to protect human health by monitoring and assessing the levels of commonly found toxic compounds in fish from the state's lakes, reservoirs and rivers. The department has maintained an active fish tissue monitoring and contaminant surveillance program since 1990. As part of this program, individual fish tissue samples are collected from selected lakes, reservoirs and rivers throughout the state and analyzed for methyl-mercury. For example, in 2006 and 2007, the department cooperated with the U.S. Fish and Wildlife Service in the collection and analysis of 665 fish tissue samples from four lakes and reservoirs.

These data are then used to issue periodic species-specific fish advisories for the state's rivers, lakes and reservoirs based on risk-based consumption levels. The approach compares the estimated average daily exposure dose for specific waterbodies and species to EPA's recommended reference dose (RfD) for methyl-mercury. Using these relationships, fish tissue data are interpreted by determining the consumption rate (e.g., two meals per week, one meal per week or one meal per month) that would likely pose a health threat to the general population and to sensitive populations (i.e., children and pregnant or breast-feeding women).

NPS Pollution Management Program Monitoring

Program Background

Since the reauthorization of the Clean Water Act in 1987, the North Dakota NPS Pollution Management Program has used Section 319 funding to support more than 90 local projects throughout the state. While the size, target audience and design of the projects have varied

significantly, they all share the same basic objectives. These common objectives are to: (1) increase public awareness of NPS pollution issues; (2) reduce/prevent the delivery of NPS pollutants to waters of the state; and (3) disseminate information on effective solutions to NPS pollution where it is threatening or impairing uses.

State and local projects currently supported with Section 319 funding essentially include three different types of projects. These project types or categories are: (1) development phase projects; (2) educational projects; and (3) watershed projects. Although most projects clearly fit into one of these categories, there are also several projects which include components from all three categories. A portion of the Section 319 funds awarded to the state have also been used to assess major aquifers in the state as well as promote and implement practices that prevent groundwater contamination.

NPS Development Phase Project Monitoring

Locally sponsored NPS assessment or TMDL development projects continue to be the primary means to determine watershed priorities and to prescribe specific management measures. These local assessments, commonly referred to as “development phase projects,” provide the foundation for watershed implementation projects. The primary purposes of development phase projects are to identify beneficial use impairments or threats to specific waterbodies and to determine the extent to which those threats or impairments are due to NPS pollution.

Work activities during a development phase project generally involve an inventory of existing data and information and supplemental monitoring, as needed, to allow an accurate assessment of the watershed. Through these efforts, the local project sponsors are able to: (1) determine the extent to which beneficial uses are being impaired; (2) identify specific sources and causes of the impairments; (3) establish preliminary pollutant reduction goals or TMDL endpoints; and (4) identify practices or management measures needed to reduce the pollutant sources and restore or maintain the beneficial uses of the waterbody. Development phase projects are generally one to two years in length.

As is the case with TMDL development projects, responsibility for development and implementation of NPS assessment projects lies primarily with the department’s Surface Water Quality Management Program. Regional TMDL development staff members are also responsible for coordinating NPS assessment projects. Technical support for assessment projects and overall program coordination are provided by Surface Water Quality Management Program staff located in Bismarck.

The goals, objectives, tasks and sampling procedures associated with each NPS assessment project are described in project-specific Quality Assurance Project Plans (QAPPs).

NPS Watershed Implementation Project Monitoring

Watershed projects are the most comprehensive projects currently implemented through the NPS Pollution Management Program. These projects are typically long-term in nature (five to 10 years, depending on the size of the watershed and extent of NPS pollution impacts) and are designed to address documented NPS pollution impacts and beneficial use impairments within

approved priority watersheds. Common objectives for a watershed project are to: (1) protect and/or restore impaired beneficial uses through the promotion and voluntary implementation of best management practices (BMPs) that reduce/prevent documented NPS pollution loadings; (2) disseminate information on local NPS pollution concerns and effective solutions; and (3) evaluate the effectiveness of implemented BMPs in meeting the NPS pollutant reduction goals of the project.

To evaluate the water quality improvement effects of BMPs that are implemented as part of a Section 319 NPS watershed restoration project, Surface Water Quality Management Program staff members assist local sponsors with the development and implementation of QAPPs specific to the pollutant reduction goals or TMDL endpoints described in the watershed restoration project implementation plan. Each QAPP developed for a watershed restoration project provides a detailed description of the monitoring goals, objectives, tasks and sampling procedures.

Support Projects and Special Studies

Support projects and special studies are activities that are conducted on an as-needed basis to provide data or information to either answer a specific question or to provide program support.

Special studies provide immediate and in-depth investigations of specific water quality problems or emerging issues and usually involve practical research. In conducting practical research, the Surface Water Quality Management Program may rely on its own staff or may contract with the USGS, academia or private consultants. Examples of special studies projects conducted by the department include:

- Studies to develop nutrient criteria for streams and lakes.
- Time of travel studies, dispersion and reareation studies in support of water quality model development.
- The Lostwood National Wildlife Refuge wetland mercury assessment project.

Support projects are activities conducted or supported by the department that result in products or tools that enhance overall program efficiency or lead to new assessment methods. Examples of support projects conducted or supported by the department include:

- Studies to evaluate or compare monitoring methods.
- The watershed and sub-watershed delineation and digitization project.

Complaint and Fish Kill Investigations

Complaint Investigations

The primary purpose for the investigation of complaints is to determine (1) whether or not an environmental or public health threat exists and (2) the need for corrective action where problems are found. Since customer service is a primary focus of the department, complaint response is a very high priority. When complaints are received by the department, they may be handled by department staff, including staff in other divisions of the Environmental Health Section, or forwarded to one of the local health districts located across the state. Once the

complaint is routed to the appropriate state or local health district staff person, a field investigation is usually conducted. When problems are identified, voluntary correction is obtained in most cases. However, necessary enforcement action can be taken under the state water pollution laws (North Dakota Century Code 61-28) and regulations or under other applicable state or federal laws.

Fish Kill Investigations

Fish mortalities can result from a variety of causes and sources, some natural in origin and some induced by man. It is recognized that response time is all-important in the initial phases of a fish kill investigation. Therefore, persons reporting a fish kill are encouraged to immediately contact the department or the NDGF during normal working hours or Emergency Response through state radio. Once a fish kill is reported, staff members from the department's Surface Water Quality Management Program and/or NDGF are dispatched to investigate. The extent of a fish kill investigation is dependent on the numbers and kinds of fish involved and the resources available at the time for the investigation. Following a decision to investigate, the investigation should continue until a cause is determined or until all known potential causes have been ruled out.

Stream Flow

Stream flow data is critical to the analysis and interpretation of water quality data. Stream flow data are used to calculate critical flow conditions for TMDLs and NDPES permitting, to estimate pollutant loading and to interpret water quality results (e.g., load duration curve analysis). The USGS and agencies of the state of North Dakota have had cooperative agreements for the collection of stream flow records since 1903. During the 2007 water year (October 1, 2006 through September 30, 2007), the USGS cooperated with numerous state, federal and local agencies in the collection and reporting of stream flow data from 117 stream flow-gauging stations.

In addition to the extensive USGS stream flow gauging network, the department conducts flow monitoring at most water quality sites associated with NPS assessment and watershed implementation projects and TMDL development projects. This ensures that flow data is available for load calculations and other data analyses.

B. Assessment Methodology

Chapter 1. Introduction

As stated earlier, for purposes of 2008 Section 305(b) reporting and Section 303(d) listing, EPA encouraged states to submit an integrated report and to follow its integrated reporting guidance (U.S. EPA, 2005). The purpose of this section is to briefly summarize the assessment methodology used in this integrated report. A complete description of the state's assessment methodology for surface waters is provided in Appendix A. In general, the state's assessment methodology is consistent with the state's beneficial use designations defined in the state's water quality standards (NDDoH, 2006). The assessment methodology is also consistent with the department's interpretation of the narrative and numeric criteria described in its state water quality standards (NDDoH, 2006).

Assessments are conducted by comparing all available and existing information for an assessment unit to applicable water quality criteria (narrative and numeric). This information, which is summarized by specific lake, reservoir, river reach or sub-watershed, is integrated as beneficial use assessments that are entered into a water quality assessment "accounting"/database management system developed by EPA. This system, which provides a standard format for water quality assessment and reporting, is termed the Assessment Database Version 2.3.0 (ADB).

Chapter 2. Assessment Database (ADB)

Developed by EPA, the ADB is an Access[®] based "accounting"/database management system that provides a standard format for water quality assessment information. It includes a software program for adding and editing assessment data and transferring assessment data between the personal computer and EPA. Assessment data, as compared to raw monitoring data, describes the overall health or condition of the waterbody by describing beneficial use impairment and, for those waterbodies where beneficial uses are impaired or threatened, the causes and sources of pollution affecting the beneficial use. The ADB also allows the user to track and report on TMDL-listed waters, including their development and approval status. A complete description of the ADB is provided in the "Water Quality Assessment Methodology for North Dakota's Surface Waters" (Appendix A).

North Dakota's ADB for the 2008 assessment cycle contains 1,709 discreet assessment units (AUs) representing 54,607 miles of rivers and streams and 247 lakes and reservoirs. Within the ADB, designated uses are defined for each AU (i.e., river or stream reach, lake or reservoir) based on the state's water quality standards. Each use is then assessed using available chemical, physical and/or biological data.

As part of integrated Section 305(b) and Section 303(d) reporting to EPA, the state also provides a copy of the ADB with the 2008 assessment cycle data. While the Section 303(d) TMDL list in Tables VI-1 through VI-5 provides all Category 5 waterbodies, the listing of all Category 1, 2, 3, 4A, 4B and 4C waterbodies are provided to EPA through the ADB.

Chapter 3. Beneficial Use Designation

Water quality reporting requirements under Sections 305(b) and 303(d) of the CWA require states to assess the extent to which their lakes and reservoirs and rivers and streams are meeting water quality standards applicable to their waters, including beneficial uses as defined in their state water quality standards. In addition to beneficial uses, applicable water quality standards also include narrative and numeric standards and antidegradation policies and procedures. While Section 305(b) requires states and tribes to provide only a statewide water quality summary, Section 303(d) takes this reporting a step further by requiring states to identify and list the individual waterbodies that are not meeting applicable water quality standards and to develop TMDLs for those waters. Both Section 305(b) reporting and Section 303(d) listing accomplish this assessment by determining whether the waterbody or AU is supporting its designated beneficial uses.

Beneficial uses are not arbitrarily assigned to AUs, but rather are assigned based on the *Standards of Quality for Waters of the State* (NDDoH, 2006). These regulations define the protected beneficial uses of the state's rivers, streams, lakes and reservoirs. Six beneficial uses (aquatic life, recreation, drinking water, fish consumption, agriculture, industrial and fish consumption) were assessed for purposes of Section 305(b) reporting and Section 303(d) listing.

All waterbodies or AUs entered into the ADB and, therefore, all stream classes (I, IA, II and III) and all lake classes (1-5) are assigned aquatic life and recreation beneficial uses. All Class I, IA and II rivers and streams and all lakes are assigned the drinking water beneficial use.

While not specifically identified in state standards, fish consumption is protected through both narrative and numeric human health criteria specified in the state's water quality standards. Fish consumption has been assigned to all Class I, IA and II rivers and streams, to those Class III streams known to provide a sport fishery and to all Class 1 through 4 lakes.

Other beneficial uses identified in the state's water quality standards are agriculture (e.g., stock watering and irrigation) and industrial (e.g., washing and cooling). These uses are applicable to all stream classes and, unless available data provide evidence of impairment, are presumed to be fully supporting.

Chapter 4. Sufficient and Credible Data Requirements

Water quality assessments, done for purposes of Section 305(b) assessment and reporting and 303(d) listing, require the department to use only what it considers to be sufficient and credible data. A complete description of the department's "sufficient and credible data requirements" is provided in the "Water Quality Assessment Methodology for North Dakota's Surface Waters" (Appendix A). In general, sufficient and credible data are chemical, physical and biological data that, at a minimum, meet the following criteria:

- Data collection and analysis followed known and documented quality assurance/quality control procedures.
- Water column chemical or biological data are 10 years old or less for rivers and streams

and lakes and reservoirs, unless there is adequate justification to use older data (e.g., land use, watershed, or climatic conditions have not changed). There is no age limit for fish tissue mercury data. Data for all 10 years of the period are not required to make an assessment.

- There are a minimum of 10 chemical samples collected in the 10-year period for rivers and streams. The 10 samples may range from one sample collected in each of 10 years or 10 samples collected all in one year.
- There should be a minimum of two samples collected from lakes or reservoirs during the growing season, May through September. The samples may consist of two samples collected the same year or samples collected in separate years.
- A minimum of five fecal coliform and/or E. coli samples are collected during any calendar month from May through September. The five samples per month may consist of five samples collected during the month in the same year or five samples collected during the same calendar month, but pooled across multiple years (e.g., two samples collected in May 2000, two samples collected in May 2001 and one sample collected in May 2005).
- For all chemical criteria that are expressed as a 30-day arithmetic average (e.g., chloride, sulfate, radium 226 and 228, and boron), a minimum of four daily samples must be collected during any consecutive 30-day period.
- A minimum of two biological samples (fish and/or macroinvertebrate) are necessary in the most recent 10-year period. Samples may be collected from multiple sites within the assessment stream reach, multiple samples collected within the same year, or individual samples collected during multiple years. Samples may consist of a minimum of two fish samples, two macroinvertebrate samples, or one fish and one macroinvertebrate sample.
- There are a minimum of five fish tissue samples per species per lake, reservoir or river that represents the range in size classes present in the waterbody.

Chapter 5. Existing and Available Water Quality Data

River and Stream Assessment Data

Chemical Data

Since 1994, the department has operated a network of 26 to 33 ambient monitoring sites. Where practical, sites are co-located with USGS flow gauging stations, thereby facilitating the analysis of chemical data with stream hydrologic data. All of these sites are established as basin or subbasin integrator sites, where the chemical characteristics measured at each of these sites reflect water quality effects in the entire watershed. It is the department's intention to maintain these as long-term monitoring sites for the purpose of assessing water quality trends and to describe the general chemical character of the state's major river basins.

From 1997 through 1999, the department implemented an intensive survey approach to chemical monitoring and assessment. The approach complemented the ambient water quality monitoring network maintained by the department and other program-monitoring activities (e.g., lake water quality assessments, NPS pollution monitoring and assessment and point-source compliance monitoring). The approach integrated chemical monitoring at targeted sites with biological monitoring at sites throughout the basin. The Souris River Basin, James River Basin and the upper Missouri River Basin were sampled in 1997, 1998 and 1999, respectively.

The department also uses data collected by the USGS. The USGS maintains and operates several water quality monitoring sites that provide data used for assessment purposes. Many of these sites are maintained by the USGS through cooperative agreements with other agencies (e.g., North Dakota State Water Commission, U. S. Bureau of Reclamation and U.S. Army Corps of Engineers), through international agreements (e.g., the Souris River Bilateral Agreement) or with the department itself.

In addition to the current 34-station ambient chemical monitoring network and the intensive basin survey program, the department cooperates with local project sponsors (e.g., soil conservation districts and water resource districts) in small watershed monitoring and assessment projects and in waterbody-specific TMDL development projects. These projects entail intensive water quality monitoring, stream flow measurements, land use assessments and biological assessments. Where lake water quality is a concern, lake monitoring also is included in the sampling and analysis plan. The goal of these small watershed monitoring and assessment projects and TMDL development projects is to estimate pollutant loadings to the lake or stream and, where appropriate, set target load reductions (i.e., TMDLs) necessary to improve beneficial uses (e.g., aquatic life and recreation). Most of these projects are followed by Section 319 NPS Pollution Management Program watershed implementation projects. Water quality data collected through these cooperative efforts also are used in assessment of waterbodies for the Section 305(b) report and the TMDL list.

Based on the department's "credible and sufficient data requirements," only the previous 10 years of water column chemistry data will be used for assessments. Years of record are based on the USGS water year. Water years are from October 1 (or one year) through September 30 of the following year. It should be noted that it is preferable to split the year in the fall when hydrologic conditions are stable, rather than to use calendar years. Data for all 10 years of the period are not required to make an assessment. **For purposes of assessments conducted for 2008 Section 305(b) report and Section 303(d) list, the period of record will be from October 1, 1996 through September 30, 2007.**

Biological Data

In response to the growing need for better water quality assessment information, the department initiated a biological monitoring program in 1993 and 1994. This program, which was a cooperative effort with the Minnesota Pollution Control Agency and the USGS's Red River National Water Quality Assessment Program, involved approximately 100 sites in the Red River Basin. The result of this initial program was the development of the Index of Biotic Integrity (IBI) for fish in the Lake Agassiz Plain ecoregion of the Red River Basin. The program continued in the Red River Basin in 1995 and 1996. The Upper Red River Basin, including the

Sheyenne River and its tributaries, was sampled in 1995, while the Lower Red River Basin was sampled in 1996. Following these initial monitoring efforts in the Red River Basin, biological monitoring was expanded statewide with sampling in the Souris River Basin in 1997, the James River Basin in 1998, the Lake Sakakawea subbasin of the Missouri River Basin in 1999 and the Lake Oahe subbasin of the Missouri River Basin in 2000. Beginning in 1995, biological monitoring was expanded to include macroinvertebrate sampling in addition to fish.

Lake and Reservoir Assessment Data

From 1991 through 1996 the department conducted a Lake Water Quality Assessment (LWQA) Project. During that time, the department completed sampling and analysis for 111 lakes and reservoirs in the state. The objective of the assessment project was to describe the general physical and chemical condition of the state's lakes and reservoirs.

The lakes and reservoirs targeted for assessment were chosen in conjunction with the NDGF. Criteria used during the selection process were geographic distribution, local and regional significance, fishing and recreational potential and relative trophic condition. Lakes without much historical monitoring information were given the highest priority.

The results from the LWQA Project have been prepared in a functional atlas-type format. Each lake report discusses the general description of the waterbody, general water quality characteristics, plant and phytoplankton diversity, trophic status assessments and watershed condition.

One of the most useful measures of lake water quality is trophic condition. Trophic condition is a means of expressing a lake's productivity as compared to other lakes in a district or geographical area. In general, oligotrophic lakes are deep, clear lakes with low primary production, while eutrophic lakes are shallow and contain macrophytes and/or algae. Eutrophic lakes are considered moderately to highly productive.

The trophic condition or status was assessed for each of the lakes and reservoirs included in the LWQA. Accurate trophic status assessments are essential for making sound preservation or improvement recommendations. In order to minimize errors in classification, a multiple indicator approach was initiated.

Beginning in 1997, LWQA Project activities were integrated into the department's rotating basin monitoring strategy. Lake Darling and the Upper Des Lacs Reservoir were sampled as the department focused its monitoring activities in the Souris River Basin in 1997. Pipestem Dam and Jamestown Reservoir were sampled in 1998; Lake Sakakawea was sampled in 1999; and Bowman-Haley Reservoir, Patterson Lake and Lake Tschida were sampled in 2000.

In addition to its inclusion in the annual LWQA Project, Devils Lake and Lake Sakakawea have received special attention. Devils Lake has increased in elevation approximately 25 feet since 1993 and is now spilling over into East and West Stump Lakes. In response to questions regarding water quality changes resulting from these water level increases, the department initiated a comprehensive water quality monitoring program in 1993 for Devils Lake. Devils Lake is sampled approximately four times per year, including once during the winter.

While Devils Lake has increased in elevation during the last 12 years, Lake Sakakawea's lake level has dropped significantly since 2002. This drop has been due to drought conditions in the upper Missouri River Basin of Montana resulting in reduced runoff and by the U.S. Army Corps of Engineers' operating policies, which favor downstream navigation interests over the health and condition of the upper Missouri River reservoirs. Of particular concern in North Dakota is the quality of Lake Sakakawea's cold water fishery. Since 2002, the department and the NDGF have cooperated in a project to monitor the condition of the lake. Sampling consists of weekly DO/temperature profiles and water quality samples collected once each month at seven locations. Beginning in 2003 through 2007, the U.S. Army Corps of Engineers also conducted water quality monitoring at several fixed-station sites on Lake Sakakawea.

Beginning in 2005 and continuing in 2006 and 2007 the department initiated a cooperative Lake Water Quality Assessment Project with the NDGF Fisheries Division. The goal of this long-term monitoring and assessment project is to: (1) monitor the chemical, physical and biological character of the state's lakes and reservoirs; (2) use chemical, physical and biological indicators to assess the current water quality condition and trophic status of monitored lakes and reservoirs; (3) determine spatial differences among lakes and reservoirs; and (4) determine temporal trends in lake water quality by comparing project data to Lake Water Quality Assessment data or other historic water quality data. Assessment information generated from this project will be used by both the NDGF and the North Dakota Department of Health's Division of Water Quality to prioritize lakes, reservoirs and their watersheds for lake maintenance and improvement projects (i.e., Save Our Lakes, Total Maximum Daily Loads, Section 319 Nonpoint Source Pollution Management Program). Samples are collected from each lake or reservoir two to four times per year and are coordinated with existing NDGF district lake sampling activities (e.g., standard adult fish population sampling, summer water quality sampling, fall reproduction sampling and winter water quality sampling). At a minimum, two samples are collected during the year, one during the summer (June, July and/or August) and one during the winter under ice cover (January or February). Sixty lakes within five of the six NDGF districts were targeted for sampling in 2005/2006. Ten lakes were targeted for sampling in 2006/2007, and six lakes were targeted in 2007/2008.

Fish Consumption Use Assessment Data

The department has maintained an active fish tissue monitoring and contaminant surveillance program since 1990. As part of this program, individual fish tissue samples are collected from the state's major lakes, reservoirs and rivers and analyzed for methyl-mercury. These data are then used to issue annual species-specific fish advisories for the state's rivers, lakes and reservoirs. The state's statewide fish consumption advisory applies to all waters known to provide a sport fishery.

Of the three rivers and 15 lakes and reservoirs for which there were sufficient credible methylmercury data, only Devils Lake, Lake Sakakawea, the Missouri River (including Lake Oahe) and the Red River were assessed for the integrated report. Creel survey reports were not available for the other lakes and rivers. Weighted average concentrations for each waterbody are presented in Appendices C-F.

Other Agency/Organization Assessment Data

In addition to the water quality data available through existing department programs and projects and that provided by the USGS, the department also requested data from other agencies and organizations. In a letter dated June 12, 2007, the department requested all readily available and credible data from 23 agencies and organizations believed to have water quality data (Appendix B). In response to this request, the department received no other additional data. While the North Dakota State Water Commission did respond to the request for additional data, it was determined that their data had already been provided to the department by the USGS.

Chapter 6. Beneficial Use Assessment Methodology

The assessment methodology or decision criteria used to assess aquatic life, recreation, drinking water, fish consumption, agricultural, and industrial uses where they are assigned to the state's surface waters is provided in Appendix A. All water quality assessments entered into the ADB for Section 305(b) reporting and Section 303(d) TMDL listing are based on "sufficient and credible" monitoring data. Physical and chemical monitoring data used for these assessments included conventional pollutants (e.g., DO, pH, temperature, ammonia, and fecal coliform and *E. coli* bacteria) and toxic pollutants (e.g., trace elements and pesticides) data collected between October 1, 1996 and September 30, 2006. Biological monitoring data used for this report included fish community data collected by the department from the Red River Basin between 1993 and 1996 and macroinvertebrate community data collected throughout the state between 1995 and 2000. If more than one site occurred within a delineated AU, data from all sites and for all years are pooled for analysis.

Chapter 7. Assessment Categories

Key to integrated reporting is an assessment of all of the state's waters and placement of those waters into one of five assessment categories. Guidance provided by the U.S. EPA (U.S. EPA, 2005) provides for five assessment categories representing varying levels of water quality standards attainment. These assessment categories range from Category 1, where all of a waterbody's designated uses are met, to Category 5, where a pollutant impairs a waterbody and a TMDL is required (Table IV-3). These category determinations are based on consideration of all existing and readily available data and information consistent with the state's assessment methodology (Appendix A).

For purposes of the 2008 Integrated Report and Section 303(d) list of impaired waterbodies needing TMDLs, the department has identified a subcategory to Category 5 waterbodies. This subcategory, termed Subcategory 5A, includes rivers, streams, lakes or reservoirs that were assessed and listed in previous Section 303(d) lists, including the 2006 list, but where the original basis for the assessment decision and associated cause of impairment is questionable.

These Subcategory 5A waterbodies include rivers and streams listed for biological impairments based on only one sample for the entire segment or on samples collected more than 10 years ago, waterbodies listed for sediment/siltation impairments, or lakes and reservoirs where the assessments are based on one sampling event or on data that are greater than 10 years old. These waterbodies will remain on the 2008 Section 303(d) list, but will be targeted for additional monitoring and assessment during the next two to four years.

Table IV-3. Assessment Categories for the Integrated Report.

Assessment Category	Assessment Category Description
Category 1	All of the waterbody's designated uses have been assessed and are met.
Category 2	Some of the waterbody's designated uses are met, but there is insufficient data to determine if remaining designated uses are met.
Category 3	There is insufficient data to determine whether any of the waterbody's designated uses are met.
Category 4	<p>The waterbody is impaired or threatened, but a TMDL is not needed. This category has been further subcategorized as:</p> <ul style="list-style-type: none"> • 4A - waterbodies that are impaired or threatened, but TMDLs needed to restore beneficial uses have been approved or established by EPA. • 4B - waterbodies that are impaired or threatened, but do not require TMDLs because the state can demonstrate "other pollution control requirements (e.g., BMPs) required by local, state or federal authority." (see 40 CFR 130.7[b][1][iii]) are expected to address all waterbody-pollutant combinations and attain all water quality standards in a reasonable period of time. • 4C - waterbodies that are impaired or threatened, but the impairment is not due to a pollutant.
Category 5	<p>The waterbody is impaired or threatened for at least one designated use, and a TMDL is needed.</p> <ul style="list-style-type: none"> • 5A – waterbodies currently listed on the Section 303(d) list, but are targeted for additional monitoring and assessment during the next two to four years.

PART V. SECTION 305(b) WATER QUALITY ASSESSMENT

A. Rivers and Streams Water Quality Assessment

Chapter 1. Assessment Category Summary

In EPA's guidance for preparing the Integrated Report, the states were encouraged to report on their waters based on five assessment categories (Table IV-1). In broad terms, the five assessment categories are as follows:

- Category 1: All designated uses are met.
- Category 2: Some designated uses are met, but there are insufficient data to determine if remaining designated uses are met.
- Category 3: There are insufficient data to determine whether any designated uses are met.
- Category 4: Water is impaired or threatened, but a TMDL is not needed for one of three reasons: (a) a TMDL already has been approved for all pollutants causing impairment; (b) the state can demonstrate that "other pollutant control requirements required by local, state or federal authority" are expected to address all waterbody-pollutant combinations and attain all water quality standards in a reasonable period of time; or (c) the impairment or threat is not due to a pollutant.
- Category 5: The waterbody is impaired or threatened for at least one designated use, and a TMDL is needed.

In addition to these five broad categories, the department has identified a subset of Category 5 waterbodies as Subcategory 5A. This subcategory includes rivers, streams, lakes or reservoirs that were assessed and listed in previous Section 303(d) lists, including the 2006 list, but where the original basis for the assessment decision and associated cause of impairment is questionable. These Subcategory 5A waterbodies include rivers and streams listed for biological impairments based on only one sample for the entire segment or on samples collected more than 10 years ago, waterbodies listed for sediment/siltation impairments, or lakes and reservoirs where the assessments are based on one sampling event or on data that are greater than 10 years old. These waterbodies will remain on the 2008 Section 303(d) list, but they will be targeted for additional monitoring and assessment during the next two to four years.

The ADB that has been submitted to EPA as part of this Integrated Report provides an assessment category for each lake, reservoir, river or stream AU.

Table V-1 provides a summary of the number of river and stream AUs and total miles of rivers and streams in each category that were assessed for this report. One AU, totaling 28.56 miles, was classified as Category 1, meaning all uses were assessed and fully supporting. One-thousand two-hundred fifty (1250) AUs totaling 47,639.62 miles were assessed as Category 2. These are AUs where at least one designated use was assessed as fully supporting, but the other uses were not assessed. In most cases, agriculture and industrial uses were assessed as fully supporting with the remaining aquatic life, recreation and/or municipal water supply uses not

assessed. A total of 17 AUs were assessed as Category 4 where at least one designated use was impaired or threatened, but where a TMDL is not required. Of these, four AUs do not need TMDLs because TMDLs have already been completed and approved by EPA (Category 4A) and 13 AUs do not need a TMDL because the cause of the impairment is not a pollutant (Category 4C). These are typically river and stream reaches where habitat degradation or flow alteration is impairing aquatic life use. A total of 194 AUs (6,615.58 miles) were assessed where at least one beneficial use is impaired and a TMDL is required. These Category 5 AUs are provided in a list in Tables VI-1 through VI-4.

Table V-1. Assessment Category Summary for Rivers and Streams in North Dakota (Miles)

Category	Description	Number AUs	Total Size (miles)
1	All uses met	1	28.56
2	Some uses met, others not assessed	1250	47,640.75
3	No uses assessed	0	0
4A	Some or all uses impaired or threatened, but a TMDL(s) has been approved for all impaired uses.	4	77.28
4B	Some or all uses impaired or threatened, but other pollutant controls will result in water quality standards attainment.	0	0
4C	Some or all uses impaired or threatened, but impairment is not due to a pollutant.	13	246.00
5	Some or all uses impaired or threatened, and a TMDL is required. Includes category 5A waterbodies.	194	6,613.64

Chapter 2. Water Quality Summary

Eighty-four percent (4,004 miles) of the rivers and streams assessed for this report fully support the beneficial use designated as aquatic life (Table V-2). Of the streams assessed as fully supporting aquatic-life use, a little more than 60 percent (2,394 miles) are considered threatened. In other words, if water quality trends continue, the stream may not fully support its use for aquatic life in the future. The remaining 16 percent (762 miles) of rivers and streams assessed for this report were assessed as not supporting aquatic life use (Table V-2).

Table V-2. Individual Use Support Summary for Rivers and Streams in North Dakota (Miles)

Use	Fully Supporting	Fully Supporting but Threatened	Not Supporting	Not Assessed	Insufficient Information for Assessment	Total Size
Aquatic Life	1,610.22	2,393.87	762.20	44,897.01	4,943.74	54,607.04
Fish Consumption	0	0	401.48	3,693.65	0	4,095.13
Recreation	1,535.57	3,421.04	1,660.38	46,461.41	1,528.64	54,607.04
Drinking Water Supply	1,651.78	85.74	0	3,388.20	434.18	5,559.90
Agriculture	54,607.04	0	0	0	0	0
Industrial	54,607.04	0	0	0	0	0

NPS pollution (e.g., siltation/sedimentation and stream habitat loss or degradation) was the primary cause of aquatic life use impairment (Table V-3). Other forms of pollution causing impairment are trace element contamination, flow alteration and oxygen depletion. Organic enrichment creates conditions in the stream that cause dissolved oxygen (DO) to be depleted. Rivers and streams impaired by siltation/sedimentation, organic enrichment, eutrophication due to excess nutrients and habitat alteration also will result in a degradation of the biological community. Typically, species composition will shift from an aquatic community comprised of intolerant species (e.g., mayflies, caddisflies, stoneflies and darters) to an aquatic community dominated by tolerant species (e.g., midges, carp and bullheads).

Table V-3. Impairment Summary for Rivers and Streams in North Dakota

Impairment	Miles
Total Fecal Coliform/E. coli	5,023.27
Physical Habitat Alterations	2,527.00
Sedimentation/Siltation	1,783.11
Biological Indicators	1,290.50
Oxygen Depletion	461.66
Mercury in Fish Tissues	401.48
Flow Alterations	274.25
Nutrients	51.40
Trace Metals in the Water Column	191.10
Total Dissolved Solids/Chloride	35.89
Ammonia	34.14
Non-native Aquatic Plants	5.53

The primary sources of pollutants affecting aquatic life use in the state are cropland erosion and runoff, animal feeding operations and poor grazing management (Table V-4). Poor grazing management includes riparian grazing and season-long grazing, which result in the deterioration of the plant community or cause a shift in the plant community away from native grass and forbe species to non-native invader species. Evidence of poor grazing practices would include cattle trailing, gully erosion, poor water infiltration rates resulting from soil compaction and severe streambank erosion. Other sources linked to aquatic-life use impairment are point-source discharges, urban runoff and hydrologic modifications (e.g., upstream impoundments, low-head dams, channelization, flow regulation and diversion, riparian vegetation removal and wetland drainage) (Table V-4).

Recreation use was assessed on 6,617 miles of rivers and streams in the state. Recreation use was fully supporting, fully supporting but threatened and not supporting on 1,536 miles, 3,421 miles and 1,660 miles, respectively (Table V-2). Fecal coliform bacteria data collected from monitoring stations across the state were the primary indicators of recreation use attainment (see Part IV. B., Chapter 6. “Beneficial Use Assessment Methodology”). For this reason, pathogens (as reflected by fecal coliform bacteria) are the primary cause of recreation use impairment in North Dakota (Table V-3). Other factors affecting the use of the state’s rivers and streams for recreation would be eutrophication from excessive nutrient loading, resulting in nuisance algae and plant growth. The primary sources of fecal coliform bacteria contamination are animal feeding operations and riparian area grazing (Table V-4). Point-source discharges also have been linked to exceedances of the fecal coliform bacteria standard of 200 colonies per 100 milliliters (ml). These exceedances occur when a municipality discharges from its sanitary sewer directly to the receiving stream, bypassing the wastewater treatment facility. These

circumstances generally occur in the spring when flooding problems cause infiltration to the sanitary sewer.

Drinking water supply use is classified for 5,560 miles of rivers and streams in the state. Of the 1,738 miles assessed for this report, only 86 miles (4.9 percent) were assessed as threatened for drinking water supply use (Table V-2). The primary threats are taste and odor problems. While the source of taste and odor has not been specifically identified, potential sources include agricultural field runoff, reservoir releases, wetland drainage and industrial and/or municipal discharges.

A total of 4,095 miles of rivers and streams were identified as capable of supporting a sport fishery from which fish could be used for consumption (Table V-2). The Red River of the North (401.48 miles) and the Missouri River from Garrison Dam to Lake Oahe are the only two rivers listed in the state's fish consumption advisory. Methyl-mercury data collected for these advisories were used, along with fish population estimates provided by the North Dakota Game and Fish Department, to estimate the weighted average methyl-mercury concentration for fish in each of these rivers (see Part IV. B. Chapter 6. "Beneficial Use Assessment Methodology – Fish Consumption Assessment Methodology for Rivers and Lakes," page IV-32 and Appendices B-E). Based on the EPA fish tissue of $0.3 \mu\text{g}$ methyl-mercury/gram of fish tissue, only the Red River of the North was assessed as not supporting fish consumption. While there are many potential sources of methyl-mercury, both anthropogenic and natural, to date there have been no specific causes or sources identified for the mercury present in North Dakota fish (Tables V-3 and V-4).

Table V-4. Impairment Source Summary for Rivers and Streams in North Dakota

Source	Miles
Riparian Grazing	5,156.30
Animal Feeding and Handling Operations	3,439.50
Crop Production (Dryland)	2,576.93
Loss of Riparian Habitat	2,524.20
Stormwater Runoff	885.91
Source Unknown	884.17
Highway and Road Runoff	634.89
Channel Erosion/Incision from Upstream Hydromodifications	584.86
Streambank Modification	582.51
On-site Treatment Systems (Septic Systems)	524.81
Wetland Loss (Drainage/Filling)	449.83
Rangeland/Pastureland Grazing	411.87
Upstream Impoundments	368.04
Channelization	292.41
Natural	264.91
Hydrostructure Flow Regulation/Modification	264.75
Municipal Point Source Discharges	224.64
Land Development	125.30
Industrial Point Source Discharge	79.60
Source Outside State Jurisdiction or Border	59.56
Flow Alteration for Water Diversion	27.15
Dam Construction	13.05
Golf Courses	13.04

Chapter 3. Environmental Monitoring and Assessment Program Western Pilot Project Results

Introduction

The Environmental Monitoring and Assessment Program - Western Pilot Project (EMAP-West) was initiated in North Dakota in 2000 by the EPA in cooperation with the USGS and the department to develop and demonstrate monitoring tools that would be used to produce unbiased estimates of ecological conditions in surface waters of the state. Information from EMAP-West was used to establish baseline biological, chemical and physical habitat condition estimates and can be used to make comparisons with similar condition estimates obtained from future monitoring activities. These baseline condition estimates can also be used to evaluate possible ecological changes associated with regulation, restoration, and conservation practices.

Study Area Description

As mentioned previously, North Dakota is covered by four level III ecoregions as defined by Bryce and others (1998). The ecoregions are the Northwest Glaciated Plains (42), the Northwest Great Plains (43), the Northern Glaciated Plains (46) and the Lake Agassiz Plain (48) (Figure V-1). In a recent ecological assessment of western streams and rivers, EPA indicated that North Dakota exists within two broader ecological areas: (1) the Cultivated Plains including the Northern Glaciated Plains and Lake Agassiz Plain ecoregions and (2) the Rangeland Plains, including the Northwest Glaciated Plains and Northwest Great Plains ecoregions (Stoddard and others, no date).

The Cultivated Plains region (ecoregions 46 and 48) (Figure V-1) is a flat to gently rolling landscape composed of glacial till that has high concentrations of temporary and seasonal wetlands. This area contains landscape features such as the Turtle Mountains, the Prairie Coteau, a former glacial lake basin now occupied by Devils Lake and the Lake Agassiz Plain. The Lake Agassiz Plain was once filled with Glacial Lake Agassiz, the last in a series of glacial lakes to fill the valley in the past three million years (Omernik, 1987). Thick beds of lake sediments created the extreme low relief of the Lake Agassiz Plain. Once covered with tall prairie grass, the Lake Agassiz Plain now has intensive small grain and row crop agriculture (Figure V-1).

The Rangeland Plains region is located in the southwest half of North Dakota (ecoregions 42 and 43) (Figure 1). The eastern portion of this ecological area is dominated by the Coteau du Missouri, a series of glacial moraines left after several glacial advances. Most of the Coteau has little or no integrated drainage but does have numerous wetlands formed mostly by melting glacial ice. The remainder of the Rangeland Plains region is made up of the Northwest Great Plains ecoregion. The landscape in this ecoregion is rolling to hilly with numerous streams but few wetlands or lakes. This area supports shortgrass grazing lands and occasional, intensively-cultivated agricultural lands (Figure V-1). The southwest portion of the Rangeland Plains is dominated by rugged badlands and grasslands (Figure V-1), which includes Theodore Roosevelt National Park and the Little Missouri National Grasslands.

Aside from impacts from the larger cities and towns, the greatest impact to streams in North Dakota likely comes from the agricultural production of both crops and animals. Crop production practices can enhance soil erosion, which permits excess nutrients, some pesticides and

sediments to enter streams. Construction of ditches, especially in the Lake Agassiz Plain ecoregion, has allowed fields to drain faster than would occur naturally. Ditching has likely increased the volume of water and sediment load that enter streams and likely caused alterations of the natural hydrology and geomorphology of streams. Animal production, especially in restricted areas, can produce excess nutrients, oxygen-consuming organic matter, pathogens and sediments that may enter streams.




Figure V-1. Landcover Categories and Ecoregions in North Dakota.

Sampling Site Determination

Probability and hand-picked sampling sites were selected in order to develop indicators and derive threshold values for estimating stream condition. Probability sites were chosen by statistical design and may not have been sampled because of access restrictions or lack of water.

In the entire state of North Dakota, access was denied to four sites by the landowner, and one site was inaccessible. Reference and stressed sampling sites were hand-picked based on anthropogenic land use practices. Reference sites represented streams in a more natural condition with little anthropogenic disturbance. Stressed sites represented streams that were in poor condition because of human or biological impacts that could potentially degrade biotic integrity. Stressors to the streams included chemical, physical and biological components such as excess nutrients, excess river-bank erosion and invasion of non-native species.

Extent of Stream Resource Assessed

A total of 111 sampling sites were selected and sampled in North Dakota (Figure V-3). Forty-one (41) probability sites and 27 hand-picked sites were selected in the Cultivated Plains region, and 23 probability sites and 20 hand-picked sites were selected in the Rangeland Plains region. All 23 probability sites in the Rangeland Plains and 40 of the 41 probability sites in the Cultivated Plains were used in the chemical stressor assessment. All 23 probability sites in the Rangeland Plains and 39 of the 41 probability sites in the Cultivated Plains were used in the physical habitat stressor assessment. Of the approximately 6,900 kilometers (km) of perennial streams within the state, an estimated 6,583 km were represented in the chemical stressor assessment (3,955 km in the Cultivated Plains and 2,627 km in the Rangeland Plains), 6,555 km in the physical habitat stressor assessment (3,928 km in the Cultivated Plains and 2,627 km in the Rangeland Plains), 6,555 km in the macroinvertebrate condition assessment (3,928 km in the Cultivated Plains and 2,627 km in the Rangeland Plains) and 6,583 km in the periphyton condition assessment (2,627 km in the Rangeland Plains and 3,928 km in the Cultivated Plains) (Figure V-2). Approximately 6,116 km of stream length were represented in the statewide assessment of mercury in fish tissue.




Figure V-2. North Dakota Stream Length Assessed.




Figure V-3. EMAP Western Pilot Project Sampling Sites in North Dakota.

Reference Site Determination

The original designation for reference and stressed sites done by EPA was revised by the North Dakota Department of Health using local knowledge and by incorporating its own method. North Dakota's method for defining reference and stressed sites involved defining a set of metrics selected from the landscape, physical habitat and chemical data (Table V-5). Biological data were considered response variables and were not used in the designation of reference and stressed sampling sites. Redundant metrics were eliminated using a correlation matrix. Each metric was standardized using the 5th and 95th percentiles as the floor and ceiling for each metric. The final Reference Index combined all of the final landscape, physical habitat and chemical metrics (Table V-5) into a composite score ranging from 0-100. Sites with Reference Index scores in the top 10 percent were designated as "reference." Sites with scores in the bottom 10 percent were designated as "stressed" for each region.

Table V-5. Landscape, Physical Habitat, and Chemical Metrics Used in the North Dakota Reference Site Index.

Landscape Metrics	Physical Habitat Metrics	Chemical Metrics
Percent Urban Cover	Percent Embeddedness	Total Suspended Solids
Percent Agricultural Cover	Percent Large Woody Debris	Total Phosphorus
Percent Forest Cover	Percent Fines	Ammonia
Percent Wetland Cover	Sinuosity	Sulfate
	Percent Side Channels	Total Nitrogen
	Bank Canopy Cover	Nitrate

Because the reference site selection process which uses the Reference Site Index is only capable of identifying 20 percent of the total sites in each ecoregion as reference or stressed (10 percent reference and 10 percent stressed), this approach may possibly miss some potential site designations. To account for this limitation, adjustments were made to the final reference and stressed site selection process. Sites were evaluated by comparing field sheets, reference scores and consulting the original EPA designations to identify sites as reference or stressed. For example, if the score for a site was just outside the threshold for stressed but the field sheet comments were strongly indicating the site being designated as stressed, the site was designated as stressed. A designation of stressed or reference was also given to a site if both the EPA designation and the field sheets agreed, but the North Dakota Reference Index was inconclusive.

The location of a sampling site in agricultural or grassland areas did not fully determine whether a site was considered stressed. Streams in croplands may have well-preserved riparian zones and have a considerable amount of buffer vegetation and, therefore, could be considered nearly undisturbed. Conversely, streams in grasslands may have extensive activity along the riparian zone from grazing animals or development and could therefore be considered highly disturbed.

In order to accurately display error for each stream km estimate, the statistical program “R” was used to provide the upper and lower limits of each condition class estimate and allowed graphical display of each estimate with confidence. All estimates were calculated using an “R” script written specifically for this assessment.

Condition Indicators

Macroinvertebrate Biotic Integrity

A separate multi-metric index (MMI) was developed for the Cultivated Plains and the Rangeland Plains. The MMI's were developed specific to this dataset and this assessment unit (i.e., North Dakota). The original EPA-ORD plains macroinvertebrate MMI did not perform well at the scale of this assessment unit with the revised reference and stressed site list.

For MMI development in both the Cultivated Plains and the Rangeland Plains regions, datasets were separated into calibration and validation datasets. The first step involved comparing box-plots of individual metrics to show separation between least disturbed (reference) and most disturbed (stressed) sites.

The next step involved submitting the metrics to a redundancy test to determine which metrics were not independently adding value to the MMI. The MMI was created using the most responsive, non-redundant metrics. Additionally, an attempt was made to cover as many ecological categories as possible, which include richness, composition, diversity, tolerance, feeding and habit guild.

The final MMI for the Cultivated Plains region consists of six metrics: percent EPT (Ephemeroptera, Plecoptera, Trichoptera) taxa, percent abundance of individuals that are predators, percent abundance of individuals rated 8 or 9 on the tolerance scale (more tolerant end), percent abundance of individuals rated 6 or 7 on the tolerance scale, percent clinger taxa and percent abundance of individuals that are collector-filterers. There was one composition

metric, two tolerance metrics, two feeding group metrics and one habit metric. These metrics were scored from 0 to 100 based on the range of the dataset, with the six values averaged for the final score. Good and poor condition class thresholds for the Cultivated Plains region were set at the 25th and 5th percentiles of the least disturbed (reference) sites (Table V-6). In the Cultivated Plains region, scores greater than 55.7 were considered good, and scores less than 49.4 were considered poor. Macroinvertebrate MMI scores greater than or equal to 49.4 and less than or equal to 55.7 were considered fair.

Once the final score for each site was calculated, sites were classified with the appropriate condition class according to each biological community. For instance, if the total score for a site in the Cultivated Plains region was 48 (< 49.4), that site is considered to be in poor condition based on the macroinvertebrate indicator.

Table V-6. Multi-metric Index Threshold Values Used to Determine Condition Classes for Biological Indicators in the Cultivated Plains Region of North Dakota.

Biological Community	Condition Class		
	Poor	Fair	Good
Macroinvertebrate MMI	5 th percentile OR <49.4	≥49.4 and ≤55.7	25 th percentile OR >55.7
Aquatic Vertebrate MMI	5 th percentile OR <52.1	≥52.1 and ≤61.5	25 th percentile OR >61.5
Periphyton MMI	10 th percentile OR <21.4	≥21.4 and ≤44.1	25 th percentile OR >44.1

Table V-7. Multi-metric Index Threshold Values Used to Determine Condition Classes for Biological Indicators in the Rangeland Plains Region of North Dakota.

Biological Community	Condition Class		
	Poor	Fair	Good
Macroinvertebrate MMI	10 th percentile OR <22.5	≥22.5 and ≤38.2	25 th percentile OR >38.2
Aquatic Vertebrate MMI	NA	NA	NA
Periphyton MMI	10 th percentile OR <32.3	≥32.3 and ≤42.5	25 th percentile OR >42.5

The final MMI for the Rangeland Plains region also consists of six metrics: EPT richness, percent abundance of individuals that are clingers, percent abundance of individuals that are collector-gatherers, percent predator taxa, percent of taxa rated 0 to 5 on the tolerance scale and percent abundance of individuals rated 8 or 9 on the tolerance scale. There was one richness metric, two tolerance metrics, two feeding group metrics and one habitat metric. These metrics were also scored from 0 to 100 based on the range of the dataset, with the six values averaged for the final score. Good and poor condition class thresholds for the Rangeland Plains were set at the 25th and the 10th percentile of the least disturbed (reference) sites (Table V-7). In the Rangeland Plains region, scores greater than 38.2 were considered good, and scores less than 22.5 were considered poor. Macroinvertebrate MMI scores greater than or equal to 22.5 and less than or equal to 38.2 were considered fair.

Each site visited was scored independently according to region (Cultivated Plains or Rangeland Plains). Thresholds were determined for indicators within each region, and data were then pooled as good, fair or poor for an overall statewide assessment. Thirty-six (36) sites, representing 3,928 km, were used to assess the Cultivated Plains region, while 23 sites, representing 2,627 km, were used to assess the Rangeland Plains region. A total of 6,555 km of

streams were assessed statewide using the macroinvertebrate MMI.

For the state of North Dakota, 3,141 stream km (48 percent) were considered to be in good condition with regard to macroinvertebrate biotic integrity; 1,602 km (24 percent) are estimated to be in fair condition; and 1,813 km (28 percent) are estimated to be in poor condition (Figure V-4).




Figure V-4. Condition Class Summary for Perennial Streams in North Dakota Based on Macroinvertebrate MMI Scores.

Vertebrate Biotic Integrity

For the fish indicator, the MMI developed by EPA-ORD for the EMAP-West assessment (for the entire plains) was used for ecoregions 46 and 48 (Cultivated Plains region of North Dakota (Stoddard et al., 2005). The fish metrics in this MMI are: native rheophilic (prefers running water) species richness; percent abundance of individuals considered super-tolerant; percent abundance of individuals that are nontolerant invertivores or piscivores; sensitive spawner species richness; native catostomid/ictalurid species richness (corrected for stream size); percent abundance of all species that are native, sensitive and migrators; non-tolerant species richness (corrected for stream size); and percent abundance of individuals that are alien (non-native or foreign individuals).

In the Cultivated Plains region, a total of 3,591 out of 4,116 stream km were represented by the probabilistic sample reflecting 87 percent of the perennial stream km present in this region. Thresholds were set using the 25th and the 5th percentile of the reference sites (Table V-6). A score of greater than 61.5 is considered good, while scores ranging between 61.5 and 52.1 are considered fair. Scores less than the fifth percentile, or less than 52.1, are considered to be in

poor condition.

Within the Cultivated Plains region of North Dakota, 1,690 stream km (47 percent) are considered poor, followed by 1,192 stream km (33 percent) estimated to be in fair condition. Only 709 stream km (20 percent) are considered to be in good condition with regard to vertebrate biotic integrity (Figure V-5).




Figure V-5. Condition Class Summary for Perennial Streams in the Cultivated Plains Region of North Dakota Based on Fish MMI Scores.

Due to relatively low species diversity in the Rangeland Plains region of North Dakota (ecoregions 42 and 43), the fish indicator performed poorly. Therefore, rather than reporting on the biological condition of streams in this region through the use of an MMI, a general description of the overall fisheries quality will be provided. Thirty-three (33) probability and hand-picked sites were sampled with a total of 8,388 fish collected via long-line electrofishing.

A total of 2,627 stream km were assessed for general fisheries quality in the Rangeland Plains region (ecoregions 42 and 43) of North Dakota. In this region of the state, 2,783 perennial stream km are estimated to be present. Primary sportfish represented only 3 percent of the total fish collected (271 individuals). Primary sportfish species included smallmouth bass, northern pike, channel catfish, sauger, walleye, white crappie, bluegill, yellow perch and white bass (Figure V-7).

Stream km estimates of fish occurrence are based on presence/absence of a particular species or family. For instance, if a single smallmouth bass (*Centrarchidae*) is collected at four different

locations and each location is representative of 125 stream km in the Rangeland Plains region, then the estimate for smallmouth bass will be 500 stream km. The families Cyprinidae (minnows) and Catostomidae (suckers) were the most frequently collected fish taxa. Both were equally represented in the Rangeland Plains of North Dakota and were estimated to be present in 2,023 stream km. Ictalurids (catfish and bullheads) were estimated to be present in 1,846 stream km while Centrarchids (sunfish and black bass) were estimated to inhabit 1,523 miles of stream km. Finally, Esocidae (pike) and Percidae (perch, walleye and sauger) were estimated to be present in 1,302 and 1,221 stream km, respectively (Figure V-6).




Figure V-6. Perennial Stream Length (km) Estimates of Each Family of Fish in the Rangeland Plains Region of North Dakota.

Interestingly, although the stream km estimates based only on occurrence are similar in the Rangeland Plains of North Dakota for each taxonomic group, the relative abundance of fish is dominated by the family Cyprinidae at 82.9 percent. Catostomidae was the next most abundant family and only accounted for 8.8 percent of collected individuals. Centrarchidae and Ictaluridae accounted for 3.2 percent and 2 percent, respectively, while Percidae and Esocidae families were present in 1.2 percent and 0.7 percent of the entire sample, respectively (Figure V-8).




Figure V-7. Abundance of Primary Sportfish in the Rangeland Plains Region of North Dakota.




Figure V-8. Percent Abundance of Each Family of Fish in the Rangeland Plains Region of North Dakota.

Periphyton Biotic Integrity

For the periphyton indicator, metric screening was similar to the MMI that was developed for macroinvertebrates. A total of 250 diatom metrics covering five categories were evaluated. First, box-plots of least (reference) and most disturbed (stressed) sites were compared to show which metrics provided separation. Next, through correlation analysis, non-redundant metrics were chosen with an attempt to include as many ecological categories as possible.

In the Cultivated Plains region, 55 samples, including repeat visits, were analyzed. Non-wadable sites were also included in this assessment. The periphyton metrics used in the MMI are: number of species in the old *Cymbella* genus; percent of total taxa that are highly mobile; percent of total taxa in the oxygen class 1 or 2; number of *Gomphonema* species; and percent abundance of individuals in the genus *Fragilaria*.

In the Rangeland Plains region, a total of 70 samples were analyzed for North Dakota. The final metrics chosen for the MMI were: percent of total taxa in the old genus *Cymbella*; percent abundance of total taxa in the old genus *Fragilaria*; percent abundance of individuals in the new genus *Nitzschia*; percent of total taxa that are moderate or highly motile; and percent of total taxa in the oxygen class 1 or 2.

For both the Cultivated Plains and the Rangeland Plains, good and poor condition class thresholds were set at the 25th and 10th percentile of the least disturbed (reference) sites (Table V-6). Similar to the macroinvertebrate indicator, each site visited was scored independently, according to region (Cultivated Plains or Rangeland Plains). Final scores were then calculated for each site and data was then pooled as good, fair or poor for an overall statewide assessment.

In the Cultivated Plains, scores greater than 44.1 were considered good. Scores that were greater than or equal to 21.4 and less than or equal to 44.1 were considered fair, while any site scoring less than 21.4 was considered to be in poor condition. In the Rangeland Plains, any site scoring greater than 42.5 was considered good. Site scores that were greater than or equal to 32.3 and less than or equal to 42.5 were in fair condition, and anything less than 32.3 was considered to be in poor condition.

Based on the periphyton MMI, North Dakota has 2,305 stream km (35 percent) in good condition, 2,585 km (40 percent) in fair condition and 1,665 km (25 percent) in poor condition (Figure V-9).




Figure V-9. Condition Class Summary for Perennial Streams in North Dakota Based on Periphyton MMI Scores.

Stressor Indicators

Environmental stressors are defined as (1) the chemical, physical habitat and biological components of the ecosystem that have the potential to degrade biotic integrity and (2) the pressures that human beings exert on habitat systems through their use of the surrounding environment. For this project, the environment was defined as perennial streams in North Dakota. Chemical stressors in North Dakota include excess nutrients and chemical contamination (pesticides and trace metals). Physical habitat stressors include excess sedimentation, bed and bank erosion, and loss of streamside vegetation. Biological stressors include the presence of invasive species. The key to stressor assessment is determining how common a stressor is in a region and how severely the stressor affects biotic integrity. During this assessment of perennial streams in North Dakota, no biological stressors were evaluated for their effect on biotic integrity.

Chemical Stressors

Three chemical stressors (total phosphorus, total nitrogen, and specific conductance) were assessed for both the Cultivated and Rangeland Plains regions of North Dakota. A fourth chemical stressor (mercury in fish tissue) was assessed for perennial streams in the state as a whole. These stressors were the same as those used in the South Dakota assessment. Thresholds for total phosphorus, total nitrogen and mercury in fish tissue were the same as those used in South Dakota, while the thresholds for specific conductance were the same as those used in the EMAP-West report. Thresholds for the Cultivated Plains are shown in Table V-8; thresholds for

the Rangeland Plains are shown in Table V-9; and thresholds for mercury in fish tissue are shown in Table V-10.

Table V-8. Threshold Values Used to Determine Condition Classes for Three Chemical Stressors in the Cultivated Plains Region of North Dakota.

Chemical Stressor	Poor	Fair	Good
Total phosphorus	>312 ug/L	228-312 ug/L	<228 ug/L
Total nitrogen	>2501 ug/L	1525-2501 ug/L	<1525 ug/L
Specific conductance	>2000 uS/cm	1000-2000 uS/cm	<1000 uS/cm

Table V-9. Threshold Values Used to Determine Condition Classes for Three Chemical Stressors in the Rangeland Plains Region of North Dakota.

Chemical Stressor	Poor	Fair	Good
Total phosphorus	>138 ug/L	70-138 ug/L	<70 ug/L
Total nitrogen	>1186 ug/L	886-1186 ug/L	<886 ug/L
Specific conductance	>2000 uS/cm	1000-2000 uS/cm	<1000 uS/cm

Table V-10. Threshold Values Used to Determine Condition Classes for the Mercury in Fish Tissue Stressor in North Dakota.

Poor	Fair	Good
>0.1 ug/g	Not Assessed	<0.1 ug/g

Chemical stressor assessments for perennial streams in the Cultivated Plains of North Dakota are shown in Figure V-10. Based on the total phosphorus stressor, 1,645 km (42 percent) of streams were found to be in poor condition and 1,616 km (41 percent) were in good condition. Only 668 km (17 percent) of streams were found to be in fair condition. Total nitrogen and specific conductance stressor thresholds were exceeded for poor condition for only 238 km (6 percent) and 373 km (9 percent) of streams, respectively. The remaining 3,717 km (94 percent) of streams were assessed to be in fair to good condition for total nitrogen and about 3,583 km (91 percent) of streams were assessed to be in fair to good condition for specific conductance.

Chemical stressor assessments for perennial streams in the Rangeland Plains region are shown in Figure V-11. Based on total phosphorus stressor thresholds, 1,031 km (39 percent) of perennial streams in the Rangeland Plains region were found to be in poor condition, while 1,249 km (48 percent) of assessed streams were found to be in good condition and 341 km (13 percent) were found to be in fair condition. Total nitrogen and specific conductance stressor thresholds were exceeded for poor condition for 479 km (18 percent) and 1,200 km (46 percent) of perennial streams, respectively. The remaining 2,148 km (82 percent) of perennial streams were assessed in fair to good condition for total nitrogen and about 1,427 km (54 percent) were assessed in fair to good condition for specific conductance. While a relatively small percentage of perennial streams in the Cultivated Plains were assessed as being in poor condition (9 percent) based on the specific conductance stressor, a relatively large percentage of streams in the Rangeland Plains region (46 percent) were assessed as being in poor condition based on the same stressor. This may be due to climatic differences between the Rangeland Plains and the Cultivated Plains. The Rangeland Plains area tends to be a dry, arid environment which may account for these differences.




Figure V-10. Perennial Stream Length (km) in the Cropland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Three Chemical Stressors.




Figure V-11. Perennial Stream Length (km) in the Rangeland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Three Chemical Stressors.

The chemical stressor assessment of perennial streams for North Dakota as a whole reflects the summation of the assessments for the Cultivated Plains and Rangeland Plains regions (Figure V-12). Based on total phosphorus stressor thresholds, 2,677 km (41 percent) of perennial streams in North Dakota were found to be in poor condition, while 2,866 km (43 percent) of assessed streams were found to be in good condition and 1,040 km (16 percent) were found to be in fair condition. Total nitrogen and specific conductance stressor thresholds were exceeded for poor condition for 717 km (11 percent) and 1,573 km (24 percent) of perennial streams, respectively. The remaining 5,866 km (89 percent) of perennial streams were assessed in fair to good condition for total nitrogen, while about 5,010 km (76 percent) were assessed in fair to good condition for specific conductance.




Figure V-12. Perennial Stream Length (km) in North Dakota Estimated to be in Good, Fair and Poor Condition Based on Three Chemical Stressors.

Based on the mercury in fish tissue stressor, approximately 3,457 km (56 percent) of the perennial streams in North Dakota were assessed to be in poor condition (mercury concentrations in fish tissue greater than 0.1 ug/g) (Figure V-13). Only about 1,830 km (30 percent) of the perennial streams were considered to be in good condition based on the mercury in fish tissue stressor. A total of 829 km of streams lack fish available for tissue analysis and, therefore, were not assessed.




Figure V-13. Perennial Stream Length (km) in North Dakota Estimated to be in Good and Poor Condition Based on the Mercury in Fish Tissue Stressor.

Physical Habitat Stressors

Four physical habitat stressors were assessed for North Dakota perennial streams: streambed stability, riparian vegetation, riparian disturbance and habitat complexity. These stressors are the same as those defined and used in the EMAP-West assessment. Thresholds for each physical habitat stressor for the Cultivated Plains region (Table V-11) and for the Rangeland Plains region (Table V-12) were the same as those used in the South Dakota assessment (Stoddard and others, no date).

Table V-11. Threshold Values Used to Determine Condition Classes for Four Physical Habitat Stressors in the Cultivated Plains Region of North Dakota.

Physical Habitat Stressor	Poor	Fair	Good
Streambed Stability	<-2.58	-2.58 to -2.20	>-2.20
Riparian Disturbance	>1.8	1.31-1.8	<1.31
Habitat Complexity	<0.136	0.136-0.214	>0.214
Riparian Vegetation	<0.041	0.041-0.236	>0.236

Table V-12. Threshold Values Used to Determine Condition Classes for Four Physical Habitat Stressors in the Rangeland Plains Region of North Dakota.

Physical Habitat Stressor	Poor	Fair	Good
Streambed Stability	<-3.01	-3.01 to -2.54	>2.54
Riparian Disturbance	>1.57	1.43-1.57	<1.43
Habitat Complexity	<0.152	0.152-0.278	>0.278
Riparian Vegetation	<0.124	0.124-0.276	>0.276

Physical habitat stressor assessments for perennial streams in the Cultivated Plains region of North Dakota are shown in Figure V-14. Approximately 2,148 km (55 percent) of streams were assessed in poor condition, while only 1,148 km (29 percent) of streams were assessed in good condition for streambed stability. Condition assessments based on riparian disturbance and riparian vegetation indicated that perennial streams in the Cultivated Plains region were considered in good condition for 2,180 km (55 percent) and 2,986 km (76 percent), respectively.

Only 77 km (2 percent) of stream length was assessed in poor condition for riparian vegetation. Habitat complexity was more evenly divided with 1,482 km (38 percent) of streams assessed in poor condition, 1,561 km (40 percent) assessed in fair condition and 885 km (22 percent) assessed in good condition.




Figure V-14. Perennial Stream Length (km) in the Cultivated Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Four Physical Habitat Stressors.

Physical habitat stressor assessments for perennial streams in the Rangeland Plains region are shown in Figure V-15. Approximately 1,221 km (46 percent) of streams were assessed in good condition, and only 781 km (30 percent) of streams were assessed in poor condition for streambed stability. Assessments of riparian disturbance and riparian vegetation stressors indicated streams were considered in good condition for 1,427 km (54 percent) and 898 km (34 percent) of stream length, respectively. The habitat complexity stressor assessment indicated that 1,773 km (68 percent) of stream length were considered in poor condition.

The riparian vegetation stressor assessment showed the greatest difference between the Rangeland Plains and Cultivated Plains regions. Approximately 76 percent of the assessed streams in the Cultivated Plains region were considered to be in good condition while only about 34 percent of the streams assessed in the Rangeland Plains region were considered to be in good condition. Because livestock grazing is more dominant in the Rangeland Plains region than in the Cultivated Plains region, it is likely that over-grazing by livestock in the Rangeland Plains region may be degrading riparian vegetation.

The physical habitat stressor assessment for North Dakota as a whole reflects the summation of the assessments for the Cultivated Plains and Rangeland Plains regions (Figure V-16).

Approximately 2,369 km (36 percent) of streams were assessed in good condition, while 2,929 km (45 percent) of streams were assessed in poor condition for streambed stability. Assessments of riparian disturbance and riparian vegetation stressors indicated streams were considered in good condition for 3,607 km (55 percent) and 3,884 km (59 percent) of stream length, respectively. The habitat complexity stressor assessment indicated that 3,255 km (50 percent) of stream length were considered in poor condition.




Figure V-15. Perennial Stream Length (km) in the Rangeland Plains Region of North Dakota Estimated to be in Good, Fair and Poor Condition Based on Four Physical Habitat Stressors.




Figure V-16. Perennial Stream Length (km) in North Dakota Estimated to be in Good, Fair and Poor Condition Based on Four Physical Habitat Stressors.

B. Lakes and Reservoirs Water Quality Assessment

Chapter 1. Assessment Category Summary

Of the 247 public lakes and reservoirs included in the Assessment Database (ADB), only 197 are included in the state's water quality standards as classified lakes and therefore are assigned designated beneficial uses. The remaining 50 lakes and reservoirs, while included in the state's estimate of total lake acres, were not assessed for this report. Table V-13 provides an assessment category summary for the 197 classified lakes and reservoirs in the state. One lake was classified as Category 1, meaning all uses were assessed and were fully supporting. One-hundred-fifty-two (152) lakes and reservoirs totaling 189,619.1 acres were assessed as Category 2. These are lakes and reservoirs where at least one designated use, mostly agriculture use and industrial use, was assessed as fully supporting, but the other uses were not assessed. A total of 11 lakes and reservoirs were assessed as Category 4A, meaning at least one designated use was impaired or threatened, but a TMDL is not required because a TMDL already has been completed and approved by EPA. Thirty-three (33) lakes and reservoirs totaling 506,572.1 acres were assessed where at least one beneficial use is impaired and a TMDL is required. These Category 5 lakes and reservoirs are provided in the state's TMDL list (Tables VI-1 through VI-4).

Table V-13. Assessment Category Summary for Lakes and Reservoirs in North Dakota (Acres)

Category	Description	Number AUs	Total Size (acres)
1	All uses met	1	885.3
2	Some uses met, others not assessed	152	189,521.4
3	No uses assessed	0	0
4A	Some or all uses impaired or threatened, but a TMDL(s) has been approved for all impaired uses.	11	3,323.83
4B	Some or all uses impaired or threatened, but other pollutant controls will result in water quality standards attainment.	0	0
4C	Some or all uses impaired or threatened, but impairment is not due to a pollutant.	0	0
5	Some or all uses impaired or threatened and a TMDL is required.	33	506,487.7

Chapter 2. Water Quality Summary

A total of 197 lakes and reservoirs, representing 700,315.89 surface acres, were assessed for this report. The remaining 50 lakes and reservoirs, included in the ADB, but not assessed, represent 61,455.61 acres or only 5.5 percent of the total lake and reservoir acres in the state.

For purposes of this report, the term “aquatic life use” is synonymous with biological integrity and is defined as the ability of a lake or reservoir to support and maintain a balanced, adaptive community of aquatic organisms (e.g., fish, zooplankton, phytoplankton, macroinvertebrates, vascular plants) having a species composition, diversity and functional organization comparable to that of least-impaired reference lakes and reservoirs in the region (modified from Karr et al., 1981). One-hundred-twenty-four (124) lakes and reservoirs, representing 686,115.1 acres, were assessed as fully supporting aquatic life use (Table V-14); in other words, they are considered capable of supporting and maintaining a balanced community of aquatic organisms. Of this total, 30 lakes and reservoirs representing 376,606.3 acres are considered threatened (Table V-14). A threatened assessment means that if water quality and/or watershed trends continue, it is unlikely these lakes will continue to support aquatic life use. The lakes and reservoirs will begin to experience more frequent algal blooms and fish kills. They will display a shift in trophic status from a mesotrophic or eutrophic condition to a hypereutrophic condition. Only three lakes, totaling 171.8 acres, were assessed as not supporting aquatic life use (Table V-14).

Table V-14. Individual Use Support Summary for Lakes and Reservoirs in North Dakota (Acres).

Use	Fully Supporting	Fully Supporting but Threatened	Not Supporting	Not Assessed	Insufficient Information for Assessment	Total Size
Aquatic Life	309,508.8	376,606.3	171.8	12,679.3	1,349.7	700,315.9
Fish Consumption	0	0	485,928.0	213,502.6	0	699,430.6
Recreation	545,336.9	135,366.4	5,546.8	13,418.8	647.0	700,315.9
Drinking Water Supply	481,406.2	0	0	217,385.4	229.0	699,202.6
Agriculture	700,315.9	0	0	0	0	700,315.9
Industrial	700,315.9	0	0	0	0	700,315.9

One of the primary causes of aquatic life impairment to the state’s lakes and reservoirs is low DO in the water column (Table V-15). Low DO in lakes can occur in summer (summer kills), but usually occurs in the winter under ice-cover conditions. Low-DO and winter kills occur when senescent plants and algae decompose, consuming available oxygen. Because the lake is ice covered, re-aeration is minimal, and the lake goes anoxic resulting in a fish kill. Fish kills are the most apparent impact to sensitive fish species (e.g., walleye, trout, bass, bluegill, crappie, northern pike), but impacts to other DO-sensitive aquatic organisms also may occur. When fish kills occur, low DO-tolerant fish species (e.g., carp, bullhead, white suckers) will be favored, resulting in a lake dominated by these rough fish species.

Pollutants that stimulate the production of organic matter also can cause aquatic life impairment.

Two secondary pollutant causes are excessive nutrient loading and siltation (Table V-15). Major sources of nutrient loading to the state's lakes and reservoirs are erosion and runoff from cropland, runoff from animal feeding operations (e.g., concentrated livestock feeding and wintering operations) and hydrologic modifications (Table V-16). Hydrologic modifications, such as wetland drainage, channelization and ditching, increase the runoff and delivery rates to lakes and reservoirs in effect increasing the size of a lake's watershed. Nutrients, sediment and organic matter that would be retained in wetlands under normal conditions become part of the lake's external budget.

Other sources of nutrient loading that affect lakes in the state are point source discharges from municipal wastewater treatment facilities, urban/stormwater runoff and shoreline development (Table V-16).

Table V-15. Impairment Summary for Lakes and Reservoirs in North Dakota.

Impairment	Acres
Oxygen Depletion	374,506.7
Temperature	368,231.0
Nutrients	140,968.4
Sedimentation/Siltation	5,499.9
Turbidity	1,567.8
Total Dissolved Solids	36.8
Mercury in Fish Tissues	485,928.0

Shoreline or cabin development directly contributes nutrients to lakes in many ways. Typically, lake cabins or homes use septic systems (tanks and drain fields) to contain their wastewater. Many of these systems are poorly designed, poorly maintained or nonexistent. Poorly designed septic systems provide a direct path of nutrients from the cabin to the lake. In addition, cabins or homes along lakes can contribute nutrients through fertilizer runoff from lawns.

Shoreline development can indirectly lead to increased nutrient loading when development results in a loss of the natural vegetation surrounding the lake. This buffer, between the lake and its watershed, provides for the assimilation of nutrients and retention of sediments contained in the runoff from the surrounding landscape. When this buffer is lost or degraded due to development, nutrients, sediment and other chemicals (e.g., pesticides, road salts) are afforded a direct path to the lake.

The previously mentioned sources are considered external or watershed-scale sources of nutrient loading. Another source that can represent a significant portion of the nutrient budget at times is internal cycling, particularly in those lakes that periodically go anoxic either during ice cover or through thermal stratification in the summer. Under these circumstances, phosphorus and reduced forms of nitrogen (e.g., ammonia) can be released into the water column. The increased nutrient concentrations impair use by stimulating noxious weed growth and algal blooms.

Recreation use (e.g., swimming, waterskiing, boating, sailing, sunbathing) was assessed for 686,250.1 lake and reservoir acres in the state. Of this total, two (2) lakes, representing 5,546.8 acres, were assessed as not supporting use for recreation (Table V-14). The primary cause of use impairment is excessive nutrient loading, which results in nuisance algal blooms and noxious aquatic plant growth (Table V-15). Sources of nutrients causing algal blooms and weed growth were described earlier (Table V-16). Thirty-seven (37) lakes and reservoirs, totaling 135,366.4 acres, were assessed as threatened (Table V-14). Nutrient loading also is linked to the negative water quality trends these lakes are experiencing. If left unchecked, these lakes will degrade to the point where frequent algal blooms and/or excessive weed growth will negatively affect recreation.

One-hundred and ninety-six (196) lakes and reservoirs, representing 699,430.6 acres, were assigned the use for fish consumption (Table V-14). Lakes not assigned the fish consumption use are saline lakes that cannot support a sport fishery. These lakes are also not assigned the use for municipal drinking water supply.

Of the 196 lakes and reservoirs entered into the ADB and assigned a use for fish consumption, only Devils Lake and Lake Sakakawea had sufficient methyl-mercury fish tissue data and fish population survey data necessary to calculate weighted average concentrations and to assess fish consumption use. Based on these data (see Appendices B-E), both were assessed as not supporting fish consumption use (Table V-14). The remaining 194 lakes and reservoirs that support a sport fishery were not assessed for this report.

Sources of methyl-mercury in fish remain largely unknown. Potential sources of mercury include natural sources and atmospheric deposition. Results of a report prepared by the department show an increase in mercury concentrations in the fillets of walleye, northern pike and chinook salmon in Lake Sakakawea following the drought and recent filling of the lake (Pearson et al., 1997). One possible reason for the higher mercury concentrations in fish is that the lake may be experiencing an increase in the rate of mercury methylation due to greater amounts of organic matter in the lake following flooding. The drought of the late 1980s and early 1990s lowered the lake level, allowing vast areas of dry lake bed to re-vegetate. When the lake began refilling in 1993, the vegetation was flooded and began decomposing. The organic matter provided to the lake during this period is thought to have favored the methylation process. This is a microbial process whereby bacteria present in the lake convert elemental mercury to its more bioavailable methyl-mercury form. The increase in bioavailable mercury in the lake is reflected in higher mercury concentrations in fish.

Five reservoirs (Lake Sakakawea, Lake Ashtabula, Homme Dam, Bisbee Dam and Mt. Carmel Reservoir) are currently used either directly or indirectly as municipal drinking water supplies, while two others (Patterson Lake and Renwick Dam) serve as back-up water supplies in the event the primary water supplies should fail.

Homme Dam, Mt. Carmel Reservoir and Lake Sakakawea were assessed as fully supporting drinking water supply use (Table V-14). Drinking water supply use was not assessed for the remaining lakes and reservoirs.

Table V-16. Impairment Source Summary for Lakes and Reservoirs in North Dakota.

Source	Acres
Source Unknown (Associated with Mercury in Fish)	485,928.0
Hydrostructure Flow Regulation/Modification	368,231.0
Crop Production (Dryland)	141,282.2
Internal Nutrient Recycling	139,777.3
Rangeland/Pastureland Grazing	126,299.7
Wetland Loss (Drainage/Filling)	125,895.3
Stormwater Runoff	117,760.3
Riparian Grazing	14,647.5
Animal Feeding and Handling Operations	13,484.2
On-site Treatment Systems (Septic Systems)	9,607.2
Anoxia Due to Thermal Stratification/Eutrophication	6,275.7
Sediment Resuspension	2,518.4
Upstream Impoundments	2,073.4
Highway and Road Runoff	413.6
Surface Mining	376.8
Streambank Modification	392.5
Loss of Riparian Habitat	194.0
Land Application of Biosolids/Septage Disposal	55.2
Flow Alteration for Water Diversion	36.8

Chapter 3. Trophic Status

Reservoirs and natural lakes were assessed for trophic status only if appropriate data were available. For purposes of this report, “trophic status” refers to the present condition or measure of eutrophication of the waterbody at the time of the assessment.

Accurate trophic status assessments are essential to making sound management decisions. In order to minimize errors in classification, all existing chemical, physical, quantitative and qualitative data were used in making final trophic status assessments.

Because there are no TSIs specific to North Dakota waters, Carlson's TSI (Carlson, R. E. 1977, “A Trophic State Index for Lakes,” *Limnology and Oceanography*, 22(2):361-369) was chosen as the initial method to describe a lake's or reservoir's trophic status. Carlson's TSI was selected because it is commonly used by limnologists and because it was developed for Minnesota, a state geographically close to North Dakota.

An attempt was made to gather enough chemical and ancillary data to group as many of North Dakota's 197 classified lakes/reservoirs into one of four trophic states (Table V-17). The four trophic states, in order of increasing productivity, are oligotrophic, mesotrophic, eutrophic and hypereutrophic. Adequate data was available to assess the trophic status of 128 of the 197 lakes entered into the ADB database. The majority of the state's assessed lakes and reservoirs range from eutrophic to hypereutrophic. Forty-one (41) lakes and reservoirs were assessed as mesotrophic. There were no oligotrophic lakes assessed in the state.

Table V-17. Trophic Status Summary for Lakes and Reservoirs in North Dakota

Trophic Status	Number of Lakes	Acreage of Lakes
Oligotrophic	0	0.0
Mesotrophic	41	509,461.0
Eutrophic	61	47,283.7
Hypereutrophic	26	128,925.9
Not Assessed	69	14,645.3
Total Number of Lakes	197	700,315.9

Chapter 4. Control Methods

NPS pollution, particularly from agricultural lands and feedlots, is the main source of pollutants leading to the degradation of the state's lakes and reservoirs. North Dakota's Section 319 NPS Pollution Management Program is very active in reducing agricultural NPS pollution (see Part III. C. Chapter 3. "NPS Pollution Management Program"). This program has kept thousands of tons of soil, along with attached contaminants, out of the state's lakes and reservoirs.

Currently, the Section 319 NPS Pollution Management Program is providing cost-sharing for four (4) watershed restoration projects that have a direct impact on lakes or reservoirs in the state. These include Lake Hoskins, Pheasant Lake, Powers Lake and McDowell Dam. These projects treat entire watersheds through the promotion of sustainable agricultural and sound land management practices. Landowner participation is voluntary, with incentives provided by cost-share programs.

Point source pollution has the potential to severely impact individual lakes and reservoirs and is the second largest pollution problem. Protection of lakes and reservoirs from point source discharges is accomplished through the NDPDES Program (see Part III. C. Chapter 2. "Point Source Control Program"). While the NDPDES Program is thought of as regulating only industrial and municipal discharges, permits also are required for stormwater discharges and large animal feeding operations.

Chapter 5. Restoration/Rehabilitation Efforts

The primary intent of the Section 319 NPS Pollution Management Program is to control NPS pollution to lakes and reservoirs on a watershed scale. This program is complemented by the North Dakota Game and Fish Department's "Save Our Lakes" program. The main goal of the "Save Our Lakes" program is "to enhance and restore North Dakota's aquatic habitat resources in order to protect the fishery of North Dakota." In general, this encompasses shoreline enhancement projects, sediment dam installation, sediment removal, grass and tree plantings, cross fencing, alternate water sources, the installation of passive low water draw-downs, cost-share assistance for animal waste management systems and the establishment of exclusion areas in riparian corridors.

Chapter 6. Acid Effects on Lakes and Reservoirs

Acid precipitation and acid mine drainage pose significant threats to some of the nation's lakes and streams. Most surface waters in North Dakota are naturally alkaline ($\text{pH} > 7$), while rainfall is naturally acidic ($\text{pH} < 7$). Surface waters are able to resist acidification by what is termed "buffering capacity." In surface waters, buffering capacity is maintained largely by the carbonate (CO_3^{2-}) and bicarbonate (HCO_3^{-1}) ions in solution. These ions are collectively measured with hydroxide ions (OH^{-1}) as total alkalinity. Acidification in surface waters occurs when the buffering capacity is exhausted, thus causing a reduction in pH. North Dakota's lakes are highly alkaline and, as a result, do not show acidity caused by anthropogenic sources.

Chapter 7. Toxic Effects on Lakes and Reservoirs

Currently, mercury is the only contaminant assessed as causing lake and reservoir use impairment. As stated previously, elevated mercury concentrations in the tissues of fish have resulted in site-specific consumption advisories for Devils Lake, Lake Sakakawea and Lake Oahe and a general fish consumption advisory for all lakes and reservoirs in the state. Again, very little is known about the source of the mercury contamination in fish from these lakes. It is likely, however, that sources are both natural and anthropogenic.

In 1991, the department initiated the LWQA Project, by which the state's lakes and reservoirs were systematically sampled and assessed for trophic status and watershed condition. In addition to data for assessing the general condition of each lake, data were also collected on the type, concentration and location of contaminants like trace elements and organic compounds.

To date, sediments and fish have been collected from 113 lakes and reservoirs throughout the state. This data should provide useful information for determining baseline contaminant concentrations and examining patterns in contaminant concentrations in lakes and reservoirs.

C. Wetlands Assessment Program

Chapter 1. Background

Wetlands have long been regarded as nuisance areas or wastelands which only serve to impede agriculture, urban or transportation development. It is only recently that the ecological and social functions and values of wetlands been realized. It is now scientifically proven that wetlands are important for the storage of flood waters, for providing fish and wildlife habitat, for recharging ground water and for retaining and cycling chemical pollutants and particulates. Recently, wetlands have been recognized as a significant source for carbon sequestration. This could make wetlands an important component in the campaign to prevent global warming.

While these are important wetland functions, probably the best known function of wetlands in North Dakota is that of waterfowl production. Most of North Dakota's remaining wetlands are located in an area known as the Prairie Pothole Region. This area extends from the Missouri Coteau in central North Dakota eastward to the glacial Lake Agassiz Plain, also known as the Red River Valley. The region covers roughly 300,000 square miles and exists as a wide band extending from central Alberta southwest into northwestern Iowa (Figure V-17). The Prairie Pothole Region, with its many types of wetlands, is arguably the most biologically diverse and productive habitat in North America.




Figure V-17. Prairie Pothole Region

Chapter 2. Extent of Wetland Resources

There seem to be as many ways to classify wetlands as there are wetlands themselves. The U. S. Fish and Wildlife Service first began to classify wetlands based on a system developed by Martin et al. (1953). This classification system was then modified by Stewart and Kantrud (1971), specifically for the Prairie Pothole Region of North America. With the Stewart and Kantrud classification system, vegetational zones are described in detail, along with the plant species most commonly found in the zone. These zones are used to identify phases which indicate the wetland's water regime or disturbed bottom soil (e.g., cropland tillage). Seven wetland classes are identified with the Stewart and Kantrud system. These include the familiar Class I - ephemeral ponds, Class II - temporary ponds, Class III - seasonal ponds and lakes, Class IV - semi-permanent ponds and lakes, and Class V - permanent ponds and lakes. Also included in the Stewart and Kantrud system are Class VI - alkali ponds and lakes, and Class VII - fens. Along with each class, there are five subclasses, A through E, based on variations in surface water salinity. Those familiar with the Stewart and Kantrud classification system refer to temporary depressional wetlands as Class II wetlands, seasonal wetlands as Class III wetlands and semi-permanent wetlands as Class IV.

In 1979, the U.S. Fish and Wildlife Service adopted the Cowardin et al. (1979) classification system for wetlands and deep water habitats of the United States. The Cowardin et al. classification system was developed to be used with the National Wetlands Inventory. In the highest level of classification, wetlands are grouped into five ecological systems: palustrine, lacustrine, riverine, estuarine and marine. The palustrine class includes only wetlands, whereas each of the four other systems includes wetlands and associated deep-water habitats. For purposes of classification, deep-water habitats are defined as areas where water is greater than 6.6 feet deep. In North Dakota, only the palustrine, lacustrine and riverine wetland types exist.

Brinson (1993) developed a classification system for use by the U.S. Army Corps of Engineers. This classification system, termed the Hydrogeomorphic (HGM) classification system, is based upon the wetland's position in the landscape (i.e., geomorphic setting), dominant source of water and the flow and fluctuation of water in the wetland. Brinson (1993) describes seven HGM wetland classes: riverine, depressional, slope, mineral soil flats, organic soil flats, estuarine fringe and lacustrine fringe.

In North Dakota, wetlands are classified into four broad categories according to the State Engineer's drainage rules. The state wetland classification includes temporary wetlands, seasonal wetlands, semi-permanent wetlands and permanent wetlands. The following are brief descriptions of each wetland class, as adopted by the North Dakota State Game and Fish Director and the State Engineer.

"Temporary wetlands" are shallow depressions which hold water or are waterlogged from spring runoff until early June. In years with normal runoff and precipitation, these areas may be tilled for crop production. In years with high runoff or heavy spring rain, these areas may not dry out until mid-July. They cannot be tilled, but may be used for hayland or pasture. Temporary wetlands frequently reflood during heavy summer and fall rains. Sheet water, as defined in North Dakota's Century Code 61-32-02, does not fall under the temporary wetland classification.

“Seasonal wetlands” are depressions, which normally hold water from spring runoff until mid-July. In years with normal runoff and precipitation, these wetlands cannot be tilled but may be used for hayland and pasture. In low runoff or dry years, these areas may be tilled for crop production but commonly reflood with heavy summer and fall rains.

“Semi-permanent wetlands” are located in well-defined depressions or basins. In normal years, these areas hold water throughout the summer. Semi-permanent wetlands generally become dry only in years of below normal runoff and precipitation. Freshwater semi-permanent wetlands (commonly called cattail sloughs) are characterized by a predominance of cattail and bulrush vegetation in scattered areas of open water. Saline semi-permanent wetlands have a preponderance of alkali bulrush in scattered areas of open water.

“Permanent wetlands” are located in well-defined basins which characteristically hold water throughout the year. The wetlands become dry only after successive years of below normal runoff and precipitation. Freshwater permanent wetlands typically have a border of aquatic vegetation and predominant open-water areas in the interior. Saline permanent wetlands are typically devoid of emergent vegetation and exhibit a white, salt-encrusted shoreline.

Currently, there are no accurate estimates of state wetland acreage based on wetland class. Statewide, it is estimated there are approximately 2.5 million acres of wetlands. When compared to the approximately 4.9 million acres of wetlands which covered North Dakota prior to development, this represents a 49 percent reduction in wetlands. Stewart and Kantrud (1973) divided the state into four biotic regions: the Prairie Pothole Region, the Lake Agassiz Plain Region, the Coteau Slope Region and the Southwestern Slope Region. They estimated that 81 percent of the wetlands in the state are located in the Prairie Pothole Region. More than 90 percent of all wetlands in the state are considered natural basin wetlands, commonly referred to as prairie potholes. Furthermore, it is estimated that 78 to 79 percent of wetland basins in the Prairie Pothole Region are less than one acre in size (Ron Reynolds, personal communication). While the rate of wetland loss in the state seems to be decreasing, it is safe to assume that wetland losses still exceed wetland gains.

Chapter 3. Integrity of Wetland Resources

Wetland integrity should be thought of in terms of whether a wetland performs a set of functions or uses which would be expected for natural or “reference” wetlands of a similar class or type. The USDA NRCS and the U.S. Army Corps of Engineers have described 11 specific functions within three general functional categories for temporary and seasonal Prairie Pothole wetlands (Lee et al., 1997) (Table V-18). Therefore, whenever a wetland’s function is diminished, it can be said that wetland integrity is diminished.

Hydrologic manipulation (e.g., drainage, wetland consolidation, channelization, filling) continues to be the greatest impact on the integrity of the state’s wetlands. While not as dramatic, other factors such as chemical contamination, nutrient loading (i.e., eutrophication) and sedimentation can also affect a wetland’s function and, therefore, its chemical, physical and biological integrity.

Landscape level changes outside the edge of the wetland basin can also negatively affect wetland integrity. Changes to the landscape, such as road construction, cropland conversion, urbanization or the drainage of adjacent wetlands, all affect wetland functions. Cowardin et al. (1981) found 40 percent of wetlands were cultivated to the wetland edge, 33 percent were in pasture and 7 percent were hayed within a 3,877-square-mile area of the Prairie Pothole Region.

When viewed on a larger scale, wetlands are part of a larger unit known as a wetland complex. Wetland complexes are aggregates of individual wetland basins which are hydrologically connected. A typical wetland complex includes recharge wetlands, flow-through wetlands and discharge wetlands. Recharge wetlands are typically located at higher elevations in the landscape and receive the majority of their hydrologic budgets from precipitation and surface runoff. Recharge wetlands get their name because they recharge ground water. Flow-through wetlands, as their name implies, receive surface- and ground-water inflow and then outflow to both surface and ground water. Discharge wetlands receive the majority of their hydrologic budgets from ground-water discharge and rarely outflow to surface water. Because recharge wetlands receive most of their water through precipitation and surface-water inflow, they tend to be fresher. Discharge wetlands, which receive most of their water from ground water, tend to be higher in total dissolved solids.

Due to this hydraulic linkage in the landscape, any land use change which affects or changes the hydrologic relationship of wetlands in the complex can and will affect the hydrologic or physical integrity of each wetland basin in the complex. This, in turn, affects both the chemical and biological integrity of wetlands in the complex.

Table V-18. Definitions of Functions for Temporary and Seasonal Prairie Pothole Wetlands (Lee et al., 1997).

Physical/Hydrologic Functions
<p>Maintenance of Static Surface Water Storage. The capacity of the wetland to maintain a hydrologic regime that supports static storage, soil moisture in the unsaturated zone and ground water interactions.</p> <p>Maintenance of Dynamic Surface Water Storage. The capacity of the wetland to maintain a hydrologic regime that supports dynamic storage, soil moisture in the unsaturated zone and ground water interactions.</p> <p>Retention of Particulates. Deposition and retention of inorganic and organic particulates ($>0.45\ \mu\text{m}$) from the water column, primarily through physical processes.</p>
Biogeochemical Functions
<p>Elemental Cycling. Short- and long-term cycling of elements and compounds on-site through the abiotic and biotic processes that convert elements (e.g., nutrients and metals) from one form to another; primarily recycling processes.</p> <p>Removal of Imported Elements and Compounds. Nutrients, contaminants, and other elements and compounds imported to the wetland that are removed from cycling processes.</p>
Biotic and Habitat Functions
<p>Maintenance of Characteristic Plant Community. Characteristic plant communities not dominated by non-native or nuisance species. Vegetation is maintained by mechanisms, such as seed dispersal, seed banks and vegetative propagation which respond to variations in hydrology and disturbances, such as fire and herbivores. The emphasis is on the temporal dynamics and structure of the plant community as revealed by species composition and abundance.</p> <p>Maintenance of Habitat Structure Within Wetland. Soil, vegetation and other aspects of ecosystem structure within a wetland required by animals for feeding, cover and reproduction.</p> <p>Maintenance of Food Webs Within Wetland. The production of organic matter of sufficient quantity and quality to support energy requirements of characteristic food webs within a wetland.</p> <p>Maintenance of Habitat Interspersion and Connectivity Among Wetland. The spatial distribution of an individual wetland in reference to adjacent wetlands within the complex.</p> <p>Maintenance of Taxa Richness of Invertebrates. The capacity of a wetland to maintain characteristic taxa richness of aquatic and terrestrial invertebrates.</p> <p>Maintenance of Distribution and Abundance of Vertebrates. The capacity of a wetland to maintain characteristic density and spatial distribution of vertebrates (aquatic, semi-aquatic and terrestrial) that utilize wetlands for food, cover and reproduction.</p>

Chapter 4. Wetland Water Quality Standards

As the lead water quality agency in the state, the department is responsible for developing and implementing water quality standards. In general, the *State Water Quality Standards* (NDDoH, 2006) are regulations which specify the beneficial uses of lakes, reservoirs, rivers and streams in North Dakota. The standards include narrative descriptions, numeric criteria and an antidegradation policy to protect beneficial uses. Common beneficial uses for the state's lakes and rivers are recreation (e.g., swimming, wading, boating, skiing), fishing, drinking water supply and aquatic life. Agriculture (i.e., stock watering and irrigation) and industrial uses for water are also recognized.

The *State Water Quality Standards* already include wetlands in the state's definition of waters of the state. However, beneficial uses have not yet been assigned to wetlands, nor have numeric limits been assigned to protect those uses. Wetlands have been provided some water quality protection by applying North Dakota's narrative standards to wetlands. These narrative standards, also known as the "free from" standards, prohibit the disposal of garbage, oil or any toxic pollutant to wetlands.

Chapter 5. Wetland Monitoring and Assessment Program

Current and Historic Program

Wetlands are often ignored in state water quality monitoring and assessment programs. However, with more than 2.5 million acres of wetlands in the state, the department believes wetland monitoring and assessment should be an important component of its overall water quality monitoring and assessment strategy. To meet its monitoring and assessment goals and objectives for wetlands, the department began developing a Wetland Monitoring and Assessment Program in the early 1990s.

Key to the Wetland Monitoring and Assessment Program has been the development of an Index of Biotic Integrity (IBI) for macroinvertebrates and plants to be used as a tool for assessing the ecological condition of wetlands. While the development of widely applicable and robust indicators for macroinvertebrates has met with limited success, the development of an IBI for wetland plants has been extremely successful. Working in collaboration with the department and with funding provided by EPA's Wetland Program Development Grants, researchers in the North Dakota State University (NDSU) Animal and Range Sciences Department have developed IBIs for plants for temporary, seasonal and semi-permanent depressional wetlands. These IBIs can be applied throughout the Northern Glaciated Plains and Northwestern Glaciated Plains ecoregions.

While an IBI approach to wetland assessment can provide very precise information on the biological condition of individual wetlands or populations of wetlands within regions (e.g., watersheds or ecoregions), it does require the use of personnel skilled in wetland plant identification and can be costly to implement, especially on large regional scales. In order to find a wetland assessment method that is less costly to implement, the department has collaborated with NDSU's Animal and Range Sciences Department to develop a regional-scale wetland assessment methodology using satellite remotely sensed data and GIS tools. This approach has been developed by assembling calibration and verification plant IBI data from wetlands sampled previously and by using multi-spectral Landsat TM and ETM+ satellite data.

Regional Scale Wetland Assessment Pilot Project

With the development of plant IBIs completed for temporary, seasonal and semi-permanent depressional wetlands in the Northern Glaciated Plains and Northwestern Glaciated Plains ecoregions, the department began implementing a regional-scale wetland assessment pilot project. The purpose of this project is to: (1) assess the biological condition of wetlands on a large geographic scale using a probabilistic study design to select and sample wetlands; and (2) apply the plant IBI to assess wetland condition. Results of this regional assessment will then be compared to wetland assessment results that will be conducted using the remote sensing methodology.

Other Program Plans

In advance of the EPA-sponsored National Wetlands Survey, the department has plans to continue to work with NDSU in the selection and development of IBIs for additional wetland

classes with unique ecological regions in the state. The department would also like to refine existing, more labor-intensive wetland assessment methods into a “rapid assessment method” (RAM) for use by volunteer monitoring groups and the regulated community.

D. Public Health/Aquatic Life Concerns

Examples of public health or aquatic life concerns include fishing advisories or bans, pollution-caused fish kills or abnormalities, known sediment contamination, discontinued use of drinking water supplies, closure of swimming areas or incidents of waterborne disease. Unlike many other states, North Dakota has had no reported incidents of drinking water supply restrictions or swimming beach closures for the reporting period 2006 to 2007.

Fish kills occur periodically in the lakes and rivers of the state. When they do occur, it is generally the result of low-water conditions, heavy snow cover or both. Because most fish kills occur during the winter, documenting their occurrence and extent is difficult. In most instances, the occurrence of fish kills is inferred through spring test netting by the North Dakota Game and Fish Department.

The primary public health concern in the state associated with lakes and streams in North Dakota is mercury contamination. In March 1991, the state issued its first fish consumption advisory for lakes and rivers. As new data are collected and analyzed, the department updates the consumption advisory. As stated previously, the consumption advisory for all rivers and lakes in the state is due to elevated concentrations of methyl-mercury in fish tissues. To date, no specific source of mercury contamination has been identified.

PART VI. NORTH DAKOTA SECTION 303(d) LIST OF WATER QUALITY-LIMITED WATERS NEEDING TMDLs

A. Background

Section 303(d) of the CWA and its accompanying regulations (CFR Part 130, Section 7) require each state to list waterbodies (i.e., lakes, reservoirs, rivers, streams and wetlands) that are considered water quality limited and require load allocations, waste load allocations and total maximum daily loads (TMDLs). This list has become known as the “TMDL list” or “Section 303(d) list.”

A waterbody is considered water quality limited when it is known that its water quality does not or is not expected to meet applicable standards. Waterbodies can be water quality limited due to point source pollution, NPS pollution or both.

In considering whether or not applicable water quality standards are being met, the state should consider not only the narrative and numeric criteria set forth in the standards but also the classified uses defined for the waterbody and whether the uses are fully supported or not supported due to any pollutant source or cause. Therefore, a waterbody could be considered water quality limited when it can be demonstrated that a beneficial use (e.g., aquatic life or recreation) is impaired, even when there are no demonstrated exceedances of either the narrative or numeric criteria. In cases where there is a use impairment but no exceedance of the numeric standard, the state should provide information as to the cause of the impairment. Where the specific pollutant (e.g., copper or phosphorus) is unknown, a general cause category (e.g., metals or nutrients) should be included with the waterbody listing.

Section 303(d) and accompanying EPA regulations and policy require only impaired and threatened waterbodies to be listed, and TMDLs are developed when the source of impairment is a pollutant. Pollution, by federal and state definition, is “any man-made or man-induced alteration of the chemical, physical, biological and radiological integrity of water.” Based on the definition of a pollutant provided in Section 502(6) of the CWA and in 40 CFR 130.2(d), pollutants would include temperature, ammonia, chlorine, organic compounds, pesticides, trace elements, nutrients, biochemical oxygen demand (BOD), sediment and pathogens. Waterbodies impaired by habitat and flow alteration and the introduction of exotic species would not be included in the Section 303(d) TMDL list, as these impairment categories would be considered pollution and not pollutants. In other words, all pollutants are pollution, but not all pollution is a pollutant.

Where a waterbody is water quality limited, the state is required to determine in a reasonable time frame the reduction in pollutant loading necessary for that waterbody to meet water quality standards, including its beneficial uses. The process by which the pollutant-loading capacity of a waterbody is determined and the load is allocated to point and nonpoint sources is called a total maximum daily load (TMDL). While the term “total maximum daily load” implies that loading capacity is determined on a daily time scale, TMDLs can range from meeting an instantaneous concentration (i.e., an acute standard) to computing an acceptable annual phosphorus load for a lake or reservoir.

Section 303(d) requires states to submit their lists of water quality-limited waterbodies “from time to time.” Federal regulations have clarified this language; therefore, beginning in 1992 and by April 1 of every even-numbered year thereafter, states are required to submit a revised list of waters needing TMDLs. North Dakota’s 2006 TMDL list was submitted to EPA in April 2006 and was approved in June 2006. This 2008 Section 303(d) list includes waterbodies not meeting water quality standards, waterbodies needing TMDLs and waterbodies that have been removed from the 2006 list. Reasons for removing a waterbody from the 2006 list include: (1) a TMDL was completed for the waterbody/pollutant combination; (2) the applicable water quality standard is now attained and/or the original basis for the listing was incorrect; (3) the applicable water quality standard is now attained due to a change in the water quality standard and/or assessment methodology; (4) the applicable water quality standard is now attained due to restoration activities; or (5) sufficient data and/or information is lacking to determine water quality status and/or the original basis for listing was incorrect.

Along with the TMDL list, states are required to provide documentation to the EPA Regional Administrator in support of the state’s decision to list or not list waterbodies. Information supporting North Dakota’s 2008 TMDL list is provided in Part IV. B. “Assessment Methodology.” At a minimum, a state’s supporting information should include: (1) a description of the methodology used to develop the list; (2) a description of the data and information used to develop the list; (3) the rationale for any decision to not use this information; (4) the rationale for removing waterbodies previously listed as water quality limited; and (5) a summary of comments received on the list during the state’s public comment period.

Following opportunity for public comment, the state must submit its list to the EPA Regional Administrator. The EPA Regional Administrator then has 30 days to either approve or reject the listings. If the EPA Regional Administrator rejects a state submittal, EPA has 30 days to develop a list for the state. This list also is required to undergo public comment prior to finalization.

B. Prioritization of TMDL-Listed Waters

When a state prepares its list of water quality-limited waterbodies, it is required to prioritize waterbodies for TMDL development and to identify those “High” priority waterbodies that will be targeted for TMDL development within the next two to four years. Factors to be considered when prioritizing waterbodies for TMDL development include: (1) the severity of pollution and the uses which are impaired; (2) the degree of public interest or support for the TMDL, including the likelihood of implementation of the TMDL; (3) recreational, aesthetic and economic importance of the waterbody; (4) the vulnerability or fragility of a particular waterbody as an aquatic habitat, including the presence of threatened or endangered species; (5) immediate programmatic needs, such as waste load allocations needed for permit decisions or load allocations for Section 319 NPS project implementation plans; and (6) national policies and priorities identified by EPA.

After considering each of the six factors, the state has developed a two-tiered priority ranking. Assessment units (AUs) listed as “High” priority are: (1) lakes and reservoirs and river and stream segments for which TMDLs are scheduled to be completed and submitted to EPA in the next two years; or (2) lakes and reservoirs and river and stream segments for which TMDL development projects are scheduled to be started in the next two years. The majority of these

“High” priority AUs were identified as such based largely on their degree of public support and interest and the likelihood of implementation of the TMDL once completed. “Low” priority AUs are those river and stream segments and lakes and reservoirs that are scheduled for completion in the next eight years.

The department has also identified a subcategory to Category 5 waterbodies. This subcategory, termed Subcategory 5A, includes “Low” priority lakes and reservoirs and river and stream segments that were assessed and listed in previous Section 303(d) lists, including the 2006 list, but where the original basis for the assessment decision and associated cause of impairment is questionable. These Subcategory 5A waterbodies include: (1) rivers and streams listed for biological impairments based on only one sample for the entire segment or on samples collected more than 10 years ago; (2) waterbodies listed for sediment/siltation impairments; or (3) lakes and reservoirs where the assessments are based on one sampling event or on data that are greater than 10 years old. These waterbodies will remain on the 2008 Section 303(d) list, but they will be targeted for additional monitoring and assessment during the next two to four years.

Waterbodies for which fish consumption use is impaired due to methyl-mercury are also considered “Low” priority. TMDL development for methyl-mercury-contaminated waterbodies is complicated by several factors, including: (1) the uncertainty regarding the fate and transport of atmospheric sources of mercury and (2) the complexity of the biological and geochemical interactions that affect the conversion of elemental mercury to methyl-mercury and its bioaccumulation rate in fish. Due to these complexities and the interstate and international nature of atmospheric mercury sources, it is the department’s recommendation that EPA take the lead in developing mercury TMDLs.

C. Public Participation Process

Public comments were solicited on the draft 2008 TMDL list through a public notice published in the following daily newspapers: Fargo Forum, Grand Forks Herald, Bismarck Tribune, Minot Daily News, Dickinson Press and Williston Daily Herald (Appendix F). The public notice encouraged interested parties to obtain a copy of the draft TMDL list by contacting the department in writing, by phone or by accessing the list through the department’s website at www.health.state.nd.us.

Comments on the draft TMDL list also were requested through mail or email from individuals and specific agencies and organizations. These included the South Dakota Department of Environment and Natural Resources, Minnesota Pollution Control Agency (Detroit Lakes Regional Office), the Natural Resources Conservation Service, the U.S. Fish and Wildlife Service, the U.S. Forest Service, the North Dakota Game and Fish Department, the North Dakota State Water Commission, the Red River Basin Commission, individuals on the North Dakota State Water Pollution Advisory Board and EPA Region VIII. Comments on the draft 2008 TMDL list were only received from EPA Region VIII. These comments and the Department’s response are provided in Appendix H. When appropriate, these comments were incorporated in the final 2008 Integrated Report.

D. Listing of Impaired Waters Needing TMDLs

As stated previously for 2008 Section 305(b) reporting and Section 303(d) TMDL listing, states were encouraged to follow the “Guidance for 2006 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d), 305(b) and 314 of the Clean Water Act” (EPA, 2005). This guidance suggests that states place their assessed waterbodies into one of five assessment categories (Table IV-3). Waterbodies (also referred to as AUs) assessed as Category 5 (including subcategory 5A) form the basis of the state’s Section 303(d) TMDL list. Tables VI-1, VI-2, VI-3 and VI-4 provide a list of AUs in the Souris, Red, Missouri and James River Basins, respectively, that are impaired and in need of TMDLs. These impaired waters also are depicted graphically for the Souris River Basin (Figure VI-1), the Upper and Lower Red River Basins (Figures VI-2 and VI-3), the Lake Sakakawea and Lake Oahe subbasins of the Missouri River Basin (Figures VI-4 and VI-5) and the James River Basin (Figure VI-6). The 2008 TMDL list is represented by 226 AUs (32 lakes and reservoirs and 194 river and stream segments) and 389 individual waterbody/beneficial use/pollutant combinations. For purposes of TMDL development, each waterbody/beneficial use/pollutant combination requires a TMDL. Of the 367 individual waterbody/pollutant combinations listed in Tables V-1 through V-4, 108 waterbody/pollutant combinations were further identified as Category 5A. These waterbodies are targeted for additional monitoring in the next two to four years to verify the current use impairment assessments and pollutant causes.

E. De-listing of 2006-Listed TMDL Waters

Table VI-5 provides a list of lakes, reservoirs, rivers and streams that were listed in the previous 2006 TMDL list but that have been removed from this year’s Section 303(d) list submittal. AUs were removed from the TMDL list for a number of reasons. The following are the primary reasons for de-listing an AU:

- A TMDL was completed for the waterbody/pollutant combination.
- The applicable water quality standard is now attained and/or the original basis for the listing was incorrect.
- The applicable water quality standard is now attained due to a change in the water quality standard and/or assessment methodology.
- The applicable water quality standard is now attained due to restoration activities.
- Sufficient data and/or information is lacking to determine water quality status and/or the original basis for listing was incorrect.

In most cases, when the original assessment was judged not to be representative of current water quality conditions due to a lack of sufficient credible data, one of the following usually occurred:

1. The data used to conduct the assessment are now more than 12 years old for rivers and streams and 14 years old for lakes and reservoirs. Based on best professional judgment, the assessment is no longer believed to be valid. This would occur if it is believed that water quality has been altered due to significant changes in land use and/or due to climatic changes.
2. The original assessment was based only on best professional judgment.

3. The original assessment was based on data extrapolated from a monitoring station(s) located in an adjacent AU.

F. TMDL Development and Monitoring Schedule

The responsibility for TMDL development in North Dakota lies primarily with the department's Division of Water Quality - Surface Water Quality Management Program. TMDL development staff are located in three regional field offices in Dickinson, Fargo and Towner, N.D. Technical support for TMDL development projects and overall program coordination are provided by Surface Water Quality Management Program staff located in Bismarck, N.D.

Historically, the technical and financial resources necessary to complete the state's TMDL development priorities have hampered the pace of TMDL development in the state. Recently, however, the state's TMDL program has seen an improvement in the financial resources available for TMDL development projects. While still significantly short of the funding necessary to meet the state's TMDL development schedule, EPA and the state of North Dakota have made available additional grants and funding to complete TMDLs. Examples of these new financial resources include the TMDL development grants available through EPA Regional VIII and CWA Section 319 grants administered by the state's Nonpoint Source Pollution Management Program.

With the continued commitment to adequate TMDL development staffing and with a continuation in the growth of funding for TMDL development projects in the state, the department is confident it will meet its TMDL development schedule.

The 2008 Section 303(d) TMDL list for North Dakota has targeted 81 waterbodies or 105 waterbody/pollutant combinations for completion in the next two to four years. These "High" priority waterbodies are AUs for which the monitoring is either completed, near completion or has recently been initiated. Of this total, it is expected that TMDLs will be completed for 36 waterbodies (57 waterbody/pollution combinations) in the next two years. These "High" priority waterbodies represent 16 percent of all "High" and "Low" priority Category 5 waterbodies on the list. Based on an anticipated TMDL completion schedule of approximately 40 additional waterbody/pollutant combinations per year following 2010, the department expects to complete TMDLs for all 2008-listed waters in the next eight years.

Table VI-1. 2008 List of Section 303(d) TMDL Waters for the Souris River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09010001-001-L_00	Short Creek Dam	111.5 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	High	No
					Sedimentation/Siltation	High	No
					Nutrient/Eutrophication Biological Indicators	High	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-09010001-001-S_00	Souris River from the N.D./Saskatchewan border downstream to Lake Darling.	43.4 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	No
					Oxygen, Dissolved	High	No
			Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09010001-006-S_00	Souris River from Lake Darling downstream to its confluence with the Des Lacs River. Located in Northern Ward County.	20.3 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09010002-001-S_00	Des Lacs River from lower Des Lacs Reservoir downstream to its confluence with the Souris River. Located in Ward and SW Renville Counties.	71.5 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
					Fecal Coliform	Low	No
			Recreation	Fully Supporting But Threatened			
ND-09010003-001-S_00	Souris River from its confluence with Oak Creek downstream to its confluence with the Wintering River. Located in McHenry County.	51 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	High	No
					Fecal Coliform	High	No
			Recreation	Not Supporting			
ND-09010003-003-S_00	Wintering River, including all tributaries. Located in SW McHenry and NE McLean Counties.	207.8 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened			
			Recreation	Not Supporting			

Table VI-1 (cont.). 2008 List of Section 303(d) TMDL Waters for the Souris River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09010003-005-S_00	Souris River from its confluence with the Wintering River downstream to its confluence with Willow Creek. Located in NE McHenry County.	74.9 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09010004-001-S_00	Willow Creek from its confluence with Ox Creek downstream to its confluence with the Souris River. This ID originally was assigned to the entire Willow Creek reach. The upper reach is assigned the ID of ND-09010004-003-S_01	46.75 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09010004-002-S_00	Oak Creek from its confluence with Willow Creek, upstream to Lake Metigoshe, including all tributaries. Located in Eastern Bottineau County.	82.4 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No




Figure VI-1. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Souris River Basin.

Table VI-2. 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020101-001-S_00	Bois De Sioux River from the ND-SD border, downstream to its confluence with the Rabbit River on MN side. Located in the SE corner of Richland County.	13.05 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation Combination Benthic/Fishes Bioassessments	Low Low	Yes Yes
ND-09020101-002-S_00	Bois De Sioux River from its confluence with the Rabbit River (MN), downstream to its confluence with the Ottertail River. Located on the Eastern border of Richland County.	15.31 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation Benthic-Macroinvertebrate Bioassessments	Low Low	Yes Yes
ND-09020104-001-S_00	Red River of the North from its confluence with the Ottertail River downstream to its confluence with the Whiskey Creek. Located in Eastern Richland County.	27.3 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
			Fish and Other Aquatic Biota	Fully Supporting But Threatened	Combination Benthic/Fishes Bioassessments	Low	Yes
			Fish Consumption	Not Supporting	Methylmercury	Low	No
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020104-002-S_00	Red River of the North from its confluence with Whiskey Creek, downstream to its confluence with the Wild Rice River. Located in NE Richland and SE Cass Counties.	52.3 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020104-003-S_00	Red River of the North, from its confluence with the Wild Rice River, downstream to the 12th Ave bridge in Fargo, ND (just upstream from Moorhead, MN waste water discharge). Eastern Cass County.	21 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
			Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020104-004-S_00	Red River of the North, from the 12th Ave N. bridge in Fargo, ND downstream to its confluence with the Sheyenne River. Eastern Cass County.	21.1 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	High	No
					BOD, carbonaceous	High	No
					Ammonia (Un-ionized)	High	No
			Fish Consumption	Not Supporting			
			Recreation	Fully Supporting But Threatened	Methylmercury	Low	No
					Fecal Coliform	High	No
			Fish Consumption	Not Supporting			
					Methylmercury	Low	No
ND-09020104-005-S_00	Red River of the North from its confluence with the Sheyenne River, downstream to its confluence with the Buffalo River. Located in NE Cass County.	10.45 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020105-001-L_00	Lake Elsie	376.8 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	No
					Turbidity	Low	No
ND-09020105-001-S_00	Wild Rice River from its confluence with the Colfax Watershed, downstream to its confluence with the Red River Of The North. Located in NE Richland and SE Cass Counties.	38.6 miles	Fish and Other Aquatic Biota	Not Supporting			
					Combination Benthic/Fishes Bioassessments	Low	Yes
					Sedimentation/Siltation	Low	Yes
			Recreation	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
					Fecal Coliform	High	No
ND-09020105-002-L_00	Mooreton Pond	36.8 acres	Fish and Other Aquatic Biota	Not Supporting	Total Dissolved Solids	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020105-003-S_00	Wild Rice River from its confluence with a tributary about 3.6 miles NE of Great Bend, ND downstream to its confluence with the Colfax Watershed. Located in Eastern Richland County.	47.5 miles	Fish and Other Aquatic Biota	Not Supporting	Combination Benthic/Fishes Bioassessments	Low	Yes
					Oxygen, Dissolved	Low	No
					Sedimentation/Siltation	Low	Yes
			Recreation	Fully Supporting But Threatened			
					Fecal Coliform	High	No
ND-09020105-005-S_00	Antelope Creek, in Richland County, from its headwaters downstream to its confluence with the Wild Rice River.	40.73 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
					Sedimentation/Siltation	Low	Yes
					Temperature	Low	Yes
			Recreation	Fully Supporting But Threatened			
					Fecal Coliform	High	No
ND-09020105-009-S_00	Wild Rice River from Elk Creek (ND-09020105-010-S_00), downstream to its confluence with a tributary 3.5 miles NE of Great Bend, ND (ND-09020105-008-S_00). Located in South Central Richland County.	53.4 miles	Fish and Other Aquatic Biota	Not Supporting	Oxygen, Dissolved	Low	No
					Sedimentation/Siltation	Low	Yes
			Recreation	Not Supporting			
					Fecal Coliform	High	No
ND-09020105-012-S_00	Wild Rice River from its confluence with Shortfoot Creek (ND-09020105-016-S_00) downstream to its confluence with Elk Creek (ND-09020105-010-S_00).	45.68 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation	Low	Yes
			Recreation	Not Supporting			
					Fecal Coliform	High	No
ND-09020105-016-S_00	Shortfoot Creek from its confluence with the Wild Rice River upstream to the ND-SD border, including all tributaries.	16.16 miles	Recreation	Not Supporting			
					Fecal Coliform	High	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020105-017-S_00	Unnamed tributaries to the Wild Rice River (ND-09020105-015-S), including Crooked Creek.	16.17 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020105-018-S_00	Wild Rice River from its confluence with the Silver Lake Diversion, downstream to Lake Tewaukon.	18.82 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020105-019-S_00	Wild Rice River upstream from its confluence with Wild Rice Creek,	57.06 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020105-020-S_00	Wild Rice Creek from its confluence with the Wild Rice River upstream to the ND-SD border, including all tributaries.	118.17 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020105-022-S_00	Wild Rice River from its confluence with Wild Rice Creek downstream to its confluence with the Silver Lake Diversion.	5.54 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020107-001-S_00	Red River of the North from its confluence with the Buffalo River downstream to its confluence with the Elm River.	29.4 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020107-006-S_00	Elm River from the dam NE of Galesburg, ND downstream to its confluence with the South Branch Elm River.	29.9 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Combination Benthic/Fishes Bioassessments	Low	Yes
					Sedimentation/Siltation	Low	Yes
ND-09020107-008-S_00	Elm River from the dam NW of Galesburg, ND downstream to the dam NE of Galesburg.	20.49 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation	Low	Yes
					Combination Benthic/Fishes Bioassessments	Low	Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020107-011-S_00	North Branch Elm River, downstream to its confluence with the Elm River.	33.4 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation Combination Benthic/Fishes Bioassessments	Low Low	Yes Yes
ND-09020107-014-S_00	Red River of the North from its confluence with the Elm River, downstream to its confluence with the Marsh River.	29.83 miles	Fish Consumption	Not Supporting			
ND-09020109-001-S_00	Goose River from a tributary upstream from Hillsboro, ND downstream to its confluence with the Red River of the North.	27.68 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Methylmercury	Low	No
			Recreation	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes
ND-09020109-007-S_00	North Branch Goose River, downstream to its confluence with the Goose River.	37.12 miles	Fish and Other Aquatic Biota	Not Supporting	Fecal Coliform	Low	No
ND-09020109-011-S_00	Goose River from its confluence with Beaver Creek, downstream to its confluence with the South Branch Goose River.	19.38 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
ND-09020109-015-S_00	South Branch Goose River downstream to its confluence with the Middle Branch Goose River.	33.35 miles	Fish and Other Aquatic Biota	Not Supporting	Combination Benthic/Fishes Bioassessments Sedimentation/Siltation	Low Low	Yes Yes
ND-09020109-017-S_00	Middle Branch Goose River, from its confluence with a tributary watershed near Sherbrooke, ND (ND-09020109-019-S_00), downstream to its confluence with the South Branch Goose River.	17.99 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
			Fish and Other Aquatic Biota	Fully Supporting But Threatened	Combination Benthic/Fishes Bioassessments	Low	Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020109-024-S_00	Beaver Creek from the Golden Lake Diversion, downstream to its confluence with the Goose River.	24.81 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020109-027-S_00	Beaver Creek, downstream to the Golden Lake diversion channel.	37.01 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation Fishes Bioassessments	Low Low	Yes Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020109-029-S_00	Spring Creek, including tributaries	123.75 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-09020109-034-S_00	Little Goose River, from Little Goose River National Wildlife Refuge, downstream to the Goose River.	28.64 miles	Fish and Other Aquatic Biota	Not Supporting	Fishes Bioassessments Sedimentation/Siltation	Low Low	Yes Yes
ND-09020201-006-L_00	Devils Lake	117697 acres	Fish Consumption	Not Supporting	Methylmercury	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-09020202-001-L_00	Warsing Dam	53.4 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved Nutrient/Eutrophication Biological Indicators Sedimentation/Siltation	Low Low Low	No No No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-09020202-001-S_00	Sheyenne River from its confluence with the Warsing Dam Watershed, downstream to the end of the hydrologic unit. Located along the Benson and Eddy County line.	8.9 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020202-004-S_00	Sheyenne River from its confluence with Big Coulee (ND-09020202-007-S_00), downstream to its confluence with the Warsing Dam Watershed (ND-09020202-003-S)	40.37 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	High	Yes
ND-09020202-006-S_00	Sheyenne River from Harvey Dam, downstream to its confluence with Big Coulee (ND-09020202-007-S_00). Located near the Pierce, Benson and Wells County junction.	35.06 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	High	Yes
ND-09020202-012-S_00	Sheyenne River from Coal Mine/Sheyenne Lakes downstream to Harvey Dam. Located along the Sheridan and Wells County border.	20.8 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-09020202-013-S_00	Unnamed tributary watershed to the Sheyenne River (ND-09020202-012-S). Located in Eastern Sheridan County.	36.24 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-09020202-015-S_00	Sheyenne River, downstream to Sheyenne Lake. Located in North Central Sheridan County.	16.7 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-09020203-001-L_00	Lake Ashtabula	5467 acres	Recreation	Not Supporting	Nutrient/Eutrophication Biological Indicators	High	No
ND-09020203-001-S_00	Sheyenne River from Tolna Dam outlet (ND-09020203-020-S) downstream to Lake Ashtabula. Located in Southern Nelson and Eastern Griggs County.	93.81 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020203-002-S_00	Baldhill Creek from tributary watershed (ND-09020203-005-S_00) downstream to Lake Ashtabula. Located in Griggs and Barnes County.	30.21 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020203-004-S_00	Silver Creek, including Gunderson Creek and all tributaries. Located in southern Griggs County.	38.51 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-09020203-007-L_00	McVile Dam	36.7 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
					Sedimentation/Siltation	Low	No
					Oxygen, Dissolved	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-09020203-008-L_00	Tolna Dam	152 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
					Oxygen, Dissolved	Low	No
					Sedimentation/Siltation	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-09020203-008-S_00	Unnamed tributary watershed to Baldhill Creek (ND-09020203-007-S). Located in NW Griggs County.	16.07 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-09020203-009-S_00	Unnamed tributaries to Baldhill Creek (ND-09020203-007-S). Located in eastern Foster and western Griggs County.	30.5 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020203-012-S_00	Pickereel Lake Creek, including all tributaries. Located in NE Griggs County.	28.04 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-09020203-013-S_00	Unnamed tributary watershed to the Sheyenne River (ND-09020203-001-S). Located in northern Griggs County.	33.92 miles	Recreation	Not Supporting	Fecal Coliform	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020203-018-S_00	Sheyenne River, downstream to the Tolna Dam outlet (ND-09020203-020-S). Located in Benson, Eddy, and Nelson Counties.	56.61 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation Fishes Bioassessments	Low Low	Yes Yes
ND-09020204-001-S_00	Sheyenne River, from its confluence with an unnamed tributary watershed (ND-09020204-014-S), downstream to its confluence with the Maple River. Located in SE Cass County.	26.74 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020204-003-L_00	Brewer Lake	117.8 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation Nutrient/Eutrophication Biological Indicators Oxygen, Dissolved	High High High	No No No
ND-09020204-003-S_00	Sheyenne River from its confluence with the Maple River, downstream to its confluence with the Red River Of The North. Located in Eastern Cass County.	19.01 miles	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-09020204-004-S_00	Rush River from its confluence with an unnamed tributary watershed (ND-09020204-011-S), downstream to its confluence with the Sheyenne River.	17.6 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-09020204-007-S_00	Rush River, downstream to an unnamed tributary watershed (ND-09020204-012-S_00). Located in north central Cass County.	41.4 miles	Fish and Other Aquatic Biota	Not Supporting	Combination Benthic/Fishes Bioassessments Sedimentation/Siltation Fecal Coliform	High High Low	Yes Yes No
			Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation Fishes Bioassessments	High High	Yes Yes
			Recreation	Not Supporting	Fecal Coliform	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020204-015-S_00	Sheyenne River, from its confluence with tributary watershed (ND-09020204-016-S_00), downstream to tributary ND-09020204-014-S_00. Located along the Richland and Cass County border.	28.03 miles					
			Fish and Other Aquatic Biota	Fully Supporting But Threatened			
					Sedimentation/Siltation	Low	Yes
			Recreation	Fully Supporting But Threatened			
					Fecal Coliform	Low	No
ND-09020204-017-S_00	Sheyenne River from unnamed tributary (ND-09020204-018-S_00), downstream to unnamed tributary watershed (ND-09020204-016-S_00). Located in northern Ransom and Richland County.	57.5 miles					
			Fish and Other Aquatic Biota	Fully Supporting But Threatened			
					Benthic-Macroinvertebrate Bioassessments	Low	Yes
					Sedimentation/Siltation	Low	Yes
ND-09020204-022-S_00	Sheyenne River from tributary near Lisbon (ND-09020204-0024-S_00), downstream to its confluence with Dead Colt Creek (ND-09020204-021-S_00). Located in central Ransom County.	11.5 miles					
			Fish and Other Aquatic Biota	Fully Supporting But Threatened			
					Fishes Bioassessments	Low	Yes
			Recreation	Fully Supporting But Threatened			
					Fecal Coliform	Low	No
ND-09020204-023-S_00	Tiber Coulee, including all tributaries. Located in south central Ransom County.	32.7 miles					
			Recreation	Fully Supporting But Threatened			
					Fecal Coliform	Low	No
ND-09020204-025-S_00	Sheyenne River, from its confluence with a tributary near Highway 46 (ND-09020204-025-S_00) downstream to its confluence with a tributary near Lisbon, ND (ND-09020204-024-S_00).	46.96 miles					
			Recreation	Fully Supporting But Threatened			
					Fecal Coliform	Low	No
ND-09020204-027-S_00	Sheyenne River, from its confluence with a tributary watershed below Valley City (ND-09020204-028-S_00), downstream to its confluence with a tributary near Highway 46 (ND-09020204-026-S_00). Located in south central Barnes County.	34.04 miles					
			Fish and Other Aquatic Biota	Fully Supporting But Threatened			
					Sedimentation/Siltation	Low	Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020204-034-S_00	Sheyenne River from its confluence with a tributary above Valley City, near railroad bridge, (ND-09020204-038-S_00) downstream to its confluence with a tributary below Valley City (ND-09020204-028-S_00). Located in Central Barnes County.	14.73 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Benthic-Macroinvertebrate Bioassessments	Low	Yes
			Recreation	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
					Fecal Coliform	Low	No
ND-09020204-040-S_00	Sheyenne River from Lake Ashtabula downstream to its confluence with a tributary above Valley City, near rail road bridge (ND-09020204-038-S_00). Located in Central Barnes County.	13.41 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-09020205-001-S_00	Maple River, from its confluence with Buffalo Creek downstream to its confluence with the Sheyenne River. Located in Eastern Cass County.	27.92 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation	Low	Yes
					Combination Benthic/Fishes Bioassessments	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020205-010-S_00	Maple River, from its confluence with a tributary near Leonard, ND (ND-09020205-011-S_00) downstream to its confluence with Buffalo Creek. Located in South Central Cass County.	48.9 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
					Fishes Bioassessments	Low	Yes
ND-09020205-012-S_00	Maple River from its confluence with the South Branch Maple River downstream to its confluence with a tributary near Leonard, ND. Located in S.W. Cass County.	26.15 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020205-015-S_00	Maple River from its confluence with a tributary watershed near Buffalo, ND (ND-09020205-019-S_00) downstream to its confluence with the South Branch Maple River. Located in Western Cass County.	40.06 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes
ND-09020205-024-S_00	Maple River downstream to its confluence with a tributary near the Steele, Cass, and Barnes County Line (ND-09020205-023-S_00)	28.28 miles	Fish and Other Aquatic Biota	Not Supporting	Oxygen, Dissolved Fishes Bioassessments	Low Low	No Yes
ND-09020301-001-S_00	Red River of the North, from its confluence with the Marsh River, downstream to its confluence with the Sand Hill River. Located in Eastern Trail County.	21.35 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020301-002-S_00	English Coulee from its confluence with a tributary upstream from Grand Forks, ND downstream to its confluence with the Red River of the North (Lower Reach).	5.53 miles	Fish and Other Aquatic Biota	Not Supporting	Total Dissolved Solids Sedimentation/Siltation Oxygen, Dissolved	High High High	No Yes No
			Recreation	Not Supporting	Fecal Coliform Sedimentation/Siltation	High High	No Yes
ND-09020301-007-S_00	Red River of the North from its confluence with the Sand Hill River, downstream to its confluence with Cole Creek.	31.13 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020301-010-S_00	Red River of the North from its confluence with Cole Creek, downstream to its confluence with the Red Lake River.	8.06 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020301-014-S_00	Red River of the North from its confluence with the Red Lake River, downstream to its confluence with English Coulee.	4.02 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
					Methylmercury	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020306-001-S_00	Red River of the North from its confluence with English Coulee, downstream to the confluence with Grand Marais Creek.	8.65 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020306-003-S_00	Red River of the North from its confluence with Grand Marais River, downstream to its confluence with the Turtle River.	12.62 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020306-004-S_00	Red River of the North from its confluence with the Turtle River, downstream to its confluence with the Forest River.	31.94 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020306-005-S_00	Red River of the North from its confluence with the Forest River, downstream to its confluence with the Park River.	22.02 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020307-001-L_00	Larimore Dam (TR #9)	76 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-09020307-001-S_00	Turtle River from its confluence with Salt Water Coulee, downstream to its confluence with the Red River of the North.	30.36 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium	Low	No
					Sedimentation/Siltation	Low	Yes
					Selenium	Low	No
			Municipal and Domestic	Fully Supporting But Threatened	Cadmium	Low	No
					Chloride	Low	No
					Arsenic	Low	No
					Selenium	Low	No
ND-09020307-006-S_00	Turtle River from its confluence with Kelly Slough, downstream to its confluence with Salt Water Coulee.	0.65 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium	Low	No
					Sedimentation/Siltation	Low	Yes
					Selenium	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020307-007-S_00	Fresh Water Coulee from its confluence with Salt Water Coulee downstream to its confluence with the Turtle River.	6.5 miles	Fish and Other Aquatic Biota	Not Supporting	Cadmium Selenium	Low Low	No No
ND-09020307-016-S_00	Kelly Slough from the control structure at Kelly Slough National Wildlife Refuge downstream to its confluence with the Turtle River.	2.69 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium Selenium	Low Low	No No
ND-09020307-019-S_00	Turtle River from its confluence with a tributary NE of Turtle River State Park, downstream to its confluence with Kelly Slough.	25.27 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium Selenium Combination Benthic/Fishes Bioassessments	Low Low Low	No No Yes
ND-09020307-021-S_00	Turtle River from its confluence with South Branch Turtle River downstream to its confluence with A tributary NE of Turtle River State Park.	13.9 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium Selenium	Low Low	No No
			Municipal and Domestic	Fully Supporting But Threatened	Arsenic Selenium	Low Low	No No
ND-09020307-024-S_00	South Branch Turtle River downstream to Larimore Dam.	18.42 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Selenium Cadmium	Low Low	No No
ND-09020307-031-S_00	North Branch Turtle River from its confluence with Whiskey Creek, downstream to its confluence with South Branch Turtle River.	15.26 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium Selenium Fishes Bioassessments	Low Low Low	No No Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020308-001-L_00	Fordville Dam	197 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-09020308-001-S_00	Forest River from Lake Ardoch, downstream to its confluence with the Red River of the North.	16.17 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation Benthic-Macroinvertebrate Bioassessments	Low Low	Yes Yes
ND-09020308-002-L_00	Whitman Dam	143 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	Yes
ND-09020308-003-L_00	Matejcek Dam	130 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	Yes
ND-09020308-015-S_00	Forest River from its confluence with South Branch Forest River, downstream to its confluence with a tributary near Highway 18.	13.26 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes
ND-09020308-023-S_00	Middle Branch Forest River from Matejcek Dam, downstream to its confluence with North Branch Forest River.	8.85 miles	Fish and Other Aquatic Biota	Not Supporting	Fishes Bioassessments	Low	Yes
ND-09020310-001-L_00	Homme Dam	194 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators Sedimentation/Siltation	Low Low	No No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020310-001-S_00	Park River from its confluence with Salt Lake Outlet (ND-09020310-009-S_00), downstream to its confluence with the Red River of the North.	15.06 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium Selenium Lead Copper	Low Low Low Low	No No No No
ND-09020310-010-S_00	Park River from its confluence with a tributary east of Grafton, ND (ND-09020310-012-S_00), downstream to its confluence with the outlet from Salt Lake (ND-09020310-009-S_00).	14.68 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Lead Selenium Copper Cadmium	Low Low Low Low	No No No No
ND-09020310-013-S_00	Park River from the confluence of the South Branch Park River and the Middle Branch Park River, downstream to its confluence with a tributary east of Grafton, ND (ND-09020310-012-S_00).	6.83 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Copper Lead Selenium Cadmium	Low Low Low Low	No No No No
ND-09020310-020-S_00	South Branch Park River from its confluence with a tributary watershed near Adams, ND (ND-09020310-022-S_00), downstream to Homme Dam.	16.9 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes
ND-09020310-029-S_00	Middle Branch Park River from a tributary near Highway 32, downstream to tributary near Highway 18.	26.18 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes
ND-09020310-039-S_00	North Branch Park River from a dam near Milton, ND downstream to its confluence with a tributary near Highway 32.	15.52 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020311-001-S_00	Red River of the North from its confluence with the Park River, downstream to its confluence with a small tributary north of Drayton, ND.	19.02 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020311-003-S_00	Red River of the North from its confluence with a small tributary north of Drayton, ND downstream to its confluence with Two Rivers.	30.3 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020311-005-S_00	Red River of the North from its confluence with Two Rivers, downstream to its confluence with the Pembina River.	17.99 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020311-007-S_00	Red River of the North from its confluence with the Pembina River, downstream to the US/Canada border.	3 miles	Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-09020313-001-S_00	Pembina River from its confluence with the Tongue River downstream to its confluence with the Red River of the North	8.76 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium	Low	No
					Copper	Low	No
					Selenium	Low	No
					Lead	Low	No
			Municipal and Domestic	Fully Supporting But Threatened	Lead	Low	No
					Arsenic	Low	No
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020313-002-L_00	Renwick Dam	220 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No

Table VI-2(cont.). 2008 List of Section 303(d) TMDL Waters for the Red River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-09020313-006-S_00	Tongue River from its confluence with a tributary N.E. of Cavalier, ND downstream to its confluence with Big Slough.	22.54 miles	Fish and Other Aquatic Biota	Not Supporting	Sedimentation/Siltation Combination Benthic/Fishes Bioassessments	Low Low	Yes Yes
ND-09020313-009-S_00	Tongue River from Renwick Dam, downstream to a tributary N.E. of Cavalier, ND.	15.91 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Combination Benthic/Fishes Bioassessments Sedimentation/Siltation	Low Low	Yes Yes
ND-09020313-021-S_00	Pembina River from its confluence with a tributary west of Neche, ND downstream to its confluence with the Tongue River.	32.72 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Cadmium Copper Lead Selenium Sedimentation/Siltation Benthic-Macroinvertebrate Bioassessments	Low Low Low Low Low Low	No No No No Yes Yes
			Municipal and Domestic	Fully Supporting But Threatened	Lead Arsenic	Low Low	No No
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-09020313-023-S_00	Pembina River from its confluence with a tributary N.E. of Walhalla, ND downstream to its confluence with a tributary west of Neche, ND.	36.97 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes
ND-09020313-025-S_00	Pembina River from its confluence with Little South Pembina River, downstream to its confluence with a tributary N.E. of Walhalla, ND.	13.09 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Fishes Bioassessments	Low	Yes




Figure VI-2. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Upper Red River Basin




Figure VI-3. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Lower Red River Basin.

Table VI-3. 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10110101-001-L_00	Powers Lake	950.6 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	High	No
					Sedimentation/Siltation	High	No
					Nutrient/Eutrophication Biological Indicators	High	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-10110101-019-L_00	McGregor Dam	54.3 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	High	No
					Nutrient/Eutrophication Biological Indicators	High	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-10110101-021-L_00	Lake Sakakawea	368231 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Temperature, water	Low	No
					Oxygen, Dissolved	Low	No
			Fish Consumption	Not Supporting	Methylmercury	Low	No
ND-10110101-056-S_00	Handy Water Creek, including all tributaries. Located in Eastern McKenzie County.	42.41 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
ND-10110101-080-S_00	Little Knife River from Stanley Reservoir, downstream to Lake Sakakawea. Located in Central Mountrail County.	45.44 miles	Recreation	Not Supporting	Fecal Coliform	Low	Yes
ND-10110102-001-S_00	Little Muddy River from its confluence with East Fork Little Muddy River, downstream to Lake Sakakawea. Located in Central Williams County.	24 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	Yes

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10110102-003-L_00	Blacktail Dam	160 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	High	No
					Sedimentation/Siltation	High	No
					Nutrient/Eutrophication Biological Indicators	High	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-10110203-001-S_00	Little Missouri River from its confluence with Little Beaver Creek downstream to its confluence with Deep Creek. Located in Slope County.	75.79 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10110203-003-S_00	Deep Creek from the confluences of East Branch Deep Creek and West Brach Deep Creek downstream to its confluence with the Little Missouri River. Located in Slope County.	42.51 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10110203-004-S_00	West Branch Deep Creek, including tributaries. Located in Slope County.	117.25 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10110203-025-S_00	Little Missouri River from its confluence with Deep Creek, downstream to its confluence with Andrew's Creek. Located in Billings and Slope Counties.	48.25 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-10110205-001-S_00	Little Missouri River from its confluence with Beaver Creek downstream to highway 85. Located in McKenzie County.	58.94 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10110205-033-S_00	Little Missouri River from HWY 85 downstream to its confluence with Cherry Creek. Located in McKenzie and Dunn Counties.	23.79 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130101-002-L_00	Brush Lake	200 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
			Recreation	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
					Nutrient/Eutrophication Biological Indicators	Low	No
ND-10130101-002-S_00	Square Butte Creek from its confluence with Otter Creek downstream to its confluence with the Missouri River.	1.79 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened			
			Recreation	Not Supporting	Sedimentation/Siltation	Low	Yes
					Fecal Coliform	Low	No
ND-10130101-003-L_00	Crooked Lake	375 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
					Nutrient/Eutrophication Biological Indicators	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-10130101-009-S_00	Square Butte Creek from Nelson Lake downstream to its confluence with Otter Creek. Located in Oliver and Morton Counties.	38.15 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened			
			Recreation	Not Supporting	Sedimentation/Siltation	Low	Yes
					Fecal Coliform	Low	No
ND-10130103-003-L_00	Braddock Lake	69.5 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
					Sedimentation/Siltation	Low	No
					Nutrient/Eutrophication Biological Indicators	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-10130103-007-S_00	Hay Creek downstream to its confluence with Apple Creek. Located in Burleigh County.	15.78 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened			
					Sedimentation/Siltation	Low	Yes

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130103-010-L_00	Lake Isabel	805.7 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
					Oxygen, Dissolved	Low	No
					Nutrient/Eutrophication Biological Indicators	Low	No
			Recreation	Fully Supporting But Threatened			
ND-10130104-001-L_00	Beaver Lake	953.1 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
					Sedimentation/Siltation	Low	No
					Nutrient/Eutrophication Biological Indicators	Low	No
			Recreation	Fully Supporting But Threatened			
ND-10130104-001-S_00	Beaver creek from its confluence with Sand Creek downstream to Lake Oahe. Located in Emmons County.	8.43 miles	Recreation	Fully Supporting But Threatened			
ND-10130104-003-S_00	Beaver Creek from its confluence with Spring Creek downstream to its confluence with Sand Creek. Located in Emmons County.	14.9 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130104-004-S_00	Sand Creek and tributaries, located in Emmons County.	108.56 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130104-005-S_00	Spring Creek and tributaries, located in Emmons County.	63.14 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130104-007-S_00	Beaver Creek from its confluence with the South Branch Beaver Creek downstream to its confluence with Spring Creek. Located in Emmons County.	37.68 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130104-008-S_00	Clear Creek and tributaries, located in Emmons County.	108.95 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
					Fecal Coliform	High	No

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130104-010-S_00	Beaver Creek from Beaver Lake downstream to its confluence with the South Branch Beaver Creek. Located in Emmons and Logan Counties.	38.92 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130104-012-S_00	Unnamed tributary on the south side of Beaver Lake, Logan and McIntosh Counties.	158.02 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130104-014-S_00	South Branch Beaver Creek from its confluence with the South Branch Beaver Creek Watershed (ND-10130104-015-S) downstream to its confluence with Beaver Creek. Located in McIntosh and Emmons Counties.	43.45 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130201-002-S_00	Knife River from its confluence with Antelope Creek downstream to its confluence with the Missouri River.	19.83 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130201-003-S_00	Knife River from its confluence with Spring Creek downstream to its confluence with Antelope Creek. Located in Mercer County.	17.83 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130201-014-S_00	Antelope Creek from its confluence with East Branch Antelope Creek Watershed (ND-10130201-016-S) downstream to its confluence with the Knife River. Located in Mercer County.	8.57 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-10130201-016-S_00	East Branch Antelope Creek upstream from Antelope Creek, including tributaries. Located in Mercer County.	83.04 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-10130201-017-S_00	Antelope Creek main stem downstream to its confluence with East Branch Antelope Creek Watershed (ND-10130201-016-S). Located in Mercer County.	21.32 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-10130201-035-S_00	Knife River from its confluence with Coyote Creek downstream to its confluence with Spring Creek. Located in Mercer County.	14.65 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130201-036-S_00	Brush Creek and tributaries, located in Mercer and Oliver Counties.	61.06 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130201-037-S_00	Coyote Creek from its confluence with Beaver Creek downstream to its confluence with the Knife River. Located in Mercer County.	17.24 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130201-042-S_00	Knife River from its confluence with Branch Knife River downstream to its confluence with Coyote Creek. Located in Dunn and Mercer Counties.	35.99 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130201-045-S_00	Elm Creek and tributaries, located in Mercer County.	137.89 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130201-046-S_00	Willow Creek and tributaries, located in Mercer County.	29.54 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130202-001-L_00	Lake Tschida	5018 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-10130202-050-S_00	Heart River from Patterson Lake, downstream to its confluence with the Green River. Located in Stark County.	24.7 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
ND-10130203-002-L_00	Crown Butte Dam	31.2 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	High	No
					Nutrient/Eutrophication Biological Indicators	High	No
			Recreation	Fully Supporting But Threatened	Oxygen, Dissolved	High	No
					Nutrient/Eutrophication Biological Indicators	High	No

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130203-005-L_00	Sweetbriar Reservoir	270.6 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-10130203-007-L_00	Danzig Dam	147.5 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
					Sedimentation/Siltation	Low	No
					Nutrient/Eutrophication Biological Indicators	Low	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	No
ND-10130204-007-S_00	Cannonball River from its confluence with Sheep Creek downstream to its confluence with Snake Creek. Located in Grant County.	46.7 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130204-014-S_00	Thirty Mile Creek from its confluence with Springs Creek downstream to its confluence with the Cannonball River. Located in Hettinger County.	39.97 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10130204-017-S_00	Thirty Mile Creek from tributary watershed (ND-10130204-019-S_00), downstream to its confluence with Springs Creek. Located in Hettinger County.	19.75 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10130204-032-S_00	Cannonball River from its confluence with Philbrick Creek downstream to its confluence with Indian Creek. Located in Hettinger and Slope County.	54.25 miles	Recreation	Fully Supporting But Threatened	Escherichia coli	Low	No
ND-10130204-044-S_00	Dead Horse Creek, including all tributaries. Located in Hettinger County.	40.18 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130205-001-S_00	Cedar Creek from its confluence with Hay Creek, downstream to its confluence with the Cannonball River. Located on border of Grant and Sioux Counties.	40.3 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130205-003-L_00	Cedar Lake	198.5 acres	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10130205-006-S_00	Crooked Creek, including all tributaries. Located in Grant County.	40.68 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130205-021-S_00	Plum Creek, including all tributaries. Located in Adams County.	79.34 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10130205-024-S_00	Cedar Creek from its confluence with Chanta Peta Creek, downstream to its confluence with Duck Creek. Located in Adams County.	67.56 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130205-033-S_00	Cedar Creek from Cedar Lake, downstream to its confluence with Chanta Peta Creek. Located in Adams County.	43.06 miles	Fish and Other Aquatic Biota	Not Supporting	Benthic-Macroinvertebrate Bioassessments	Low	Yes
ND-10130205-042-S_00	Cedar Creek from its confluence with South Fork Cedar Creek, downstream to Cedar Lake. Located in Slope and Bowman County.	30.86 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
			Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130205-043-S_00	North Fork Cedar Creek, including all tributaries. Located in Slope County.	14.5 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130205-044-S_00	Unnamed tributaries to Cedar Creek (ND-10130205-042-S_00). Located in Slope and Bowman Counties.	81.25 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10130205-045-S_00	South Fork Cedar Creek, including all tributaries. Located in Bowman County.	21.99 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10130205-046-S_00	Cedar Creek upstream from its confluence with South Fork Cedar Creek, including all tributaries. Located in Bowman and Slope Counties.	49.23 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10130205-047-S_00	North Cedar Creek, including all tributaries. Located in Slope County.	115.13 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10130206-001-S_00	Cannonball River from its confluence with Dogtooth Creek, downstream to Lake Oahe. Border of Morton and Sioux County.	20.83 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130206-007-S_00	Cannonball River from its confluence with a tributary watershed near Shields, ND (ND-10130206-028-S_00), downstream to its confluence with Dogtooth Creek.	21.15 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10130206-027-S_00	Cannonball River from Cedar Creek, downstream to a tributary near Shields, ND.	23.52 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10130303-001-L_00	Mirror Lake	63.3 acres	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
			Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
					Sedimentation/Siltation	High	No
					Oxygen, Dissolved	High	No
			Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No

Table VI-3(cont.). 2008 List of Section 303(d) TMDL Waters for the Missouri River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10130303-001-S_00	Flat Creek, downstream to Mirror Lake. Located in Adams County.	21.03 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10130303-003-S_00	Flat Creek from Mirror Lake downstream to the ND-SD border. Located in Adams County.	24.11 miles					




Figure VI-4. Graphic Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Lake Sakakawea/Missouri River Basin.




Figure VI-5. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the Lake Oahe/Missouri River Basin.

Table VI-4. 2008 List of Section 303(d) TMDL Waters for the James River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10160001-002-L_00	Jamestown Reservoir	2073.4 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No
ND-10160001-002-S_00	James River downstream from Jamestown Reservoir to its confluence with Pipestem Creek.	3.33 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Benthic-Macroinvertebrate Bioassessments	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160001-003-S_00	James River from Arrowwood Lake, downstream to Mud Lake.	3.01 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved	Low	No
ND-10160001-013-S_00	James River from its confluence with Big Slough, downstream to its confluence with Rocky Run.	20.27 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160001-015-S_00	Rocky Run from its confluence with Rosefield Slough downstream to its confluence with the James River.	10.2 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160001-018-S_00	Rocky Run from its confluence with a tributary watershed west of Cathay, ND, downstream to its confluence with Rosefield Slough.	14.53 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160001-021-S_00	Rocky Run from its beginning, downstream to its confluence with a tributary watershed located west of Cathay, ND (ND-10160001-020-S_00).	24.3 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160001-023-S_00	James River from its confluence with Rocky Run, downstream to its confluence with Lake Juanita Outlet (ND-10160001-027-S_00)	21.94 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160002-001-L_00	Pipestem Reservoir	1877 acres	Recreation	Fully Supporting But Threatened	Nutrient/Eutrophication Biological Indicators	High	No

Table VI-4(cont.). 2008 List of Section 303(d) TMDL Waters for the James River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10160002-001-S_00	Pipestem Creek, from its beginning, downstream to Sykeston Dam (Lake Hiawatha).	25.21 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160002-007-S_00	Pipestem Creek from Pipestem Dam #3 (ND-10160002-005-L_00), downstream to its confluence with Little Pipestem Creek.	7.22 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-10160002-008-S_00	Little Pipestem Creek, downstream to its confluence with Pipestem Creek.	24.28 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160002-010-S_00	Pipestem Creek from its confluence with Little Pipestem Creek, downstream to Pipestem Dam #4 (ND-10160002-006-L_00).	29.22 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160002-012-S_00	Unnamed tributary watershed to Pipestem Creek (ND-10160002-013-S_00).	40.74 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160002-013-S_00	Pipestem Creek from Pipestem Dam #4 (ND-10160002-006-L_00), downstream to Pipestem Reservoir.	21 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160003-001-S_00	James River from its confluence with Pipestem Creek, downstream to its confluence with Seven Mile Coulee.	13.04 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Oxygen, Dissolved Ammonia (Un-ionized)	High High	No No
ND-10160003-003-S_00	Cottonwood Creek, downstream to Lake LaMoure.	67.67 miles	Recreation	Not Supporting	Fecal Coliform	Low	No
ND-10160003-008-S_00	Buffalo Creek from its beginning, downstream to its confluence with Beaver Creek (ND-10160003-005-S_00).	32.0 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No

Table VI-4(cont.). 2008 List of Section 303(d) TMDL Waters for the James River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10160003-013-S_00	Seven Mile Coulee, including all tributaries. Located in Eastern Stutsman County.	39.87 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160003-025-S_00	Bone Hill Creek, downstream to its confluence with the James River.	39.33 miles	Recreation	Not Supporting	Fecal Coliform	High	No
ND-10160003-029-S_00	James River from its confluence with Bone Hill Creek, downstream to its confluence with Cottonwood Creek.	38.65 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	Low	No
ND-10160003-032-S_00	Bear Creek from tributary watershed (ND-10160003-035-S_00), downstream to its confluence with the James River.	30.35 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160003-034-S_00	Bear Creek, upstream from tributary watershed (ND-10160003-035_00), including all tributaries.	58.42 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160003-035-S_00	Unnamed tributary watershed to Bear Creek.	33.36 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
ND-10160004-001-S_00	Elm River from Pheasant Lake, downstream to the ND/SD border and Elm Lake.	5.56 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-002-S_00	Maple River from its confluence with South Fork Maple River, downstream to the ND/SD border.	41.59 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-005-S_00	Elm River, downstream to Pheasant Lake. Located in Dickey County.	13.79 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-006-S_00	Upper Elm River, including all tributaries. Located in Dickey County.	15.24 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes

Table VI-4(cont.). 2008 List of Section 303(d) TMDL Waters for the James River Basin in North Dakota.

Assessment Unit ID	AU Description	AU Size	Designated Use	Use Support	Impairment	Priority	5A
ND-10160004-007-S_00	Bristol Gulch, including all tributaries. Located in Dickey County.	45.93 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-008-S_00	Unnamed tributaries to the Elm River (ND-10160004-005-S_00). Located in Dickey County.	21.69 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-009-S_00	Unnamed tributary to Pheasant Lake. Located in Dickey County.	2.53 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-013-S_00	Maple River from its confluence with Maple Creek, downstream to its confluence with South Fork Maple River. Located in Dickey County.	15.79 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
ND-10160004-015-S_00	South Fork Maple River from its confluence with three tributaries, downstream to its confluence with the Maple River. Located in Dickey County.	14.53 miles	Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No
			Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
			Recreation	Not Supporting	Fecal Coliform	High	No
ND-10160004-022-S_00	Maple Creek, downstream to its confluence with the Maple River. Located in Lamoure County.	33.91 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
			Recreation	Not Supporting	Fecal Coliform	High	No
ND-10160004-026-S_00	Maple River from Schlect-Thom Dam, downstream to its confluence with Maple Creek. Located in Lamoure County.	20.01 miles	Fish and Other Aquatic Biota	Fully Supporting But Threatened	Sedimentation/Siltation	Low	Yes
			Recreation	Fully Supporting But Threatened	Fecal Coliform	High	No




Figure VI-6. Graphical Depiction of 2008 Section 303(d) Listed Waters Needing TMDLs in the James River Basin.

Table VI-5. 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

Assessment Unit ID/Description	AU Size	Impaired Use	Pollutant	Rationale for De-listing
ND-09010002-002-L_00 - Northgate Dam	150.8 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 22, 2006.
			<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 22, 2006. The TMDL includes a linkage analysis between phosphorus and low dissolved oxygen that provides justification that meeting the phosphorus target will also meet the dissolved oxygen TMDL target.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 22, 2006. The TMDL report also includes de-listing justification for the sediment/siltation impairment suggesting the narrative criteria for sediment is currently being met.
			<i>Recreation</i>	
			<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 22, 2006.
ND-09010003-001-L_00 - Carbury Dam	130 acres	<i>Fish and Other Aquatic Biota</i>	<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL includes a linkage analysis between phosphorus and low dissolved oxygen that provides justification that meeting the phosphorus target will also meet the dissolved oxygen TMDL target.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL report also includes de-listing justification for the sediment/siltation impairment suggesting the narrative criteria for sediment is currently being met.
			<i>Recreation</i>	
			<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-09010003-003-S_00 - Wintering River, including all tributaries. Located in SW McHenry and NE McLean counties.	195.9 miles	<i>Fish and Other Aquatic Biota</i>	<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. An analysis of available suspended sediment, TSS, and biological monitoring data does not support suspended sediment as a cause of aquatic life use impairment.
ND-09010003-005-S_00 - Souris River from its confluence with the Wintering River downstream to its confluence with Willow Creek. Located in NE McHenry County.	76.2 miles	<i>Fish and Other Aquatic Biota</i>	<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to change in WQS. Based on the revised dissolved oxygen standard whereby up to 10% of representative samples during any 3-year period can exceed the standard, the water quality standard is now being attained.
ND-09010004-002-L_00 - Long Lake	287 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on the most recent water quality assessment data collected in 2006 as part of the North Dakota Game and Fish Dept District Lake Water Quality Assessment Project, the lake is assessed as mesotrophic which based on the state's assessment methodology means both aquatic life and recreation uses are met.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; threatened water no longer threatened. Based on the most recent water quality assessment data collected in 2006 as part of the North Dakota Game and Fish Dept District Lake Water Quality Assessment Project, the lake is assessed as mesotrophic which based on the state's assessment methodology means both aquatic life and recreation uses are met.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-09020105-002-L_00 - Mooreton Pond	36.8 acres	<i>Fish and Other Aquatic Biota</i>	<i>Turbidity</i>	Applicable WQS attained; according to new assessment method. Based on recent monitoring data collected by the Department and the ND Game and Fish Dept the lake is assessed as mesotrophic. Therefore, the previous TMDL listing for turbidity is no longer valid. The lake remains listed as impaired for aquatic life due to elevated TDS which is negatively affecting the growth and reproduction of fish.
ND-09020109-002-L_00 - South Golden Lake	323.5 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and the trend is improving, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to change in WQS. Based on the trophic status assessment for this waterbody and a revision to the dissolved oxygen standard as it applies to Class 3 lakes which became effective October 1, 2006 the lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions in the hypolimnion.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and the trend is improving, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-09020202-002-L_00 - Balta Dam	108 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to change in WQS. Based on the trophic status assessment for this waterbody and a revision to the dissolved oxygen standard as it applies to Class 3 lakes which became effective October 1, 2006 the lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions in the hypolimnion.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
ND-09020203-004-L_00 - Red Willow Lake	130 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to change in WQS. Based on the trophic status assessment for this waterbody and a revision to the dissolved oxygen standard as it applies to Class 3 lakes which became effective October 1, 2006 the lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions in the hypolimnion.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-09020204-004-S_00 - Rush River from its confluence with an unnamed tributary watershed (ND-09020204-011-S), downstream to its confluence with the Sheyenne River.	17.44 miles	<i>Fish and Other Aquatic Biota</i>	<i>Organic Enrichment (Sewage) Biological Indicators</i>	Applicable WQS attained; original basis for listing was incorrect. Based on 96 dissolved oxygen measurements taken between 2006 and 2007 as part of the water quality assessment project there is no evidence of organic enrichment or low (less than 5 mg/L) oxygen concentrations in the river.
ND-09020204-005-L_00 - Dead Colt Creek Dam	124 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL for this waterbody was completed and approved by EPA on September 22, 2006.
			<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL was completed and approved by EPA on September 22, 2006. The TMDL contains a linkage analysis between phosphorus loading and low dissolved oxygen in reservoirs. It is anticipated that meeting the phosphorus load reduction will result in the dissolved oxygen target being
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. A nutrient (phosphorus) and dissolved oxygen TMDL was completed and approved by EPA on September 22, 2006. The TMDL report also includes de-listing justification for the sediment/siltation impairment suggesting the narrative criteria for sediment is currently being met.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL for this waterbody was completed and approved by EPA on September 22, 2006.
ND-09020204-007-S_00 - Rush River, downstream to an unnamed tributary watershed (ND-09020204-012-S_00). Located in north central Cass County.	40.92 miles	<i>Fish and Other Aquatic Biota</i>	<i>Organic Enrichment (Sewage) Biological Indicators</i>	Applicable WQS attained; original basis for listing was incorrect. Based on 96 dissolved oxygen measurements taken between 2006 and 2007 as part of the water quality assessment project there is no evidence of organic enrichment or low (less than 5 mg/L) oxygen concentrations in the river.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-09020307-001-S_00 - Turtle River from its confluence with Salt Water Coulee, downstream to its confluence with the Red River of the North.	30.36 miles	<i>Fish and Other Aquatic Biota</i>	<i>Total Dissolved Solids</i>	Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). Based on the water quality assessment that was conducted by the Grand Forks County SCD, there is no basis for the listing of the aquatic life impairment due to TDS. There is no existing water quality standard for TDS and therefore, no basis for the original assessment and TMDL listing. The waterbody has been listed as impaired for aquatic life due to several trace metals and for drinking water due to exceedences of the state chloride standard.
ND-09020307-006-S_00 - Turtle River from its confluence with Kelly Slough, downstream to its confluence with Salt Water Coulee.	0.65 miles	<i>Fish and Other Aquatic Biota</i>	<i>Total Dissolved Solids</i>	Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). Based on the water quality assessment that was conducted by the Grand Forks County SCD, there is no basis for the listing of the aquatic life impairment due to TDS. There is no existing water quality standard for TDS and therefore, no basis for the original assessment and TMDL listing.
ND-09020308-001-S_00 - Forest River from Lake Ardoch, downstream to its confluence with the Red River of the North.	16.17 miles	<i>Fish and Other Aquatic Biota</i>	<i>Total Dissolved Solids</i>	Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). Based on limited biological and water quality data for this waterbody, there is no basis for the listing of the aquatic life impairment due to TDS. There is no existing water quality standard for TDS and therefore, no basis for the original assessment and TMDL listing. The waterbody remains listed as impaired for aquatic life due to biological indicators and suspended sediment. The waterbody has also be identified as Category 5A, therefore additional monitoring will be conducted in the next 2-4 years to verify aquatic life impairments.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-09020313-007-L_00 - Lake Upsilon	414 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to change in WQS. Based on the trophic status assessment for this waterbody and a revision to the dissolved oxygen standard as it applies to Class 3 lakes which became effective October 1, 2006 the lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions in the hypolimnion.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. Based on water quality assessment data collected by the North Dakota Game and Fish Department's Fisheries Division, there is no basis for the listing of the aquatic life impairment due to sedimentation/siltation. There is also no existing water quality standard for suspended sediment and therefore, no basis for the original assessment and TMDL listing. The waterbody has been assessed as eutrophic and therefore also de-listed as impaired for aquatic life due to nutrients and low dissolved oxygen.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

Assessment Unit ID/Description	AU Size	Impaired Use	Pollutant	Rationale for De-listing
ND-09020313-011-L_00 - Armourdale Dam	79.8 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007.
			<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL includes a linkage analysis between phosphorus and low dissolved oxygen that provides justification that meeting the phosphorus target will also meet the dissolved oxygen TMDL target.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL report also includes de-listing justification for the sediment/siltation impairment suggesting the narrative criteria for sediment is currently being met.
			<i>Recreation</i>	
			<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007
ND-10130101-004-L_00 - Strawberry Lake	140 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now mesotrophic and a trend that is improving, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; according to new assessment method. Based on the trophic status assessment for this waterbody and a revision to the dissolved oxygen standard as it applies to Class 3 lakes which became effective October 1, 2006 the lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions in the hypolimnion.
			<i>Recreation</i>	
			<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now mesotrophic and a trend that is improving, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-10130103-014-L_00 - McDowell Dam	55.2 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 27, 2007. Following the completion of the TMDL, the lake was restored with an alum treatment in May 2006. Intensive monitoring to date shows the lake is now mesotrophic and is fully supporting its recreational uses. Low water levels have hampered efforts to fully restore the lakes fishery.
			<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 27, 2007. The TMDL includes a linkage analysis between phosphorus and low dissolved oxygen that provides justification that meeting the phosphorus target will also meet the dissolved oxygen TMDL target.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on September 27, 2007. Following the completion of the TMDL, the lake was restored with an alum treatment in May 2006. Intensive monitoring to date shows the lake is now mesotrophic and is fully supporting its recreational uses. Low water levels have hampered efforts to fully restore the lakes fishery.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is improving, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
ND-10130106-002-L_00 - Green Lake	868.6 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on the trophic status assessment for this waterbody and recent monitoring data collected by the North Dakota Game and Fish Department which shows no recent exceedences of the dissolved oxygen standard as it applies to Class 2 lakes, lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; threatened water no longer threatened. Based on the trophic status assessment for this waterbody and recent monitoring data collected by the North Dakota Game and Fish Department which shows no recent exceedences of the dissolved oxygen standard as it applies to Class 2 lakes, lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is improving, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; threatened water no longer threatened. Based on the trophic status assessment for this waterbody and recent monitoring data collected by the North Dakota Game and Fish Department which shows no recent exceedences of the dissolved oxygen standard as it applies to Class 2 lakes, lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

Assessment Unit ID/Description	AU Size	Impaired Use	Pollutant	Rationale for De-listing
ND-10130106-003-L_00 - Lake Hoskins	553.5 acres	Fish and Other Aquatic Biota	Nutrient/Eutrophication Biological Indicators	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL was completed and approved by EPA on September 22, 2006.
			Oxygen, Dissolved	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL was completed and approved by EPA on September 22, 2006. The TMDL contains a linkage analysis between phosphorus loading and low dissolved oxygen in reservoirs. It is anticipated that meeting the phosphorus load reduction will result in the dissolved oxygen target being
		Recreation		
			Nutrient/Eutrophication Biological Indicators	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL was completed and approved by EPA on September 22, 2006.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

Assessment Unit ID/Description	AU Size	Impaired Use	Pollutant	Rationale for De-listing
ND-10130202-004-L_00 - Dickinson Dike	22 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; due to restoration activities. Through a lake restoration project sponsored by the North Dakota Game and Fish Department's "Save Our Lakes" Program, Dickinson Dike (known also as Queen City Dam) was dredged thereby deeping and removing accumulated sediments and associated nutrients. Recent monitoring data collected by the North Dakota Game and Fish Department in 2005 and 2006 shows the lake is now mesotrophic and is meeting all of its designated beneficial uses.
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to restoration activities. Through a lake restoration project sponsored by the North Dakota Game and Fish Department's "Save Our Lakes" Program, Dickinson Dike (known also as Queen City Dam) was dredged thereby deeping and removing accumulated sediments and associated nutrients. Recent monitoring data collected by the North Dakota Game and Fish Department in 2005 and 2006 shows the lake is now mesotrophic and is meeting all of its designated beneficial uses.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; due to restoration activities. Through a lake restoration project sponsored by the North Dakota Game and Fish Department's "Save Our Lakes" Program, Dickinson Dike (known also as Queen City Dam) was dredged thereby deeping and removing accumulated sediments and associated nutrients. Recent monitoring data collected by the North Dakota Game and Fish Department in 2005 and 2006 shows the lake is now mesotrophic and is meeting all of its designated beneficial uses.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; due to restoration activities. Through a lake restoration project sponsored by the North Dakota Game and Fish Department's "Save Our Lakes" Program, Dickinson Dike (known also as Queen City Dam) was dredged thereby deeping and removing accumulated sediments and associated nutrients. Recent monitoring data collected by the North Dakota Game and Fish Department in 2005 and 2006 shows the lake is now mesotrophic and is meeting all of its designated beneficial uses.
ND-10130204-001-L_00 - Sheep Creek Dam	84.4 acres	<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A nutrient (phosphorus) TMDL for this waterbody was completed and approved by EPA on May 28, 2008.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

Assessment Unit ID/Description	AU Size	Impaired Use	Pollutant	Rationale for De-listing
ND-10130204-006-L_00 - Indian Creek Dam	222 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007.
			<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL includes a linkage analysis between phosphorus and low dissolved oxygen that provides justification that meeting the phosphorus target will also meet the dissolved oxygen TMDL target.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL report also includes de-listing justification for the sediment/siltation impairment suggesting the narrative criteria for sediment is currently being met.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007.
ND-10130204-047-S_00 - North Fork Cannonball River from its confluence with White Lake Watershed (ND-10130204-049-S_00), downstream to its confluence with Philbrick Creek. Located in Slope County.	33.25 miles	<i>Recreation</i>	<i>Fecal Coliform</i>	Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). There are no fecal coliform data available to support the current listing. The current TMDL listing will be verified through additional monitoring before a TMDL is completed.
ND-10130204-051-S_00 - Philbrick Creek from its confluence with Adobe Wall Creek, downstream to its confluence with the Cannonball River. Located in Slope County.	11.7 miles	<i>Recreation</i>	<i>Fecal Coliform</i>	Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). There are no fecal coliform data available to support the current listing. The current TMDL listing will be verified through additional monitoring before a TMDL is completed.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-10130205-003-L_00 - Cedar Lake	198.5 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication. The lake remains listed as a Category 5A waterbody for sedimentation/siltation
			<i>Oxygen, Dissolved</i>	Applicable WQS attained; due to change in WQS. Based on the trophic status assessment for this waterbody and a revision to the dissolved oxygen standard as it applies to Class 3 lakes which became effective October 1, 2006 the lake has also been de-listed for aquatic life impairments due to low dissolved oxygen conditions in the hypolimnion.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	Applicable WQS attained; threatened water no longer threatened. Based on recently collected monitoring data and an assessment that shows the lake is now eutrophic and a trend that is stable, the waterbody has been de-listed for both aquatic life and recreational use impairments due to nutrient enrichment/eutrophication. The lake remains listed as a Category 5A waterbody for sedimentation/siltation
ND-10160002-005-S_00 - Pipestem Creek from Sykston Dam, downstream to a small impoundment known as Pipestem Dam #3 (ND-10160002-005-L_00) which is located near the Wells/Foster County Line .	10.53 miles	<i>Recreation</i>	<i>Fecal Coliform</i>	Applicable WQS attained; threatened water no longer threatened. Based on 54 fecal coliform samples collected in 2006 and 2007 as part of the water quality implementation project there is no evidence of a recreational use impairment due to fecal coliform bacteria.
ND-10160004-003-S_00 - Weber Gulch, including all tributaries. Located in Dickey County.	114.75 miles	<i>Recreation</i>	<i>Fecal Coliform</i>	Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). There are no data available for this waterbody to support the recreational use impairment due to fecal coliform bacteria. The current listing was presumably made by extrapolating data collected from a downstream waterbody.

Table VI-5(cont.). 2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.

<i>Assessment Unit ID/Description</i>	<i>AU Size</i>	<i>Impaired Use</i>	<i>Pollutant</i>	<i>Rationale for De-listing</i>
ND-10160004-005-L_00 - Pheasant Lake	232.1 acres	<i>Fish and Other Aquatic Biota</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007.
			<i>Oxygen, Dissolved</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL includes a linkage analysis between phosphorus and low dissolved oxygen that provides justification that meeting the phosphorus target will also meet the dissolved oxygen TMDL target.
			<i>Sedimentation/Siltation</i>	Applicable WQS attained; original basis for listing was incorrect. A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007. The TMDL report also includes de-listing justification for the sediment/siltation impairment suggesting the narrative criteria for sediment is currently being met.
		<i>Recreation</i>	<i>Nutrient/Eutrophication Biological Indicators</i>	TMDL approved or established by EPA (4A). A TMDL for nutrients (phosphorus) and dissolved oxygen was completed and approved by EPA on February 22, 2007.

PART VII. GROUND WATER ASSESSMENT

A. Ground Water Extent and Uses

Chapter 1. Aquifer Description

Ground water underlies the land surface throughout all of North Dakota and is present in both unconsolidated deposits and bedrock. Unconsolidated deposits are loose beds of sand, gravel, silt or clay that are of glacial origin. Aquifers in the unconsolidated deposits are called glacial drift aquifers and are the result of glacial outwash deposits. Glacial drift aquifers are generally more productive than aquifers found in the underlying bedrock and provide better quality water. Approximately 206 glacial drift aquifers have been identified and delineated throughout the state. The locations and aerial extent of the major glacial drift aquifers in the state are shown in Figure VII-1. It is estimated that 60 million acre-feet (AF) of water are stored in the major glacial drift aquifers in the state.




Figure VII-1. Major Glacial Drift Aquifers in North Dakota.

The bedrock underlying North Dakota consists primarily of shale and sandstone that generally (except in southwestern North Dakota) underlie the unconsolidated deposits. Bedrock aquifers underlie the entire state and tend to be more continuous and widespread than glacial drift aquifers. Water contained within bedrock aquifers occurs primarily along fractures in the rock, and the water produced is generally more mineralized and saline than water from glacial drift aquifers. The major bedrock aquifers that underlie North Dakota are shown in Figure VII-2. The amount of water available in the bedrock aquifers is unknown.




Figure VII-2. Location and Extent of North Dakota's Primary Bedrock Aquifers.

North Dakota has completed a multi-agency effort to assess and map the major ground water resources found within the state's boundaries. The County Ground Water Studies Program provides a general inventory of the state's ground water resources and was completed through a cooperative effort of the North Dakota State Water Commission (SWC), the North Dakota Geological Survey, the United States Geological Survey, county water resource districts and county commission boards. The county ground water studies identified the location and extent of major aquifers, hydraulic properties of the aquifers, water chemistry, estimated well yields and the occurrence and movement of ground water, including sources of recharge and discharge. The county studies were prepared in three parts:

- Part I describes the geology.
- Part II provides basic ground water data, including descriptive lithologic logs of test holes and wells, water levels in observation wells and water chemistry analyses.
- Part III describes the general hydrogeology.

The County Ground Water Studies are available for all counties in North Dakota. The SWC and other federal and state agencies continue to evaluate the ground water resources and expand the available knowledge of the quantity and quality of these resources.

Chapter 2. Ground Water Use

Ground water use in North Dakota has historically been categorized as agricultural (irrigation or livestock watering), industrial and domestic (private or public). Ninety-four percent of the incorporated communities in the state rely on ground water from private wells, municipal distributions systems and/or rural water systems. Ground water is virtually the sole source of all water used by farm families and residents of small communities having no public water distribution system.

As indicated in Table VII-1, the highest consumptive use of ground water is related to irrigation.

Table VII-1. 2003 Reported Ground Water Use in North Dakota.

Type of Water Use	Amount of Water Used (acre-feet)	Percent of Total Water Used (%)
Irrigation	111,581	61
Municipal	27,782	15
Livestock	17,589	10
Rural Water Systems/Other	10,479	6
Industrial	9,648	5
Rural Domestic	5,887	3
Total	182,966	100

Notes: 1 acre-foot = 325,850 gallons

Data was obtained from the North Dakota State Water Commission website.

B. Ground Water Contamination Sources

Chapter 1. Contaminant Source Description

Contamination of ground water from manmade and natural sources has been detected in every county of the state. The degree to which contamination incidents are investigated or remediated is a function of the contaminant, its impact on the beneficial use of the resource and the overall risk it poses to the public or the environment. The following are the highest priority contaminant sources which have caused adverse impacts on the beneficial use of ground water resources throughout the state:

- Agricultural chemical facilities
- Animal feedlots
- On-farm agricultural mixing and loading procedures
- Above ground and underground storage tanks
- Surface impoundments
- Large industrial facilities
- Spills and releases

Common contaminants associated with these facilities include organic pesticides, nitrates, halogenated solvents, petroleum hydrocarbon compounds, sulfates, chlorides and total dissolved solids.

Chapter 2. Ground Water Contaminant Source Databases

The major sources of ground water contamination were determined utilizing a combination of professional experience and a review of existing department computer databases. Several databases maintained by the Division of Water Quality compile information relating to the type of regulated activity, its size and location and, in some cases, regional ground water quality information. The primary databases used to identify the major sources of ground water contamination are:

Concentrated Animal Feeding Operations (CAFO) Database

Since 1972, North Dakota has maintained an active concentrated animal feeding operations (CAFO) permit program. The program is designed to protect the quality of the state's water resources through oversight of the construction and management of CAFOs. The program regulates animal feeding operations and can require design or operational modifications to protect the quality of the waters of the state. Regulatory authority is provided in North Dakota Century Code (NDCC) 61-28 and North Dakota Administrative Code (NDAC) 33-16, which can require specific actions for construction, water quality monitoring, animal disposal, contingency planning and animal waste disposal. The CAFO database provides location, operation and contact information. The database is updated as needed to reflect changes in the program, such as the approval of new operations or modifications to existing operations. At present, information regarding 715 facilities is listed in the CAFO database.

Underground Injection Control (UIC) Program Class V Database

The Underground Injection Control (UIC) Program regulates the injection of liquid waste into the ground where it may have the potential to adversely impact underground sources of drinking water. The department has regulatory primacy to oversee and enforce the Class I and Class V UIC Programs. As part of this effort, the department completed a statewide survey designed to identify the type, location and use of small industrial or commercial injection systems. The UIC Class V database was developed to catalog information obtained during the survey and is updated as needed. At present, approximately 2,500 sites are in the database, with a total of 564 identified as facilities that have discharged or currently discharge waste fluids into a Class V underground injection well.

Spill Response/Contaminant Release Database

The department maintains databases which track the initial response and subsequent follow-up action at locations where contaminants released to the environment impact water quality. Site location, contaminant type, responsible party and a historical record of activities conducted at the site are maintained.

Ambient Ground Water Quality Monitoring Database

The Ambient Ground Water Quality Program was developed to monitor ground water quality in the 50 most vulnerable aquifers in the state. In general, vulnerability was determined based upon natural geologic conditions, total appropriated water use and land use. The program was originally designed to identify the occurrence of about 60 different pesticides in ground water. New pesticides are added from time to time in response to increased production of specialty crops and/or new pest infestations. The Ambient Ground Water Quality Database contains all the data obtained through the implementation of the monitoring program. This includes sample location, analytical results and other site-specific information.

C. Ground Water Protection Programs

In 1967, North Dakota enacted legislation enabling the state regulation of activities which have caused, or which have the potential to cause, adverse impacts to the quality of the waters of the state. NDCC 61-28 entitled, “Control, Prevention and Abatement of Pollution of Surface Waters,” not only defines the statement of policy for surface and ground water quality protection, but also sets specific prohibitions and penalties for violation of the state law. Since the enactment of NDCC 61-28, the state has pursued a policy to:

“...act in the public interest to protect, maintain and improve the quality of the waters of the state for continued use as public and private water supplies, propagation of wildlife, fish and aquatic life and for domestic, agricultural, industrial and recreational and other legitimate beneficial uses....”

North Dakota has historically envisioned ground water quality protection to include a mix of financial and technical cooperation among federal, state and local governmental agencies and private entities. Since the early 1970s, the department has continued to build upon existing ground water protection capacities through the attainment of primacy for federal programs or through cooperative working relationships with other state, federal and local entities.

The following are brief descriptions of the programs administered by the department’s Division of Water Quality.

Chapter 1. Wellhead and Source Water Protection Programs

The 1996 Amendments to the Safe Drinking Water Act established the Source Water Protection Program to serve as an overall umbrella of protection efforts for all public water systems, including ground water- and surface water-dependent systems. In North Dakota, the Wellhead Protection Program focuses on the ground water-dependent systems, while the Source Water Protection Program addresses surface water-dependent systems. The Source Water Protection Program involves the delineation of a protection area along rivers or reservoirs that provide source water for the system and an inventory of potential contaminant sources within the protection area. Under both wellhead and source water protection, the department assesses the system’s susceptibility to potential contaminant sources found in the protection area.

The 1996 Amendments to the Safe Drinking Water Act required all states to complete the minimum elements of wellhead and source water protection (delineation, contaminant source inventory and susceptibility) by May 2003. The department completed the mandatory elements for all of the Community Water Systems and all of the Non-community Water Systems in the state by the required deadline.

North Dakota continues to promote and implement the Source Water Assessment Program. Public water systems are encouraged to implement the voluntary elements of wellhead and source water protection. These elements include the development of management strategies, contingency planning and public awareness programs. The department works with, and provides assistance to, all public water systems who desire to follow through with the voluntary elements of the program.

Following the completion of source water assessment requirements in 2003, the Wellhead Protection Program began conducting source water monitoring and contaminant source studies for ground water-dependent community public water systems that have been rated as susceptible or for systems that have had detections of organic or inorganic contaminants regulated by the Safe Drinking Water Act National Primary Drinking Water Regulations. Source water monitoring typically involves the use of existing monitoring wells at contaminant release sites or the use of private water supply wells in or near the wellhead protection area. Source water monitoring is accomplished through coordination with the local public water system and the department's divisions of Municipal Facilities and Waste Management.

D. Ground Water Quality

Chapter 1. Ambient Ground Water Monitoring Program

Ambient ground water quality monitoring activities are conducted by several agencies, with the primary activities being conducted by the North Dakota SWC and the department. The monitoring programs have been developed to assess ground water quality and/or quantity in the major aquifer systems located throughout the state. The monitoring is designed to evaluate the condition of ground water quality as it relates to inorganic/organic chemical constituents and the occurrence of selected agricultural chemical compounds. Additional water quality information is collected as part of the Safe Drinking Water Act requirements through the monitoring of public drinking water programs.

The maintenance of a baseline description of ground water quality is an essential element of any statewide comprehensive ground water protection program. In recent years, concern for the quality of North Dakota's environment and drinking water has increased as it is learned that many states in the country have experienced ground water contamination from a variety of point and nonpoint sources of pollution.

In North Dakota, a large portion of the potable ground water resource underlies agricultural areas. Prior to the inception of the Ambient Ground Water Monitoring Program in 1992, only limited data were available to assess the impact of agricultural chemicals on the state's ground water quality. The goal of the Ambient Ground Water Monitoring Program is to provide an assessment of the quality of North Dakota's ground water resources with regard to agricultural chemical contamination.

Several glacial drift aquifers have been monitored each year of the program since 1992. The monitoring conducted in 1996 marked the completion of the first five-year cycle of monitoring high-priority glacial drift aquifers in the state. The second five-year cycle of monitoring began in 1997, during which time the aquifers sampled five years earlier in 1992 were resampled. The third five-year cycle of monitoring was completed in 2006. Conducting the monitoring on five-year cycles, preferably using most of the same wells for sampling, will provide a temporal assessment of agricultural chemical occurrence in specific aquifers. Results of each year's monitoring are described in annual ground water monitoring reports.

Chapter 2. Underground Injection Control (UIC) Program

The department's Class I and V Underground Injection Control (UIC) Programs have been administered in accordance with UIC rules and program descriptions. Program activities include administration of the program grant, permitting, surveillance and inspections, quality assurance, enforcement, data management, public participation, training, technical assistance and Class V assessment activities. The current UIC inventory includes four active Class I wells and 545 active Class V injection wells of various subclasses. The UIC Program coordinates with other programs, including the Resource Conservation and Recovery Act (RCRA), Underground Storage Tank (UST), National Pollutant Discharge Elimination System (NPDES) and Wellhead/Source Water Protection to identify activities which may threaten ground water quality.

Chapter 3. Additional Ground Water-Related Projects

Ground Water Protection Program staff work on many projects related to the protection of the ground water resources of North Dakota. Projects include special monitoring projects; review of sites for livestock feeding operations; review of sites for landfill operations; and working on emergency response, investigations and cleanup of releases to the environment.

Facility Location Reviews

The Ground Water Protection Program takes the lead or assists other programs and agencies in evaluating the impacts land use activities may have on ground water quality. Site reviews or preliminary site reviews are conducted for new feedlot or CAFO operations, landfill or waste disposal facilities and industrial facilities. The Ground Water Protection Program also conducts special monitoring projects at CAFO facilities in the state to evaluate/identify potential ground water quality changes. In addition, site reviews are conducted for on-site sewage systems in new residential subdivisions to assess potential ground water impacts.

Water Appropriation and Monitoring

The department reviews approximately 40 water appropriation permits each year to assess potential impacts to ground water quality. Proposed water uses includes agricultural, public water supply, recreational and industrial uses. A cooperative project with the SWC is underway involving the Karlsruhe aquifer to identify causes and potential solutions to nitrate increases in irrigated areas. Meetings were conducted with SWC personnel and local residents to discuss survey results and ongoing research. Currently, voluntary measures such as BMPs and reduced nutrient application rates are being implemented and evaluated in these areas. One of the irrigators has voluntarily installed shallow recovery/production wells to recover nitrate in the area of highest contamination. Residential drinking water wells are being monitored to ensure there is no danger to public health.

Contaminant Release Sites

The Ground Water Protection Program coordinates with the UST Program, RCRA/Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) Program and the Drinking Water Program to provide technical oversight relating to the assessment and remediation of ground water contamination incidents. The majority of sites are related to fuel storage facilities, although other types of storage sites include pesticides, nutrients/fertilizers, chlorinated solvents, metals and trace metals, and other inorganic compounds.

Pesticide Use Exemption Evaluations

The department also reviews applications for pesticide use exemptions (Federal Insecticides, Fungicides and Rodenticides Act Section 18 Requests) for potential impacts to surface or ground water. Approximately six requests are reviewed each year, and comments regarding each request are provided to the North Dakota Department of Agriculture.

Emergency Response and Spills

Additional project oversight is provided by the Ground Water Protection Program staff for a wide variety of emergency response and release incidents. The Ground Water Protection Program provides technical assistance to the Division of Emergency Management to address potential water quality impacts from accidental or intentional releases. The department continues to work with the North Dakota Oil and Gas Division on response to oilfield spills, using the one-stop online spill reporting capabilities which were added to the department web site, with automatic notification to appropriate department personnel. The Ground Water Protection Program also provides oversight or technical comment either directly to the responsible party or through the appropriate oversight agency on other ground water contamination projects. Typical projects include sites that require one or more of the following activities: site assessment, selection and implementation of appropriate corrective action, and sample collection and data review/evaluation.

REFERENCES

- Brinson, M.M., 1993, *A Hydrogeomorphic Classification for Wetlands*, technical report WRP-DE-4, U.S. Army Corps of Engineer Waterways Experiment Station, Vicksburg, MS.
- Carlson, R.E., 1997, *A Trophic Status Index for Lakes, Limnology and Oceanography*, Vol. 22 (Issue 2), pp. 361-364.
- Cowardin, L.M., V. Carter, F.C. Golet and E.T. LaRoe, 1979, *Classification of Wetlands and Deepwater Habitats of the United States*, U.S. Fish and Wildlife Service, Office of Biological Services – 79/31.
- , D.S. Gilmes and L.M. Mechlin, 1981, “*Characteristics of Central North Dakota Wetlands Determined from Sample Aerial Photographs and Ground Study*,” The Wildlife Society Bulletin 9 (4): pp. 280-288.
- Dahl, T.E., 1990, *Wetlands - Losses in the United States: 1780's to 1980's*, Washington, D.C., U.S. Fish and Wildlife Service Report to Congress.
- EPA, July 2005, *Guidance for 2006 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d), 305(b) and 314 of the Clean Water Act*, Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans and Watershed, U.S. Environmental Protection Agency, Washington, DC.
- Karr, J.R. and D. R. Dudley, 1981, *Ecological Perspectives on Water Quality Goals*, Env. Mgmt. 5:44-68.
- Lee, L.S. and Mark M. Brinson, William J. Kleindl, P. Michael Whited, Michael Gilbert, Wade L. Nuttons, Dennis F. Whigham, Dave DeWald, 1997, *Operational Draft Guidebook for the Hydrogeomorphic Assessment of Temporary and Seasonal Prairie Pothole Wetlands*.
- Martin, A.C., N. Hotchkiss, F.M. Uhles and W.S. Bourn, 1953, *Classification of Wetlands in the U.S.*, U.S. Fish and Wildlife Service, Resource Publication 92.
- NDDoH, 2005, *North Dakota's Water Quality Monitoring Strategy for Surface Waters: 2005-2014*, North Dakota Department of Health, Bismarck, ND.
- , 2006, *Standards of Quality for Waters of the State* (revised), North Dakota Century Code 33-16-02.1., North Dakota Department of Health, Bismarck, ND.
- Omernik, J. M. 1987. Ecoregions of the conterminous United States. *Annals of the Association of American Geographers* 77, 118-125.

Pearson, Eric and M.J. Ell, 1997, *Effects of Rising Reservoir Water Levels Resulting from the 1993 Flood on the Methyl-Mercury Concentrations in Fish Tissues in Lake Sakakwea, ND*, North Dakota Department of Health, Bismarck, ND.

Reynolds, Ron, 1997, personal communication, project leader, HAPET Office, U.S. Fish and Wildlife Service, Bismarck, ND.

Stewart, R.E. and H.A. Kantrud, 1971, *Classification of Natural Ponds and Lakes in the Glaciated Prairie Region*, Resource Publication 92, Bureau of Sport Fisheries and Wildlife, U.S. Fish and Wildlife Service, Washington, D.C.

----- and H.A. Kantrud, 1973, *Ecological Distribution of Breeding Waterfowl Populations in North Dakota*, Journal of Wildlife Management 37(1): 39-50.

Stoddard, J.L., Peck, D.V., Olsen, A.R., Larsen, D.P., Van Sickle, J., Hawkins, C.P., Hughes, R.M., Whittier, T.R., Lomnický, G.A., Herlihy, A.T., Kaufman, P.R., Peterson, S.A., Ringold, P.L., Paulsen, S.G., and Blair, R. 2005a. Environmental Monitoring and Assessment Program (EMAP) western streams and rivers statistical summary: U.S. Environmental Protection Agency Report 620/R-05/006, 1,762 p.

Appendix A

North Dakota Water Quality Assessment Methodology for Surface Waters

Water Quality Assessment Methodology for North Dakota's Surface Waters



**North Dakota Department of Health
Division of Water Quality**

Water Quality Assessment Methodology
for North Dakota's Surface Waters

John Hoeven, Governor
Terry Dwelle, M.D., State Health Officer



North Dakota Department of Health
Division of Water Quality
Gold Seal Center, 4th Floor
918 East Divide Avenue
Bismarck, ND 58501-1947

701.328.5210

Table of Contents	Page
I. Introduction	1
A. Background	1
B. North Dakota Surface Water Resources.....	1
C. Purpose and Scope	2
II. Water Quality Standards	3
A. Background	3
B. Beneficial Use Designation.....	3
C. Numeric Water Quality Standards	4
D. Narrative Water Quality Standards	5
E. Antidegradation Policies and Procedures.....	6
III. Assessment Database	6
IV. Sufficient and Credible Data Requirements and Overwhelming Evidence.....	8
A. Sufficient and Credible Data Requirements.....	8
B. Overwhelming Evidence	9
V. Beneficial Use Assessment Methodology.....	10
A. Aquatic Life Use Assessment Methodology for Rivers and Streams	10
1. Chemical Criteria	10
2. Biological Assessment Criteria.....	11
B. Recreation Use Assessment Methodology for Rivers and Streams	14
C. Aquatic Life and Recreation Use Assessment Methodology for Lakes and Reservoirs	16
D. Drinking Water Supply Use Assessment Methodology for Rivers, Lakes and Reservoirs	18
E. Fish Consumption Assessment Methodology for Rivers, Lakes and Reservoirs.....	20
F. Agricultural Use Assessment Methodology for Rivers, Lakes and Reservoirs.....	21
G. Industrial Use Assessment Methodology for Rivers, Lakes, and Reservoirs.....	22
VI. References.....	22

List of Tables**Page**

1. Assessment Categories for the Integrated Report.....	2
2. Biological Integrity Scoring Criteria and Aquatic Life Use Support Criteria Based on the Lake Agassiz Plain Ecoregion Fish IBI	13
3. Macroinvertebrate IBI Scores for Reference Sites in the Lake Agassiz Plain Ecoregion of North Dakota.....	13
4. Biological Integrity Scoring Criteria and Aquatic Life Use Support Criteria Based on the Lake Agassiz Plain Ecoregion Macroinvertebrate IBI.....	14
5. State Water Quality Standards for Chloride, Sulfate, and Nitrate	19

List of Figures

1. Map of Reach-Indexed Assessments Units Delineated in the Souris River Basin	8
2. A Graphical Representation of Carlson's TSI	18

Appendices

A. North Dakota Water Quality Standards	
---	--

I. INTRODUCTION

A. Background

The federal Clean Water Act (CWA) provides the regulatory context and mandate for state water quality monitoring and assessment programs. The North Dakota Department of Health (NDDoH) has been designated as the state water pollution control agency for purposes of the federal CWA and, as such, is authorized to take all actions necessary or appropriate to secure for the state all benefits of the CWA and similar federal acts (NDCC 61-28-04). State law establishes policy to protect, maintain, and improve the quality of waters of state, while the overall goal of the federal CWA is to “restore and maintain the chemical, physical, and biological integrity of the Nation’s waters.”

Various sections in the CWA require states to conduct specific activities to monitor, assess, and protect their waters. These activities include:

- Develop and adopt water quality standards designed to protect designated beneficial uses (Section 303);
- Establish and maintain monitoring programs to collect and analyze water quality data (Section 106). Reporting on the status of waters and the degree to which designated beneficial uses are supported (Section 305[b]);
- Identify and prioritize waters that are not meeting water quality standards (Section 303[d]);
- Assess the status and trends of water quality in lakes and identifying and classifying lakes according to trophic condition (Section 314); and
- Identify waters impaired due to nonpoint sources of pollution as well as identifying those sources and causes of nonpoint source pollution (Section 319).

B. North Dakota’s Surface Water Resources

The NDDoH currently recognizes 247 lakes and reservoirs for water quality assessment purposes. Of this total, there are 138 manmade reservoirs and 109 natural lakes. The 138 reservoirs have an areal surface of 543,156 acres, which represents about 71 percent of North Dakota’s total lake/reservoir surface acres. Of these, 480,731 acres or 62 percent of the state’s entire lake and reservoir acres are contained within the two mainstem Missouri River reservoirs (Lake Sakakawea and Lake Oahe). The remaining 136 reservoirs share 62,425 acres, with an average surface area of 459 acres. The 109 natural lakes in North Dakota cover 218,616 acres, with approximately 117,697 acres or 54 percent attributed to Devils Lake. The remaining 108 lakes average 934 acres, with half being smaller than 250 acres.

There is an estimated 54,607 miles of rivers and streams in the state. Estimates of river stream miles in the state are based rivers and streams entered into the Assessment Database and reach indexed to the 1:100,000 scale National Hydrography Dataset (NHD).

One of the most significant water resource types in the state are wetlands. There are an estimated 2.5 million acres of wetlands in the state. The majority of these wetlands are temporary, seasonal, semi-permanent, and permanent depressional wetlands located in what is commonly called the Prairie Pothole Region.

C. Purpose and Scope

Water quality standards provide the fundamental benchmarks by which the quality of all surface waters are measured. It is the water quality standards that are used to determine impairment. As a general policy, the assessment procedures described in this methodology are consistent with the NDDoH's interpretation of the state's water quality standards.

For purposes of Section 305(b) reporting and Section 303(d) listing, the US Environmental Protection Agency (EPA) encourages states to submit an integrated report and to follow its integrated reporting guidance (EPA, 2005). Key to integrated reporting is an assessment of all of the state's waters and placement of those waters into one of five assessment categories. The categories represent varying levels of water quality standards attainment, ranging from Category 1, where all of a waterbody's designated uses are fully supporting, to Category 5, where a pollutant impairs a waterbody and a TMDL is required (Table 1). These category determinations are based on consideration of all existing and readily available data and information consistent with the state's water quality assessment methodology. The purpose of this document is to describe the assessment methodology used in the state's biennial integrated report. This information, which is summarized by specific lake, reservoir, river reach or sub-watershed, is integrated as beneficial use assessments that are entered into a water quality assessment "accounting"/database management system developed by EPA. This system, which provides a standard format for water quality assessment and reporting, is termed the Assessment Database (ADB).

Table 1. Assessment Categories for the Integrated Report

Assessment Category	Assessment Category Description
Category 1	All of the waterbody's designated uses have been assessed and are fully supporting.
Category 2	Some of the waterbody's designated uses are fully supporting, but there is insufficient data to determine if remaining designated uses are fully supporting.
Category 3	Insufficient data to determine whether any of the waterbody's designated uses are met.
Category 4	At least one of the waterbody's beneficial uses is not supported or has been assessed as fully supporting, but threatened, but a TMDL is not needed. This category has been further sub-categorized as: <ul style="list-style-type: none"> • 4A - waterbodies that are impaired or threatened, but TMDLs needed to restore beneficial uses have been approved or established by EPA; • 4B - waterbodies that are impaired or threatened, but do not require TMDLs because the state can demonstrate that "other pollution control requirements (e.g., BMPs) required by local, state or federal authority" • (see 40 CFR 130.7[b][1][iii]) are expected to address all waterbody-pollutant combinations and attain all water quality standards in a reasonable period of time; and • 4C - waterbodies that are impaired or threatened, but the impairment is not due to a pollutant.
Category 5	At least one of the waterbody's beneficial uses is not supported or has been assessed as fully supporting, but threatened, and a TMDL is needed.

II. WATER QUALITY STANDARDS

A. Background

As stated previously, water quality standards are the fundamental benchmarks by which the quality of all of the state's surface waters are assessed. It is the state's water quality standards that are ultimately used to determine beneficial use impairment status.

Water quality standards were first adopted into North Dakota administrative code beginning in the late 1960's. "Water quality standards" is a term which is used in both a broad and narrow sense. In its broadest sense, water quality standards include all the provisions and requirements in water quality rules and regulations, including minimum wastewater treatment requirements and effluent limits for point source dischargers. In the more narrow sense, water quality standards define the specific uses we make of waters of the state and set forth specific criteria, both numeric and narrative, that define acceptable conditions for the protection of these uses, including antidegradation provisions (Appendix A). The term "water quality standards" is used in the more narrow sense throughout this document.

Water quality reporting requirements under Sections 305(b) and 303(d) of the CWA require states to assess the extent to which their lakes, reservoirs, rivers, and streams are meeting water quality standards applicable to their waters, including beneficial uses as defined in their state water quality standards. In addition to beneficial uses, applicable water quality standards also include narrative and numeric standards and antidegradation policies and procedures. While Section 305(b) requires states and tribes to provide only a statewide water quality summary, Section 303(d) takes this reporting a step further by requiring states to identify and list the individual waterbodies that are not meeting applicable water quality standards and to develop TMDLs for those waters. Both Section 305(b) reporting and Section 303(d) listing accomplish this assessment by determining whether a waterbody is supporting its designated beneficial uses.

B. Beneficial Use Designation

The protected beneficial uses of the state's surface waters are defined in the *Standards of Quality for Waters of the State* (Appendix A). The state's water quality standards provide for four stream classes (I, IA, II, and III) and five lake classes (1-5). While considered "waters of the state" and protected under the state's narrative standards, the state's water quality standards do not define beneficial uses for wetlands.

All classified lakes, reservoirs, rivers, and streams in the state are protected for aquatic life and recreation. Protection for aquatic life means surface waters are suitable for the propagation and support of fish and other aquatic biota, including aquatic macroinvertebrates, and that these waters will not adversely affect wildlife in the area. Protection of all surface waters, except wetlands, for recreation means waters should be suitable for direct body contact activities such as bathing and swimming and for secondary contact activities such as boating, fishing, and wading.

Class I, IA, and II rivers and streams and all classified lakes and reservoirs are designated for use as municipal and drinking water supplies. Specifically, these waters shall be suitable for use as a source of water supply for drinking and culinary purposes after treatment to a level approved by the NDDoH.

While not specifically identified in state water quality standards, fish consumption is protected through both narrative and numeric human health criteria specified in the state's water quality standards (Appendix A). The state's narrative water quality standards provide that surface waters shall be "free from materials attributable to municipal, industrial, or other discharges or agricultural practices" which will "render any undesirable taste to fish flesh or, in any way, make fish inedible." In addition, the state's statewide fish consumption advisory applies to all waters known to provide a sport fishery.

Other beneficial uses identified in the state's water quality standards are agriculture (e.g., stock watering and irrigation) and industrial (e.g., washing and cooling). These uses apply to all classified rivers, streams, lakes, and reservoirs.

Four beneficial uses (aquatic life, recreation, drinking water, and fish consumption) are typically assessed for purposes of Section 305(b) reporting and Section 303(d) listing. All waterbodies included in the assessment database (ADB) and, therefore, all stream classes (I, IA, II, and III) and all lake classes (1-5) are assigned aquatic life and recreation beneficial uses. All Class I, IA, and II rivers and streams and all classified lakes and reservoirs are assigned the drinking water beneficial use. Fish consumption use is assumed to apply to all Class I, IA, and II rivers and streams, to those Class III streams known to provide a sport fishery, and to all Class 1 through 4 lakes and reservoirs.

C. Numeric Water Quality Standards

A numeric water quality standard is considered a safe concentration of a pollutant in water, associated with a specific beneficial use. Numeric standards are associated with all use classes. Ideally, if the numeric standard is not exceeded, the use will be protected. However, nature is very complex and variable, and the NDDoH may use a variety of assessment tools (e.g., chemical and biological monitoring) to fully assess beneficial uses. With few exceptions, protection for aquatic life and/or drinking water uses will also provide protection for less sensitive uses (e.g., agriculture and industrial uses). For some pollutants, numeric standards may be applicable to more than one use and may be more stringent for one use than another. For example, the drinking water standard for selenium is 50 µg/L, while the chronic aquatic life standard is 5 µg/L.

As is the case for most states, the state of North Dakota's numeric standards for toxic pollutants are based on the EPA's aquatic life criteria. The EPA develops and publishes these criteria as required by Section 304(a) of the CWA. Most numeric standards have two parts, a chronic value and an acute value. The chronic standard is the highest concentration of a toxicant to which organisms can be exposed indefinitely with no harmful effects, including growth and reproduction. The acute standard protects aquatic organisms from potential lethal effects of a short-term "spike" in the concentration of the toxicant.

In the development of aquatic life criteria and associated standards, the EPA and the NDDoH have addressed some of the many toxicological, water chemistry, and practical realities that affect a toxicant's impact on aquatic biota. For example, pollutant concentrations and flow volumes vary in effluents and in receiving streams over time, aquatic organisms generally can tolerate higher concentrations of toxicants for shorter periods of time, and the sensitivity of aquatic organisms to toxicants often varies over their lifespan. EPA's approach for expressing water

quality standards addresses varying toxicant concentrations, length of an averaging period for the standard, and the number of acceptable exceedances over time. These concepts are highly relevant to the interpretation of water quality standards and the assessment of waterbodies based on available data. In the development and implementation of numeric water quality standards, these concepts are referred to as:

- Magnitude;
- Duration; and
- Frequency.

Magnitude refers to the concentration of a given pollutant and is represented by the numeric standard. For example, the chronic and acute standards for copper are 14.0 and 9.3 µg/L, respectively. This is the “magnitude” of copper that, if not exceeded in water, will protect aquatic biota from chronic and acute effects.

Duration refers to the period of time the measured concentration of a toxicant can be averaged and still provide the desired level of protection to the aquatic community. In the context of toxicity to aquatic organisms, it would be unrealistic to consider a standard as an instantaneous maximum concentration never to be exceeded. On the other hand, toxicant concentrations averaged over too long a time could be under-protective, if it allowed exceedingly high lethal concentrations to be masked by the average. In general, EPA recommends a 4-day averaging period for chronic standards and a 1-hour averaging period for acute standards.

Frequency refers to the number of times a standard may be exceeded over a prescribed time period and still provide adequate protection. EPA guidance and state water quality standards specify that the numeric standards, both chronic and acute, should not be exceeded more than once in three years. The three year time frame is based on studies of the time it takes for aquatic communities to recover from a major disturbance.

D. Narrative Water Quality Standards

A narrative water quality standard is a statement(s) that prohibits unacceptable conditions from occurring in or upon surface waters, such as floating debris, oil, scum, garbage, cans, trash, or any unwanted or discarded material. Narrative standards also prohibit the discharge of pollutants, which alone or in combination with other substances, can 1) cause a public health hazard or injury to the environment; 2) impair existing or reasonable beneficial uses of surface waters; or 3) directly or indirectly cause concentrations of pollutants to exceed applicable standards. Narrative standards are often referred to as “free froms” because they help keep surface waters free from very fundamental and basic forms of water pollution (e.g., sediment and nutrients).

The association between narrative standards and beneficial use impairment is less well defined than it is for numeric standards. Because narrative standards are not quantitative, the determination that one has been exceeded typically requires a “weight-of-evidence” approach to the assessment showing a consistent pattern of water quality standards violations. The narrative standards relevant to this guidance document are found in state water quality standards Section 33-16-02.1-08 (Appendix A). These standards protect surface waters and aquatic biota from:

- Eutrophication (particularly lakes and reservoirs);

- Impairment of the biological community (exemplified by the Index of Biotic Integrity); and
- Impairment of fish for human consumption.

E. Antidegradation Policies and Procedures

In addition to numeric and narrative standards and the beneficial uses they protect, a third element of water quality standards is antidegradation. The fundamental concept of antidegradation is the protection of waterbodies whose water quality is currently better than applicable standards. Antidegradation policies and procedures are in place to maintain high quality water resources and prevent them from being degraded down to the level of water quality standards.

State water quality standards has established three categories or tiers of antidegradation protection (Appendix A). Category 1 is a very high level of protection and automatically applies to all Class I and IA rivers and streams, all Class 1, 2, and 3 lakes and reservoirs, and wetlands that are functioning at their optimal level. Category 1 may also apply to some Class II and III rivers and streams, but only if it can be demonstrated that there is remaining pollutant assimilative capacity, and both aquatic life and recreation uses are currently being supported. Category 2 antidegradation protection applies to Class 4 and 5 lakes and reservoirs and to Class II and III rivers and streams not meeting the criteria for Category 1. Category 3 is the highest level of protection and is reserved for Outstanding State Resource Waters. Waterbodies may only be designated Category 3 after they have been determined to have exceptional value for present and prospective future use for public water supplies, propagation of fish or aquatic biota, wildlife, recreational purposes, or agricultural, industrial, or other legitimate beneficial uses.

III. ASSESSMENT DATABASE

North Dakota's Assessment Database (ADB) contains 1,709 discreet assessment units (AUs) representing 54,607 miles of rivers and streams and 247 lakes and reservoirs. Within the ADB, designated uses are defined for each AU (i.e., river or stream reach and lake or reservoir) based on the state's water quality standards. Each use is then assessed using available chemical, physical and/or biological data.

With an estimated 54,607 miles of rivers and streams and 761,772 acres of lakes and reservoirs, it is impractical to adequately assess each and every mile of stream or every acre of lake. However, the NDDoH believes it is important to: 1) accurately assess those waters for which beneficial use assessment information is available; and 2) account for those stream miles and lake acres that are not assessed or for which there are insufficient data to conduct an assessment. As a result, the NDDoH has adopted the ADB to manage water quality assessment information for the state's rivers, streams, lakes, and reservoirs.

Developed by EPA, the ADB is an Access[®] based “accounting”/database management system that provides a standard format for water quality assessment information. It includes a software program for adding and editing assessment data and transferring assessment data between the personal computer and EPA. Assessment data, as compared to raw monitoring data, describes the overall health or condition of the waterbody by describing beneficial use impairment and, for those waterbodies where beneficial uses are impaired or threatened, the causes and sources of pollution affecting the beneficial use. The ADB also allows the user to track and report on TMDL-listed waters, including their development and approval status and de-listing rationale.

To create North Dakota’s ADB, the state’s 54,607 miles of rivers and streams and 247 lakes and reservoirs have been delineated into 1,709 discreet AUs. An AU can be an individual lake or reservoir, a specific river or stream reach or a collection of stream reaches in a sub-watershed. North Dakota’s ADB is currently represented by 1,462 river and stream AUs and 247 lake and reservoir AUs. Each of these AUs is then assessed individually, based on the availability of sufficient and credible data. In order to delineate and define AUs used in the ADB, the NDDoH follows a general set of guidelines:

1. Each AU is within the eight-digit USGS hydrologic unit.
2. Each river and stream AU is composed of stream reaches of the same water quality standards classification (I, IA, II or III).
3. To the extent practical, each AU is within a contiguous Level IV ecoregion.
4. Mainstem perennial rivers are delineated as separate AUs. Where these rivers join with another major river or stream within the eight-digit hydrologic unit, the river was further delineated into two or more AUs.
5. Tributary rivers and streams, which are named on USGS 1:100,000 scale planimetric maps or the National Hydrography Dataset (NHD), are delineated as separate AUs. These AUs may be further delineated, based on stream order or water quality standards classification.
6. Unnamed ephemeral tributaries to a delineated AU are consolidated into one unique AU. This is done primarily for accounting purposes so that all tributary stream reaches identified in the NHD are included in the ADB.
7. Stream reaches, which are identified in the NHD and on USGS 1:24,000 scale maps and which do not form either an indirect or direct hydrologic connection with a perennial stream, are not included in the ADB. This would include small drainages that originate and flow into closed basin lakes or wetlands. (Note: These delineation criteria do not apply to tributaries to Devils Lake.)

The ADB provides an efficient accounting and data management system. It also allows for the graphical presentation of water quality assessment information by linking assessments contained in the ADB to the NHD file through “reach indexing” and geographic information systems (GIS). In order to facilitate the GIS data link, the NDDoH has “reach-indexed” each AU in the ADB to the NHD file. The product of this process is a GIS coverage that can be used to

graphically display water quality assessment data entered in the ADB. An example can be seen in Figure 1, which depicts each of the reach-indexed AUs delineated in the Souris River Basin. Assessments completed and entered into the ADB also form the basis for the state's Section 319 Nonpoint Source (NPS) Assessment Report and Management Plan. Because of the way the NDDoH's Surface Water Quality Management Program is structured, there is complete integration of the state's Section 305(b) Water Quality Assessment Report, the Section 303(d) TMDL List and the Section 319 NPS Assessment Report and Management Plan.




Figure 1. Map of Reach-Indexed Assessment Units Delineated in the Souris River Basin.

IV. SUFFICIENT AND CREDIBLE DATA REQUIREMENTS AND OVERWHELMING EVIDENCE

A. Sufficient and Credible Data Requirements

For water quality assessments, including those done for purposes of Section 305(b) assessment and reporting and 303(d) listing, the NDDoH will use only what it considers to be sufficient and credible data. Sufficient and credible data are chemical, physical, and biological data that, at a minimum, meet the following criteria:

- Data collection and analysis followed known and documented quality assurance/quality control procedures.
- Water column chemical or biological data are 10 years old or less for rivers and streams and lakes and reservoirs, unless there is adequate justification to use older data (e.g., land use, watershed, or climatic conditions have not changed). There is no age limit for fish tissue mercury data. Years of record are based on the USGS water year. Water years are from October 1 in one year through September 30 of the following year. It should be noted that it is preferable to split the year in the fall when hydrologic conditions are stable, rather than to use calendar years. Data for all 10 years of the period are not

required to make an assessment.

- There are a minimum of 10 chemical samples collected in the 10-year period for rivers and streams. The 10 samples may range from one sample collected in each of 10 years or 10 samples collected all in one year.
- There should be a minimum of two samples collected from lakes or reservoirs collected during the growing season, May-September. The samples may consist of two samples collected the same year or samples collected in separate years.
- A minimum of five fecal coliform and/or E. coli samples are collected during any calendar month from May through September. The five samples per month may consist of five samples collected during the month in the same year or five samples collected during the same calendar month, but pooled across multiple years (e.g., two samples collected in May 2000, two samples collected in May 2001 and one sample collected in May 2005).
- For all chemical criteria that are expressed as a 30-day arithmetic average (e.g., chloride, sulfate, radium 226 and 228, and boron) a minimum of four daily samples must be collected during any consecutive 30-day period. Samples collected during the same day shall be averaged and treated as one daily sample.
- A minimum of two biological samples (fish and/or macroinvertebrate) are necessary in the most recent 10-year period. Samples may be collected from multiple sites within the assessment stream reach, multiple samples collected within the same year, or individual samples collected during multiple years. Samples may consist of a minimum of two fish samples, two macroinvertebrate samples, or one fish and one macroinvertebrate sample.
- There are a minimum of 5 fish tissue samples per species per lake, reservoir, or river, that represents the range in size classes present in the waterbody.

B. Overwhelming Evidence

There are situations where a single set of data is all that is needed to make a use support determination. For example, a single set of water chemistry data may be sufficient to establish that a waterbody is not supporting aquatic life use. In such situations where a single data set irrefutably proves that impairment exists, an impairment determination may be based on this “overwhelming evidence.”

A number of factors are evaluated when making a determination as to whether data can be used as a basis for an “overwhelming evidence” assessment. Factors include the technical soundness of the methods used to collect the data and the spatial and temporal coverage of the data as it relates to the waterbody being assessed. Data quality and data currency (i.e., how old are the data?) are also factors which are considered.

Data cannot be overwhelming evidence unless the methods used for collection and analysis meets the most stringent standards for reliability and validity. The person evaluating the data must be certain that the data are representative of actual current waterbody conditions. The data

must be representative of the spatial extent of the waterbody and of relevant temporal patterns. Data more than three or four years old should not be used as overwhelming evidence unless there is a strong basis for concluding that conditions have not changed since the data were collected.

V. BENEFICIAL USE ASSESSMENT METHODOLOGY

A. Aquatic Life Use Assessment Methodology for Rivers and Streams

The following is a description of the assessment methodology or decision criteria used to assess aquatic life and recreation uses where they are assigned to rivers and streams in the state. The methodologies used to assess drinking water and fish consumption uses are the same for both rivers and lakes and are provided in separate sections of this document.

All water quality assessments entered into the ADB for Section 305(b) reporting and Section 303(d) TMDL listing are based on “sufficient and credible” monitoring data. Physical and chemical monitoring data used for these assessments includes conventional pollutant (e.g., dissolved oxygen, pH, temperature, ammonia, fecal coliform bacteria, and *E. coli* bacteria) and toxic pollutant (e.g., trace elements and pesticides) data collected for the most recent 10-year period. Biological monitoring data used for assessment includes fish community data collected by the NDDoH from the Red River Basin between 1993 and 1996, macroinvertebrate community data collected throughout the state between 1995 and 2000, and data collected between 2000 and 2004 as part of the EMAP Western Pilot Project.

As stated previously, use impairment for the state’s rivers and streams is assessed for aquatic life and recreation. The following is the beneficial use decision criteria utilized for these assessments.

The NDDoH uses both chemical and biological data when assessing aquatic life use support for the state’s rivers and streams. In some cases, both chemical data and biological data are used to make an assessment determination for an AU. Where both data are available, the NDDoH uses a weight-of-evidence approach in making an assessment decision. For example, if there are chemical data that do not show an aquatic life use impairment, but there are sufficient and credible biological data to show an impairment to the aquatic community, then the use-support decision will be to list the river or stream AU as “not supporting.”

1. Chemical Assessment Criteria

In general, aquatic life use determinations utilizing chemical data are based on the number of exceedances of the current *Standards of Quality for Waters of the State* (Appendix A) for DO, pH, and temperature and on the number of exceedances of the acute or chronic standards for ammonia, aluminum, arsenic, cadmium, copper, cyanide, lead, nickel, selenium, silver, zinc, and chromium. The acute and chronic water quality standards for trace metals are expressed as total recoverable metals and not as dissolved metals. However, where dissolved metals data are available, use support assessments are made by applying the dissolved metals data to the water quality standards expressed as the total recoverable fraction.

The following are the use support decision criteria that the NDDoH uses to assess aquatic life use based on chemical data:

- Fully Supporting:

For the conventional pollutants DO, pH, and temperature, the standards of 5 mg/L (daily minimum) for DO, 7.0 to 9.0 (Class I and IA streams and all lakes) and 6.0 to 9.0 (Class II and III streams) for pH and 29.4 °C (85 °F) (maximum) for temperature are not exceeded in the AU. Consistent with state water quality standards (Appendix A), if the DO or pH standard is exceeded, but in less than 10 percent of the samples and there is no record of lethality to aquatic biota, then the AU is also assessed as “fully supporting.”

For ammonia and other toxic pollutants (e.g., trace elements and organics), aquatic life is assessed as “fully supporting” if the acute or chronic standard is not exceeded during any consecutive three-year period.

- *Fully Supporting but Threatened:*

For DO and pH, one or more standards were exceeded in 11 to 25 percent of the measurements taken during the 10-year assessment period. The temperature standard is exceeded, but in less than 10 percent of the measurements taken during the 10-year assessment period.

For ammonia and other toxic pollutants, the acute or chronic standard was exceeded once or twice during any consecutive three-year period during the 10-year assessment period.

- *Not Supporting:*

For DO and pH, one or more standards were exceeded in more than 25 percent of the measurements taken during the 10-year assessment period. The temperature standard is exceeded in more than 10 percent of the measurements taken during the 10-year assessment period.

For ammonia and other toxic pollutants, the acute or chronic standard was exceeded three or more times during any consecutive three-year period during the 10-year assessment period.

2. Biological Assessment Criteria

Aquatic-life use, or biological integrity, can be defined as “the ability of an aquatic ecosystem to support and maintain a balanced, integrated, adaptive community of organisms having a species composition, diversity and functional organization comparable to that of the natural habitats of the region.” (Karr, 1981) When the aquatic community (e.g., fish and macroinvertebrates) is similar to that of “least disturbed” habitats in the region, termed “reference condition,” aquatic life use can be assessed as fully supporting. When the aquatic community deviates significantly from reference condition, it is assessed as either fully supporting, but threatened or not supporting, depending upon the degree of impairment.

While chemical data provides an indirect assessment of aquatic life use impairment, direct measures of the biological community are believed to be a more accurate assessment of aquatic-life use or biological integrity. The state water quality standards (Appendix A) describe a narrative biological goal that “the biological condition of surface waters shall be similar to that of sites or waterbodies determined by the NDDoH to be regional reference sites.” This narrative standard also states that it is the intent of the state, in adopting this narrative goal, “to provide an additional assessment method that can be used to identify impaired surface waters.”

The NDDoH began a stream biological monitoring and assessment program in 1993. In order to interpret these biological data and to develop a biological assessment methodology, the NDDoH has adopted the “multi-metric” index approach to assess biological integrity or aquatic-life use support for rivers and streams. The multi-metric index approach assumes that various measures of the biological community (e.g., species richness, species composition, trophic structure, and individual health) respond to human-induced stressors (e.g., pollutant loadings or habitat alterations). Each measure of the biological community, termed a “metric,” is evaluated and scored on either a 1-, 3-, 5-point scale (fish) or on a scale of 0-100 (macroinvertebrates). The higher the score, the better will be the biological condition and, presumably, the lower the pollutant or habitat impact.

To date, the NDDoH has developed final multi-metric IBIs (Index of Biotic Integrity) for only fish and macroinvertebrates in the Lake Agassiz Plain ecoregion. The Lake Agassiz Plain ecoregion is a part of the larger Red River of the North Basin. While the NDDoH continues to analyze both fish and macroinvertebrate data from other river basins and ecoregions in the state, including data collected as part of the EMAP Western Pilot Project, the lack of an adequate number of quantifiable reference sites within these regions has limited the analysis of metrics and the development and interpretation of IBIs. As a result, biological assessments based on IBI results are limited to only the Lake Agassiz Plain ecoregion in the state.

Biological Assessment Methods for the Lake Agassiz Plain Ecoregion

The fish IBI was published in a report entitled *Development of Index of Biotic Integrity Expectations for the Lake Agassiz Plain Ecoregion* (EPA, 1998). This IBI is based on 12 metrics and a 1, 3, 5 scoring criteria similar to Karr et al. (1986). This IBI results in a total possible score of 60. Table 2 provides a summary of the IBI scores and their related biological integrity classes (excellent, good, fair, poor, and very poor). Sites with biological integrity classes rated as excellent and good are assessed as fully supporting aquatic life use, while sites that were rated as poor and very poor were assessed as not supporting aquatic life use (Table 2). Sites with a biological condition class rated as fair were not assessed.

The macroinvertebrate IBI for the Lake Agassiz Plain ecoregion was published in the report entitled *Macroinvertebrate Index of Biotic Integrity for the Lake Agassiz Plain Ecoregion (46) of North Dakota* (NDDoH, 2006). This IBI was developed based on 41 samples collected from 33 sites, including five reference site samples.

To determine the biological condition or aquatic life use support of streams, threshold values are required to determine what constitutes good biological condition scores (i.e., fully supporting aquatic life use) or poor biological condition scores (i.e., not supporting

aquatic life use) in a multi-metric index. The assessment approach used for this report is outlined in Barbour et al. (1999). First, the 25th percentile of the five reference sites IBI scores was determined. Based on the reference site macroinvertebrate IBI scores for sites in the Lake Agassiz Plain ecoregion (Table 3), the 25th percentile of reference site IBI scores is 53. This value is equivalent to the dividing line between good and fair biological condition. (Note: This threshold could be set lower if there is more confidence that the reference sites truly represent non-impacted conditions. Since there is usually some doubt about the certainty of reference site population, using values above the 25th percentile was selected as a conservative approach to determine if a value at a site is within the range of reference sites.)

The thresholds between fair, poor, and very poor were then determined by dividing the range below good (0-53) into three parts (0-17.77, 17.78-35.33, and 35.34-52.99). The very poor biological condition range is represented by the lower third of the range of IBI scores from 0-17.77, the poor range by scores ranging from 17.78-35.33 and the fair range by scores ranging from 35.34-52.99. Biological condition scores were then translated into aquatic life use attainment categories by assigning the good biological condition class as fully supporting aquatic life use and the poor and very poor biological condition class as not supporting aquatic life use (Table 4). Due to uncertainty associated with the reference site population, sites classified with a biological condition score of fair should not be assessed (Table 4).

Table 2. Biological Integrity Scoring Criteria and Aquatic Life Use Support Criteria Based on the Lake Agassiz Plain Ecoregion Fish IBI.

Fish IBI Score	Biological Integrity Class	Aquatic Life Use Support
60-51	Excellent	Fully Supporting
50-41	Good	Fully Supporting
40-31	Fair	Not Assessed
30-21	Poor	Not Supporting
20-12	Very Poor	Not Supporting

Table 3. Macroinvertebrate IBI Scores for Reference Sites in the Lake Agassiz Plain Ecoregion of North Dakota.

Site ID	Reference Site Description	IBI Score
551106	Tongue River Below Renwick Dam	72.7
551226	Turtle River Near Emerado, ND	44.6
551231	Pembina River 3.75 miles West of Neche, ND	52.8
551246	Sheyenne River 7.5 miles Southeast of Lisbon, ND	79.8
551248	Sheyenne River 1.5 miles West of Ransom/Richland County Line	88.1

Table 4. Biological Integrity Scoring Criteria and Aquatic Life Use Support Criteria Based on the Lake Agassiz Plain Ecoregion Macroinvertebrate IBI.

IBI Score	Biological Integrity Class	Aquatic Life Use Support
100-53	Good	Fully Supporting
52.99-35.34	Fair	Not Assessed
35.33-17.78	Poor	Not Supporting
17.77-0	Very Poor	Not Supporting

A minimum of two samples which result in beneficial use assessments of fully supporting and/or not supporting are required to assess a waterbody based on biological data (see Section IV. Sufficient and Credible Data Requirements). For assessments based on biological monitoring data, the following use support decision criteria will apply:

- *Fully Supporting:*

Use support assessments for all samples are fully supporting.

- *Fully Supporting, but Threatened:*

Use support assessment for at least one sample is fully supporting, and use support assessments for all other samples are not supporting.

- *Not Supporting:*

Use support assessments for all samples are not supporting.

Biological Assessment Methods for Other Regions in the State

The NDDoH recognizes that there may be biological data that are available for other regions in the state that meet the sufficient and credible data requirements. Where these data are available the NDDoH encourages the use of this information to make aquatic life use support decisions. While it is not possible to assess these sites or waterbodies as fully supporting, sites that are exemplified by low taxa richness, presence of pollutant tolerant taxa and/or low density, can be assessed as not supporting aquatic life use.

B. Recreation Use Assessment Methodology for Rivers and Streams

Recreation use is any activity that relies on water for sport or enjoyment. Recreation use includes primary contact activities such as swimming and wading and secondary contact activities such as boating, fishing, and wading. Recreation use in rivers and streams is considered fully supporting when there is little or no risk of illness through either primary or secondary contact with the water. The state's recreation use support assessment methodology for rivers and streams is based on the state's numeric water quality standards for fecal coliform bacteria and E. coli bacteria (Appendix A).

For each assessment based solely on fecal coliform data, the following criteria are used:

- Assessment Criterion 1a: For each assessment unit, the geometric mean of samples collected during any month from May 1 through September 30 does not exceed a density of 200 colony forming units (CFUs) per 100 milliliters (mL). A minimum of five monthly samples is required to compute the geometric mean. If necessary, samples may be pooled by month across years.
- Assessment Criterion 2a: For each assessment unit, less than 10 percent of samples collected during any month from May 1 through September 30 exceed a density of 400 CFUs per 100 ml. A minimum of five monthly samples is required to compute the percent of samples exceeding the criteria. If necessary, samples may be pooled by month across years.

The two criteria are then applied using the following use support decision criteria:

- Fully Supporting: Both criteria 1a and 2a are met.
- Fully Supporting but Threatened: Criterion 1a is met, but 2a is not.
- Not Supporting: Criterion 1a is not met. Criteria 2a may or may not be met.

For each assessment based solely on E. coli data, the following criteria are used:

- Assessment Criterion 1b: For each assessment unit, the geometric mean of samples collected during any month from May 1 through September 30 does not exceed a density of 126 CFUs per 100 mL. A minimum of five monthly samples is required to compute the geometric mean. If necessary, samples may be pooled by month across years.
- Assessment Criterion 2b: For each assessment unit, less than 10 percent of samples collected during any month from May 1 through September 30 exceed a density of 409 CFUs per 100 ml. A minimum of five monthly samples is required to compute the percent of samples exceeding the criteria. If necessary, samples may be pooled by month across years.

The two criteria are then applied using the following use support decision criteria:

- Fully Supporting: Both criteria 1b and 2b are met.
- Fully Supporting but Threatened: Criterion 1b is met, but 2b is not.
- Not Supporting: Criterion 1b is not met. Criteria 2b may or may not be met.

For each assessment base on both fecal coliform and E. coli data, the following criteria are used:

- Fully Supporting: Criteria 1a, 1b, 2a, and 2b are all met.
- Fully Supporting but Threatened: Criterion 1 a and 1b are both met, but criterion 2a and/or 2b are not.
- Not Supporting: Criterion 1a and/or 1b are not met. Criteria 2a and/or 2b may or may not be met.

C. Aquatic Life and Recreation Use Assessment Methodology for Lakes and Reservoirs

The following is a description of the assessment methodology or decision criteria used to assess aquatic life and recreation uses for lakes and reservoirs in the state. The methodology used to assess the drinking water, fish consumption, agricultural, and industrial uses is the same for both rivers and lakes and is provided in a separate section of the document.

1. Aquatic Life and Recreation

The state's narrative water quality standards (Appendix A) form the basis for aquatic life and recreation use assessment for Section 305(b) reporting and the Section 303(d) TMDL list. State water quality standards contain narrative criteria that require lakes and reservoirs to be "free from" substances "which are toxic or harmful to humans, animals, plants, or resident aquatic biota" or are "in sufficient amounts to be unsightly or deleterious." Narrative standards also prohibit the "discharge of pollutants" (e.g., organic enrichment, nutrients, or sediment), "which alone or in combination with other substances, shall impair existing or reasonable beneficial uses of the receiving waters."

Trophic status is the primary indicator used to assess whether a lake or reservoir is meeting the narrative standards. Trophic status is a measure of the productivity of a lake or reservoir and is directly related to the level of nutrients (i.e., phosphorus and nitrogen) entering the lake or reservoir from its watershed and/or from the internal recycling of nutrients. Highly productive lakes, termed "hypereutrophic," contain excessive phosphorus and are characterized by large growths of weeds, bluegreen algal blooms, low transparency, and low dissolved oxygen (DO) concentrations. These lakes experience frequent fish kills and are generally characterized as having excessive rough fish populations (carp, bullhead, and sucker) and poor sport fisheries. Due to the frequent algal blooms and excessive weed growth, these lakes are also undesirable for recreational uses such as swimming and boating.

Mesotrophic and eutrophic lakes, on the other hand, have lower phosphorus concentrations, low to moderate levels of algae and aquatic plant growth, high transparency, and adequate DO concentrations throughout the year. Mesotrophic lakes do not experience algal blooms, while eutrophic lakes may occasionally experience algal blooms of short duration, typically a few days to a week.

Due to the relationship between trophic status indicators and the aquatic community (as reflected by the fishery) or between trophic status indicators and the frequency of algal blooms, trophic status becomes an effective indicator of aquatic life and recreation use support in lakes and

reservoirs. For purposes of this assessment methodology, it is assumed that hypereutrophic lakes do not fully support a sustainable sport fishery and are limited in recreational uses, whereas mesotrophic lakes fully support both aquatic life and recreation use. Eutrophic lakes may be assessed as fully supporting, fully supporting but threatened, or not supporting their uses for aquatic life or recreation.

Eutrophic lakes are further assessed based on: 1) the lake or reservoir's water quality standards fishery classification; 2) information provided by North Dakota Game and Fish Department Fisheries Division staff, local water resource managers and the public; 3) the knowledge of land use in the lake's watershed; and/or 4) the relative degree of eutrophication. For example, a eutrophic lake, which has a well-balanced sport fishery and experiences infrequent algal blooms, is assessed as fully supporting with respect to aquatic life and recreation use. A eutrophic lake, which experiences periodic algal blooms and limited swimming use, would be assessed as not supporting recreation use. A lake fully supporting its aquatic life and/or recreation use, but for which monitoring has shown a decline in its trophic status (i.e., increasing phosphorus concentrations over time), would be assessed as fully supporting but threatened.

It is recognized that this assessment procedure ignores the fact that, through natural succession, some lakes and reservoirs may display naturally high phosphorus concentrations and experience high productivity. While natural succession or eutrophication can cause high phosphorus concentrations, research suggests that these lakes are typically eutrophic and that lakes classified as hypereutrophic are reflecting external nutrient loading in excess of that occurring naturally.

Since trophic status indicators specific to North Dakota waters have not been developed, Carlson's trophic status index (TSI) (Carlson, 1977) has been chosen to assess the trophic status of lakes or reservoirs. To create a numerical TSI value, Carlson's TSI uses a mathematical relationship based on three indicators: 1) Secchi Disk Transparency in meters (m); 2) surface total phosphorus concentration expressed as $\mu\text{g/L}$; and 3) chlorophyll-a concentration expressed as $\mu\text{g/L}$.

This numerical value, ranging from 0-100, corresponds to a trophic condition with increasing values indicating a more eutrophic (degraded) condition. Carlson's TSI estimates are calculated using the following equations and is also depicted graphically in Figure 2.

- Trophic status based on Secchi Disk Transparency (TSIS):
$$\text{TSIS} = 60 - 14.41 \ln(\text{SD})$$

Where SD = Secchi disk transparency in meters.
- Trophic status based on total phosphorus (TSIP):
$$\text{TSIP} = 14.20 \ln(\text{TP}) + 4.15$$

Where TP = Total phosphorus concentration in $\mu\text{g L}^{-1}$.
- Trophic status based on chlorophyll-a (TSIC):
$$\text{TSIC} = 9.81 \ln(\text{TC}) + 30.60$$

Where TC = Chlorophyll-a concentrations in $\mu\text{g L}^{-1}$.




Figure 2. A Graphic Representation of Carlson's TSI

In general, of the three indicators, it is believed that chlorophyll-a is the best indicator of trophic status, since it is a direct measure of lake productivity. Secchi disk transparency should be used next, followed by phosphorus concentration. In theory, for a given lake or reservoir, the measures of chlorophyll-a, Secchi disk transparency, and phosphorus concentration are all interrelated and should yield similar trophic status index values. This, however, is usually not the case. Many lakes and reservoirs in the state are shallow and windswept causing non-algal turbidity to limit light penetration. This situation may result in a lake having a high phosphorus concentration, low Secchi disk transparency, and low chlorophyll-a concentration. In other instances, other micronutrients may be limiting algal growth even though excessive phosphorus is present.

When conducting an aquatic life and recreation use assessment for a lake or reservoir, the average trophic status index score should be calculated for each indicator. When the trophic status index scores for each indicator (chlorophyll-a, Secchi disk transparency, and phosphorus concentration) each result in a different trophic status assessment then the assessment should be based first on chlorophyll-a, followed by Secchi disk transparency. Only when there are not adequate chlorophyll-a and/or Secchi disk transparency data available to make an assessment should phosphorus concentration data be used.

D. Drinking Water Supply Use Assessment Methodology for Rivers, Lakes, and Reservoirs

Drinking water is defined as “waters that are suitable for use as a source of water supply for drinking and culinary purposes, after treatment to a level approved by the NDDoH” (Appendix A). All Class I, IA, and II rivers and streams and all lakes and reservoirs classified in the state water quality standards (Appendix A), with the exception of Lake George in Kidder County, are assigned the drinking water supply beneficial use. While most lakes and reservoirs are assigned this use, few currently are used as a drinking water supply. Lake Sakakawea is the current

drinking water supply for the Southwest Water Pipeline and the cities of Garrison, Parshall, Pick City, and Riverdale.

Drinking water use is assessed by comparing ambient water quality data to the state water quality standards (Tables 1 and 2 in Appendix A). Ambient water chemistry data are compared to the water quality standards for chloride, sulfate, and nitrate (Table 5) and to the human health standards for Class I, IA, and II rivers and streams (see Table 2 in Appendix A). Drinking water supply is not a designated use for Class III rivers and streams. The human health standard for Class I, IA, and II rivers and streams considers two means of exposure: 1) ingestion of contaminated aquatic organisms; and 2) ingestion of contaminated drinking water.

Drinking water use is also protected through the state's narrative water quality standards. To paraphrase, narrative standards provide language that waters of the state shall be free from materials that produce a color or odor, or other conditions to such a degree as to create a nuisance. Further, state narrative standards provide language that states that waters of the state shall be "free from substances....in concentrations or combinations which are toxic or harmful to **humans**, animals, plants, or resident biota." There shall also be "no discharge of pollutants, whichshall cause a public health hazard or injury to environmental resources."

Table 5. State Water Quality Standards for Chloride, Sulfate, and Nitrate (Appendix A)

Stream Classification	Water Quality Standards (mg/L)		
	Chloride ¹	Sulfate ¹	Nitrate ²
Class I	100	250	10
Class IA	175	450	10
Class II	250	450	10

¹Expressed as a 30-day arithmetic average based on a minimum of four daily samples collected during the 30-day period.

²The water quality standard for nitrite of 1 mg/L shall also not be exceeded.

In order to make beneficial use determinations for drinking water, the following decision criteria are used:

- *Fully Supporting:*

Based on Numeric Standards: No exceedances of the water quality standard for nitrate, one or fewer exceedances of the 30-day average standards for chloride or sulfate, and no exceedances of any of the human health standards.

Based on Narrative Standards: No drinking water complaints on record in the last two years.

- *Fully Supporting but Threatened:*

Based on Numeric Standards: The fully supporting, but threatened use assessment designation is not applied to the drinking water use. Waters are either assessed as fully supporting or not supporting based on chemical data applied to the numeric standards.

Based on Narrative Criteria: No impairment based on the numeric criteria, but a declining trend in water quality over time suggests a measurable increase in the cost to treat water for drinking water supply may occur if the trend continues.

- *Not Supporting:*

Based on Numeric Criteria: One or more exceedances of the water quality standard for nitrate, two or more exceedances of the 30-day average criteria for chloride or sulfate, or one or more exceedances of any of the human health standards.

Based on Narrative Criteria: Knowledge of taste and odor problems or increased treatment costs have been associated with pollutants.

E. Fish Consumption Use Assessment Methodology for Rivers, Lakes and Reservoirs

As stated previously, the state's narrative water quality standards provide that surface waters shall be "free from materials attributable to municipal, industrial, or other discharges or agricultural practices" which will "render any undesirable taste to fish flesh or, in any way, make fish inedible." Fish consumption use is assumed to apply to all Class I, IA, and II rivers and streams, to those Class III streams known to provide a sport fishery and to all Class 1 through 4 lakes and reservoirs.

Fish consumption use is assessed based on EPA guidance. To protect people from exposure to methyl-mercury, EPA recommends a fish tissue-based criterion of 0.3 micrograms (μg) methyl-mercury/gram of fish tissue. This criterion is based on national average consumption rates of fish by recreational users and adjusted for exposures due to consumption of commercial fish. To determine whether the fish tissue criterion of 0.3 $\mu\text{g/g}$ has been exceeded in a lake, reservoir, or river and therefore assessed as not supporting fish consumption, the average fish tissue concentration, weighted by distribution of catch and consumption, is determined for each species in each lake, reservoir, or river for which sufficient and credible data exist.

The weighted average methyl-mercury concentration for each fish species in each lake or river is calculated by multiplying the average methyl-mercury concentration for fish size range by the relative proportion of that size class assumed to be in the creel of fisherman catching and keeping fish from that lake or river. Data to estimate the proportion of each size class in the creel of fisherman were obtained from North Dakota Game and Fish Department creel survey reports and is specific to each lake, reservoir, or river. The weighted-average concentration for each species in each lake or reservoir is then calculated by summing the average concentrations for each size class.

F. Agricultural Use Assessment Methodology for Rivers, Lakes and Reservoirs

Agricultural uses are defined in the state water quality standards as “waters suitable for irrigation, stock watering, and other agricultural uses, but not suitable for use as a source of domestic supply for the farm unless satisfactory treatment is provided.” While not specifically stated in state water quality standards, the numeric standards for pH (6.0-9.0), boron (750 µg/L as a 30-day average), sodium (less than 50% of cation based on mEq/L), and radium (5 pCi/L as a 30-day average) are intended for the protection of agricultural uses. Further, state water quality standards provide for the protection of agricultural uses by providing language that states that waters of the state shall be “free from substances....in concentrations or combinations which are toxic or harmful to humans, **animals, plants**, or resident biota.”

In order to make beneficial use determinations for agricultural uses, the following decision criteria are used:

- *Fully Supporting:*

Based on Numeric Standards: Ten percent or less of the samples exceed the water quality standard for pH or sodium and one or fewer exceedances of the 30-day average criteria for boron or radium.

Based on Narrative Standards: Water supply supports normal crop and livestock production.

- *Fully Supporting but Threatened:*

Based on Numeric Standards: The fully supporting, but threatened use assessment designation is not applied to agricultural use. Waters are either assessed as fully supporting or not supporting based on chemical data applied to the numeric standards.

Based on Narrative Standards: No impairment based on the numeric criteria, but a declining trend in water quality over time suggests a measurable decrease in crop and/or livestock production may occur if the trend continues.

- *Not Supporting:*

Based on Numeric Standards: Greater than 10 percent of samples are exceeded for the water quality standard for pH or sodium, or two or more exceedances of the 30-day average criteria for boron or radium.

Based on Narrative Standards: At least one pollutant has been demonstrated to cause a measurable decrease in crop or livestock production.

G. Industrial Use Assessment Methodology for Rivers, Lakes and Reservoirs

Industrial uses are defined in the state water quality standards as “waters suitable for industrial purposes, including food processing, after treatment.” While there are no specific numeric criteria in the state’s water quality standards intended to protect industrial uses, it is assumed that if the state’s narrative standards are met, or if other numeric water quality standards are met, the beneficial uses for industry will also be met.

VI. REFERENCES

- Barbour, M.T., J. Gerritsen, B.D. Snyder, J.B. Stribling, 1999, *Rapid Bioassessment Protocols for Use in Wadable Streams and Rivers*, EPA 841-B-99-002.
- Carlson, R.E., 1997, *A Trophic Status Index for Lakes*, *Limnology and Oceanography*, Vol. 22 (Issue 2), pp. 361-364.
- EPA, 1998, *Development of Index of Biotic Integrity Expectations for the Lake Agassiz Plain Ecoregion*, U.S. Environmental Protection Agency Region V, Chicago, IL.
- , July 2005, *Guidance for 2006 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d), 305(b) and 314 of the Clean Water Act*, Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans and Watershed, U.S. Environmental Protection Agency, Washington, DC.
- Karr, J.R. and D. R. Dudley, 1981, *Ecological Perspectives on Water Quality Goals*, *Env. Mgmt.* 5:44-68.
- , K.D. Fausch, P.L., Angermeier, P.R. Yant, and I.J. Schlosser, 1986, *Assessing biological integrity in running waters: a method and its rationale*. Illinois Natural History Survey Special Publication 5, 22 pp.
- NDDoH, 2006, *Macroinvertebrate Index of Biotic Integrity for the Lake Agassiz Plain Ecoregion (46) of North Dakota*, North Dakota Department of Health, Bismarck, ND.

Appendix A

Standards of Quality for Waters of the State

Appendix B

Agency and Organization Data Request Letter, Form and Contacts

June 11, 2007

Contact

Dear Agency/Organization:

The Clean Water Act requires states and tribes to monitor and assess the quality of its lakes, reservoirs, rivers, streams and wetlands and to report on the status and condition of its surface waters every two years. The next report, which will be a consolidation of both the Section 305(b) Water Quality Assessment Report and Section 303(d) List of Impaired Waters Needing Total Maximum Daily Loads is due to the US Environmental Protection Agency on April 1, 2008. The North Dakota Department of Health is the primary agency for water quality monitoring and assessment in the state of North Dakota and is therefore responsible for assessing the state's surface waters and preparing the integrated report.

As part of its responsibility, the Department maintains a network of water quality monitoring sites where it collects data on the chemical, physical and biological quality. While these data will be used to provide an assessment of the state's surface water quality, the Department is also requesting additional data that may be used for the 2008 report. If your agency or organization has chemical, physical or biological water quality data that you believe would be beneficial to the state's water quality assessment then please fill out the attached form and return it to me at your earliest convenience.

If you have any questions concerning this request, please contact me at 701.328.5214. Your cooperation in this matter is appreciated.

Sincerely,

Michael J. Ell
Environmental Administrator
Division of Water Quality

Letter Contacts

Jeff Towner
Field Supervisor
North Dakota Field Office
US Fish and Wildlife Service
3425 Miriam Avenue
Bismarck, ND 58501

Dennis Breitzman
Dakotas Area Office
Bureau of Reclamation
P.O. Box 1017
304 East Broadway
Bismarck, ND 58502-1017

Dr. Mark Gozalez
Soil Scientist/Hydrologist
Dakota Prairies Grasslands
US Forest Service
240 West Century Ave
Bismarck, ND 58503

Keith Weston
Water Quality Specialist
Natural Resources Conservation Service
220 East Rosser Avenue
P.O. Box 1458
Bismarck, ND 58502-1458

Terry Steinwand
Director
ND Game and Fish Department
100 North Bismarck Expressway
Bismarck, ND 58501-5095
cc. Greg Power
Steve Dyke

Bob Backman
River Keepers
325 7th Street South
Fargo, ND 58103

Gerald Groenewold
EERC
University of North Dakota
P.O. Box 9018
Grand Forks, ND 58202-9018
cc. Wes Peck

Jim Zeigler
Minnesota Pollution Control Agency
714 Lake Avenue, No. 220
Detroit Lakes MN 56501

Dr. John Watson
School of Engineering and Mines
University of North Dakota
P.O. Box 8155
Grand Forks, ND 58202-8155

Dr. Steven Kelsch
Department of Biology
University of North Dakota
P.O. Box 9019
Grand Forks, ND 58202-8155

Dr. Carolyn E. Grygiel
Natural Resources Management Program Director
Department of Animal and Range Sciences
North Dakota State University
Hultz Hall 163
Fargo, ND 58105

Dr. Frank Yazdani, Chairman
Department Civil Engineering and Construction
North Dakota State University
Civil and Industrial Engineering 201
Fargo, ND 58105

Dr. William Bleier, Chairman
Department of Biological Sciences
North Dakota State University
Stevens Hall, Room 218
Fargo, ND 58105

Edward Murphy
North Dakota Geological Survey
600 East Boulevard Avenue
Bismarck, ND 58505-0840

Greg Wiche
US Geological Survey
821 East Interstate Avenue
Bismarck, ND 58503

Lance Yohe
Red River Basin Commission
119 5th Street South, #209
P.O. Box 66
Moorhead, MN 56561-0066

Col. David C. Press, Commander
US Army Corps of Engineers
Omaha District
106 S. 15th Street
Omaha, NE 68102-1618

Col. Jon L. Christensen
US Army Corps of Engineers
St Paul District
190 5th Street East
St. Paul, MN 55101-1638

Rosie Sada
Montana Department of Environmental Quality
Water Quality Monitoring Section
Metcalf Building Office
1520 E. Sixth Avenue
P.O. Box 200901
Helena, MT 59620

Jim Feeney
Watershed Protection Program
South Dakota Department of Environment and Natural Resources
Joe Foss Building
523 East Capitol Avenue
Pierre, SD 57501-3181

Don Rufledt
Bureau of Land Management
2933 3rd Ave West
Dickinson, ND 58601

Dale Frink
North Dakota State Water Commission
900 East Boulevard Avenue
Bismarck, ND 58505-0850

Water Quality Data Summary for North Dakota

Contact Person: _____

Address: _____

Phone: _____

Email: _____

Data Description: _____

Data Period of Record: _____

Were the data collected according standard operating procedures and/or by following a documented quality assurance/quality control plan?

Yes No Other: _____

Data Availability (e.g., electronic, report): _____

If you have any questions concerning this information, please contact Mike Ell at 701.328.5214

Please return form to: Mike Ell, North Dakota Department of Health, Division of Water Quality,
918 E Divide Ave, 4th Floor, Bismarck, ND 58501-1947

Appendix C

Estimated Weighted Average Methyl-mercury Concentrations in Fish for Lake Sakakawea

Chinook Salmon			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 63	0.173	0.236	0.041
63-72	0.298	0.646	0.192
>73	0.270	0.128	0.035
Weighted Average ⁴			0.268

Northern Pike			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 58	0.12	0.138	0.017
59-77	0.355	0.454	0.161
78-99	0.479	0.408	0.195
>99	0.895	0	0
Weighted Average ⁴			0.373

Sauger			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 37	0.17	0.028	0.005
38-47	0.337	0.873	0.294
>47	0.72	0.099	0.071
Weighted Average ⁴			0.37

¹ Based on the average methyl-mercury concentration for fish sampled in the size range.

² Estimated as the proportion of fish caught and kept by fisherman for that species and waterbody. Based on data obtained from the report entitled *Angler Use and Sport Fishing Catch Survey on Lake Sakakawea, North Dakota - May 1 Through October 24, 2000*, prepared by Larry Brooks and Jeff Hendrickson, submitted to North Dakota Game and Fish Department, Project F-2-R-47, Study 3, Number A-1275, Job C.

³ Calculated by multiplying the average concentration per size range with the weighting factor for the size range.

⁴ Calculated as the sum of the weighted concentrations for each size range.

Walleye			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 40	0.171	0.216	0.037
40-46	0.196	0.411	0.081
47-50	0.389	0.248	0.096
>50	0.508	0.125	0.064
Weighted Average ⁴			0.278

¹ Based on the average methyl-mercury concentration for fish sampled in the size range.

² Estimated as the proportion of fish caught and kept by fisherman for that species and waterbody. Based on data obtained from the report entitled *Angler Use and Sport Fishing Catch Survey on Lake Sakakawea, North Dakota - May 1 Through October 24, 2000*, prepared by Larry Brooks and Jeff Hendrickson, submitted to North Dakota Game and Fish Department, Project F-2-R-47, Study 3, Number A-1275, Job C.

³ Calculated by multiplying the average concentration per size range with the weighting factor for the size range.

⁴ Calculated as the sum of the weighted concentrations for each size range.

Appendix D

Estimated Weighted Average Methyl-mercury Concentrations in Fish for Lake Oahe and the Missouri River

Walleye			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 36	0.15	0.218	0.033
36-39	0.152	0.505	0.077
40-51	0.243	0.264	0.064
>51	0.63	0.013	0.008
Weighted Average ⁴			0.183

¹ Based on the average methyl-mercury concentration for fish sampled in the size range.

² Estimated as the proportion of fish caught and kept by fisherman for that species and waterbody. Based on data obtained from the report entitled *Angler Use and Sport Fishing Catch Survey on Lake Sakakawea, North Dakota - April 1 Through October 15, 2000*, prepared by Larry Brooks and Jeff Hendrickson, submitted to North Dakota Game and Fish Department, Project F-2-R-47, Study 3, Number A-1275, Job B.

³ Calculated by multiplying the average concentration per size range with the weighting factor for the size range.

⁴ Calculated as the sum of the weighted concentrations for each size range.

Appendix E

Estimated Weighted Average Methyl-mercury Concentrations in Fish for Devils Lake

Walleye			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 34	0.43	0.187	0.081
34-40	0.623	0.462	0.288
41-49	0.608	0.249	0.151
50-60	1.248	0.083	0.104
>60	1.79	0.019	0.034
Weighted Average ⁴			0.658

Northern Pike			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 58	0.43	0.11	0.047
59-67	0.569	0.439	0.25
68-77	0.659	0.356	0.235
>77	1.153	0.095	0.11
Weighted Average ⁴			0.642

¹ Based on the average methyl-mercury concentration for fish sampled in the size range.

² Estimated as the proportion of fish caught and kept by fisherman for that species and waterbody. Based on data obtained from the report entitled *Angler Use and Sport Fishing Catch Survey on Lake Sakakawea, North Dakota - May 1 Through October 31, 2001*, prepared by Larry Brooks and Randy Hiltner, submitted to North Dakota Game and Fish Department, Project F-2-R-49, Study 3, Number 2, October 2002.

³ Calculated by multiplying the average concentration per size range with the weighting factor for the size range.

⁴ Calculated as the sum of the weighted concentrations for each size range.

Yellow Perch			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 21	0.27	0.082	0.022
21-25	0.529	0.539	0.285
26-30	0.437	0.333	0.146
>30	0.62	0.046	0.029
Weighted Average ⁴			0.482

White Bass			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 28	0.31	0.061	0.02
28-35	0.54	0.338	0.182
36-41	0.933	0.41	0.382
>41	1.31	0.191	0.25
Weighted Average ⁴			0.834

¹ Based on the average methyl-mercury concentration for fish sampled in the size range.

² Estimated as the proportion of fish caught and kept by fisherman for that species and waterbody. Based on data obtained from the report entitled *Angler Use and Sport Fishing Catch Survey on Lake Sakakawea, North Dakota - May 1 Through October 31, 2001*, prepared by Larry Brooks and Randy Hiltner, submitted to North Dakota Game and Fish Department, Project F-2-R-49, Study 3, Number 2, October 2002.

³ Calculated by multiplying the average concentration per size range with the weighting factor for the size range.

⁴ Calculated as the sum of the weighted concentrations for each size range.

Appendix F

Estimated Weighted Average Methyl-mercury Concentrations in Fish for the Red River of the North

Walleye			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 41	0.74	0.484	0.36
41-63	0.885	0.484	0.428
>63	1.598	0.032	0.051
Weighted Average ⁴			0.839

Channel Catfish			
Size Range (cm)	Average methyl-Hg Concentration ($\mu\text{g/g}$) ¹	Weighting Factor ²	Weighted Concentration ($\mu\text{g/g}$) ³
< 38	0.17	0.276	0.046
38-46	0.287	0.141	0.04
47-56	0.381	0.245	0.093
57-68	0.527	0.252	0.133
>68	0.814	0.086	0.07
Weighted Average ⁴			0.382

¹ Based on the average methyl-mercury concentration for fish sampled in the size range.

² Estimated as the proportion of fish caught and kept by fisherman for that species and waterbody. Based on data obtained from the report entitled *Angler Use and Sport Fishing Catch Survey on Red River, North Dakota - March 15 Through October 31, 2000*, prepared by Larry Brooks and Lynn Schlueter, submitted to North Dakota Game and Fish Department, Project F-2-R-48, Study 3, June 2002.

³ Calculated by multiplying the average concentration per size range with the weighting factor for the size range.

⁴ Calculated as the sum of the weighted concentrations for each size range.

Appendix G

Public Notice Statement Requesting Public Comment on the State of North Dakota's Draft 2008 Section 303(d) List

PUBLIC NOTICE STATEMENT

Notice of submittal to the U.S. Environmental Protection Agency (EPA) and a request for public comment on the State of North Dakota's draft 2008 Section 303(d) List of Waters Needing Total Maximum Daily Loads (TMDLs).

1. Summary

Section 303(d) of the Clean Water Act (CWA) and its accompanying regulations (CFR Part 130 Section 7) requires each state to identify waterbodies (i.e., lakes, reservoirs, rivers, streams, and wetlands) which are considered water quality limited and requiring load allocations, waste load allocations, or total maximum daily loads. A waterbody is considered water quality limited when it is known that its water quality does not meet applicable water quality standards or is not expected to meet applicable water quality standards. Waterbodies can be water quality limited due to point sources of pollution, nonpoint sources of pollution, or both.

Section 303(d) of the Clean Water Act requires states to submit their lists of water quality limited waterbodies "from time to time". Federal regulations have clarified this language, therefore, beginning in 1992 and by April 1st of every even numbered year thereafter, states were required to submit a revised list of waters needing TMDLs. This list has become known as the "TMDL list" or "Section 303(d) list." The state of North Dakota last submitted its TMDL list to EPA in April 2006. This list, referred to as the "2006 list" was approved by EPA on June 27, 2006. The draft 2008 Section 303(d) list, which will be submitted to EPA as part of the integrated Section 305(b) water quality assessment report and Section 303(d) TMDL list, includes a list of waterbodies not meeting water quality standards and which need TMDLs, and a list of waterbodies which have been removed from the list submitted in as part of the 2006 list.

Following an opportunity for public comment, the state must submit its list to the EPA Regional Administrator. The EPA Regional Administrator then has 30 days to either approved or disapprove the state's listings. The purpose of this notice is to solicit public comment on the draft list prior to formally submitting the list to the EPA Regional Administrator.

2. Public Comments

Persons wishing to comment on the State's draft 2008 Section 303(d) List of Waters Needing TMDLs may do so, in writing, within thirty (30) days of the date of this public notice or by July 11, 2008. Comments must be received within this 30-day period to ensure consideration in the EPA approval or disapproval decision. All comments should include the name, address, and telephone number of the person submitting comments, and a statement of the relevant facts upon which they are based. All comments should be submitted to the attention of the Section 303(d) TMDL Coordinator, North Dakota Department of Health, Division of Water Quality, 918 East Divide Avenue, 4th Floor, Bismarck, ND 58501 or by email at mell@state.nd.us. The 2008 Section 303(d) TMDL list may be reviewed at the above address during normal business hours or by accessing it through the Department's web address (<http://www.health.state.nd.us>). Copies may also be requested by writing to the Department at the above address or by calling 701.328.5210.

Appendix H

Public Comments on the State of North Dakota's Draft 2008 Section 303(d) List and the State's Responses



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
REGION 8**

1595 Wynkoop Street
DENVER, CO 80202-1129
Phone 800-227-8917
<http://www.epa.gov/region08>

June 30, 2008

Ref: 8EPR-EP

Mike Ell
Division of Water Quality
North Dakota Department of Health
918 East Divide Ave., 4th Floor
Bismarck, ND 58501-1947

RE: EPA Comments on North Dakota's Draft 2008 Integrated Report (IR)

Dear Mr. Ell:

Thank you for the opportunity to comment on North Dakota's Draft Integrated Report (IR). Detailed comments are enclosed. A goal of the 2008 IR cycle is to achieve 100 percent on-time submittals of the Integrated Reports by April 1, 2008. To assist in attaining this goal, EPA recommends a series of best practices that have been used by States and EPA regions to meet previous IR deadlines. Timely submittal and EPA review of integrated reports is a key to demonstrating State and EPA success in accomplishing mutual goals for restoring and maintaining the nation's waters.

EPA is aware of the challenge the North Dakota Department of Health faces in terms of sufficient staffing to address jobs like compiling the draft report and releasing it for public comment. We recommend the October 12, 2006 memorandum *Information Concerning 2008 Clean Water Act Sections 303(d), 305(b), and 314 Integrated Reporting and Listing Decisions*, as a good starting point for reviewing best practices to increase timeliness. It is available here: <http://epa.gov/owow/tmdl/guidance.html#2>. We also note in places in the draft IR the most recent EPA integrated reporting guidance cited is from 2006.

Please contact me (303-312-6237) if you have any questions with regard to our comments.

Sincerely,

Kris Jensen
Water Quality Unit
Ecosystems Protection Program

Enclosure

Cc: Karen Hamilton, Vern Berry, Dave Moon, Julie Kinsey, EPA

Detailed Comments on North Dakota's 2008 Draft Integrated Report (IR)

Congratulations to the State for completing these efforts in this listing cycle:

- A. **Environmental Monitoring and Assessment Program – Western Pilot Project (EMAP – West)**
EPA commends North Dakota staff for including the entire report of the western pilot project results for North Dakota in their 2008 Integrated Report.
- B. **Assessment Methodology**
EPA appreciates the State's efforts to address the following issues in the assessment methodology for the 2008 listing cycle:
 1. Human Health Criteria
 2. Sufficient and Credible Data Requirements and Overwhelming Evidence

Comments:

1. **Comments Related to Categories 1-4:**
Review of Category 4 waters (i.e., from ADB files): Category 4a = 31 TMDLs; Category 4b = 0; Category 4c = 16 listings (13 physical substrate habitat alterations; 2 other flow regime alterations; 1 low flow alterations). Vern Berry, ND's EPA TMDL Coordinator, has reviewed all of these listings and concurs with North Dakota's categorization.
 - All of the 4a listings are consistent with EPA's TMDL approvals; and
 - All of the 4c listings appear to be for non-pollutants.
2. **Comments Related to Category 5:**
 - a. **General comment:** EPA notes many of the waterbody listings have significant mileage or acreage changes in **Table VI-3**. Please explain/correct this discrepancy.
 - b. **Table VI-5 – “2006 Section 303(d) TMDL Waters in the State That Have Been De-listed for 2008”:** EPA concurs with the waters delisted as the result of a TMDL being approved, including the sediment delistings.
 - c. **Chapter 4, p. III-28, third paragraph:** The three-tiered, numerical priority ranking system described here appears inconsistent with the two-tiered, narrative (i.e., high, low) priority ranking described elsewhere in the report and reflected in the 303(d) listing tables. Please advise or revise.
 - d. **Assessment Unit Name and Description Changes:** EPA's review of Category 5 waters, Table VI-3, has revealed several/many name and description changes of

assessment units. Please clarify and/or correct these changes.

- e. **Pollutant Name Changes/Modifications:** EPA also notes and requires confirmation from North Dakota that each of the changes to the pollutant listings (total fecal coliform => fecal coliform; organic enrichment/oxygen, dissolved => oxygen, dissolved; biological indicators => combination benthic/fishes bioassessments; etc...) was made as a refinement an existing waterbody/pollutant listing.
- f. According to EPA's review the following waterbody/pollutants from the 2006 303(d) list were missing without explanation in 2008:
- English Coulee (ND- 09020301-002S_00) – organic enrichment;
 - Turtle River (ND-09020307-001-S_00) – total dissolved solids;
 - Turtle River (ND-09020307-006-S_00) – total dissolved solids.
- g. The final TMDL approval date for Sheep Creek Dam was May 28, 2008. Please revise the ADB files and 303(d) delisting table, p. VI-55.



**North Dakota Department of Health
Response to Comments on the Draft 2008 Integrated Report**

US Environmental Protection Agency, Region 8

Comment:

EPA notes many of the waterbody listings have significant mileage or acreage changes in **Table VI-3**. Please explain/correct this discrepancy.

Department Response:

River and stream mileage estimates described in previous Section 303(d) TMDL lists were based on “reach indexing” and segment length estimates provided in Reach File 3 (RF3). In preparing for the 2008 Integrated Report, mileage estimates for many assessment units (AUs) were recalculated (using GIS) using the more accurate 1:100,00 scale National Hydrography Dataset (NHD). To accomplish this task, AUs were ‘reach indexed’ to the NHD and stream miles recalculated. In some cases the change in stream length mileage was also the result of corrections in the original reach indexing, where stream segments in the NHD were included or excluded in error. This was also the case with changes in some lake and reservoir acreage estimates. In most of the cases where lake and reservoir acreage estimates changed, the difference was due to new mapping information provided by the ND Game and Fish Department. In a few instances the change in lake acreage was due to either increasing or decreasing water elevation, thereby changing the lakes surface area.

It should also be noted that for the first time the Section 303(d) list of impaired waters needing TMDLs was generated directly from the Assessment Database (ADB) rather than from a table where the listing information was hand entered. By generating the TMDL list directly from the ADB, the Department and EPA are assured that there is complete harmonization and consistency between the two.

Comment:

Table VI-5 – “2006 Section 303(d) TMDL Waters in the State That Have Been De-listed for 2008”: EPA concurs with the waters delisted as the result of a TMDL being approved, including the sediment delistings.

Department Response:

In a telephone conference call with Vern Berry and Kris Jensen, EPA Region 8, on July 23, 2008 clarification was requested on whether this statement also meant that all of the de-listed waterbodies provided in Table VI-5 were approved by EPA Region 8. With the exception of Antelope Creek, ND-09020105-005-S_00, it was agreed that the de-listing rationale provided by the Department for all of the waterbody stream segments, lakes and reservoirs de-listed in Table VI-5 were acceptable to EPA.

The Department has reviewed the temperature de-listing for Antelope Creek, ND-09020105-005-S_00, and based on existing data and the Department’s Assessment Methodology has determined that there is no evidence for the temperature de-listing. The Department’s has restored the

temperature impairment listing to Table VI-2 and removed the de-listing from Table VI-5. Assessment data in the Assessment Database (ADB) has also been updated to reflect this change.

Comment:

Chapter 4, p. III-28, third paragraph: The three-tiered, numerical priority ranking system described here appears inconsistent with the two-tiered, narrative (i.e., high, low) priority ranking described elsewhere in the report and reflected in the 303(d) listing tables. Please advise or revise.

Department Response:

The three-tiered, numerical priority ranking system for TMDL development described in Part III, Chapter 4, reflects the previously used system. The priority ranking system that has been put in place for 2008 TMDL listed waters is described in Part VI, Section F. The narrative provided in Part III, Chapter 4, has been revised to be consistent with the narrative provided in Part VI, Section F.

Comment:

EPA's review of Category 5 waters, Table VI-3, has revealed several/many name and description changes of assessment units. Please clarify and/or correct these changes.

Department Response:

As was stated in the Department's Response to the first comment (see above), the 2008 Integrated Report represents the first time that the Section 303(d) list of impaired waters needing TMDLs was generated directly from the Assessment Database (ADB) rather than from a table where the listing information was hand entered. This "hand entered" table was first generated for the 2002 Section 303(d) listing cycle and was edited by adding and deleting waterbodies by hand for the 2004 and 2006 cycles. As a result, waterbody names and descriptions added, changed or modified in the ADB were not reflected in the 2002, 2004 or 2006 lists. Therefore, while the actual assessment unit ID remains the same from the 2006 cycle to the 2008 cycle, the waterbody name, description and/or size has change to reflect the name, description and size as it appears in the ADB. As was also stated previously, by generating the TMDL list directly from the ADB, the Department and EPA are assured that there is complete harmonization and consistency between the two.

Comment:

EPA also notes and requires confirmation from North Dakota that each of the changes to the pollutant listings (total fecal coliform => fecal coliform; organic enrichment/oxygen, dissolved => oxygen, dissolved; biological indicators => combination benthic/fishes bioassessments; etc...) was made as a refinement (of) an existing waterbody/pollutant listing.

Department Response:

The Department acknowledges that the pollutant listings in the 2008 TMDL list are a refinement of previous pollutant listings. Changes to these pollutant listings were made to reflect the more accurate pollutant descriptions provided in the Assessment Database (ADB). As was stated in previous responses to comments, the 2008 Integrated Report represents the first time that the

Section 303(d) list of impaired waters needing TMDLs was generated directly from the Assessment Database (ADB) rather than from a table where the listing information was hand entered. Listings in the “hand entered” tables include pollutant descriptions such as “total fecal coliform”, “organic enrichment/oxygen, dissolved”, and “biological indicators”. The ADB provides a pick list with a prescribed set of pollutant or cause categories from which to pick. Pollutant categories such as “total fecal coliform”, “organic enrichment/oxygen, dissolved” and “biological indicators” are not among the pollutant listings provided in the ADB pick list. Rather, the ADB user must choose “fecal coliform” instead of “total fecal coliform”, “dissolved oxygen” instead of “organic enrichment/oxygen, dissolved” and “combination benthic/fishes bioassessments” instead of “biological indicators.” It should be noted that the 2008 list also includes “benthic-macroinvertebrate bioassessments” and “fishes bioassessments” as a refinement of the “biological indicators” listing. These refined “bioassessment” pollutant cause categories simply reflects the types of biological data by which the assessment is based.

The Department also acknowledges that for future TMDL listing cycles, any further changes to pollutant cause categories (unless they are the result of changes EPA makes to the ADB pollutant cause categories) will only be made through the de-listing and listing process.

Comment:

According to EPA’s review the following waterbody/pollutants from the 2006 303(d) list were missing without explanation in 2008:

- English Coulee (ND- 09020301-002-S_00) – organic enrichment;
- Turtle River (ND-09020307-001-S_00) – total dissolved solids;
- Turtle River (ND-09020307-006-S_00) – total dissolved solids.

Department Response:

The waterbody/pollutant listing for English Coulee (ND-09020301-002-S) and organic enrichment is on the 2008 TMDL list, however the pollutant listing was changed to dissolved oxygen (see previous response to comments). As stated previously, the ADB does not provide for a pollutant cause called “organic enrichment” as was provided in the previous 2006 TMDL list. Since low dissolved oxygen is the result of organic enrichment, the category “dissolved oxygen” was selected in the ADB.

Both Turtle River segments (ND-09020307-001-S and ND09020307-006-S) listed for TDS were included in Table V-5 of the 2008 Integrated Report entitled “2006 Section 303(d) TMDL Waters in the State Which Have Been De-listed for 2008.” Justification for these two de-listings were *“Data and/or information lacking to determine water quality status; original basis for listing was incorrect (Category 3). Based on the water quality assessment that was conducted by the Grand Forks County SCD, there is no basis for the listing of the aquatic life impairment due to TDS. There is no existing water quality standard for TDS and therefore, no basis for the original assessment and TMDL listing.”*

Comment:

The final TMDL approval date for Sheep Creek Dam was May 28, 2008. Please revise the ADB files and 303(d) delisting table, p. VI-55.

Department Response:

The final TMDL approval date for Sheep Creek Dam has been changed in both the ADB and the de-listing table to reflect the actual approval date of May 28, 2008.